

50+51

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C12N 15/56, 15/63, 1/21, 9/24, 15/11</b>		<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/52178</b>
			<b>(43) International Publication Date:</b> 8 September 2000 (08.09.00)
<b>(21) International Application Number:</b> PCT/US00/03542		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
<b>(22) International Filing Date:</b> 14 February 2000 (14.02.00)			
<b>(30) Priority Data:</b> 09/258,892 1 March 1999 (01.03.99) US			
<b>(71) Applicants (for all designated States except US):</b> INSIGHT STRATEGY & MARKETING LTD. [IL/IL]; P.O. Box 2128, 76121 Rehovot (IL). HADASIT MEDICAL RESEARCH SERVICES & DEVELOPMENT LTD. [IL/IL]; P.O. Box 12000, 91120 Jerusalem (IL).			
<b>(71) Applicant (for TJ only):</b> FRIEDMAN, Mark, M. [US/IL]; 1 Alharizi Street, 43406 Raanana (IL).		<b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
<b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> PECKER, Iris [IL/IL]; 42 Wolfson Street, 75203 Rishon Lezion (IL). VLODAVSKY, Israel [IL/IL]; 34 Arbel Street, 90805 Mevaseret Zion (IL). FEINSTEIN, Elena [IL/IL]; 12/29 Hahagana Street, 76214 Rehovot (IL).			
<b>(74) Common Representative:</b> FRIEDMAN, Mark, M.; c/o Castorina, Anthony, Suite 207, 2001 Jefferson Davis Highway, Arlington, VA 22202 (US).			
<b>(54) Title:</b> POLYNUCLEOTIDE ENCODING A POLYPEPTIDE HAVING HEPARANASE ACTIVITY AND EXPRESSION OF SAME IN GENETICALLY MODIFIED CELLS			
<b>(57) Abstract</b>  A polynucleotide ( <i>hpa</i> ) encoding a polypeptide having heparanase activity, vectors including same, genetically modified cells expressing heparanase, a recombinant protein having heparanase activity and antisense oligonucleotides and constructs for modulating heparanase expression.			

52

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakistan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

POLYNUCLEOTIDE ENCODING A POLYPEPTIDE HAVING  
HEPARANASE ACTIVITY AND EXPRESSION OF SAME IN  
GENETICALLY MODIFIED CELLS

5 FIELD AND BACKGROUND OF THE INVENTION

The present invention relates to a polynucleotide, referred to hereinbelow as *hpa*, encoding a polypeptide having heparanase activity, vectors (nucleic acid constructs) including same and genetically modified cells expressing heparanase. The invention further relates to a recombinant  
10 protein having heparanase activity and to antisense oligonucleotides, constructs and ribozymes for down regulating heparanase activity. In addition, the invention relates to heparanase promoter sequences and their uses.

*Heparan sulfate proteoglycans:* Heparan sulfate proteoglycans (HSPG) are ubiquitous macromolecules associated with the cell surface and extra cellular matrix (ECM) of a wide range of cells of vertebrate and invertebrate tissues (1-4). The basic HSPG structure includes a protein core to which several linear heparan sulfate chains are covalently attached. These polysaccharide chains are typically composed of repeating hexuronic  
15 and D-glucosamine disaccharide units that are substituted to a varying extent with N- and O-linked sulfate moieties and N-linked acetyl groups (1-4). Studies on the involvement of ECM molecules in cell attachment, growth and differentiation revealed a central role of HSPG in embryonic morphogenesis, angiogenesis, neurite outgrowth and tissue repair (1-5).  
20 HSPG are prominent components of blood vessels (3). In large blood vessels they are concentrated mostly in the intima and inner media, whereas in capillaries they are found mainly in the subendothelial basement membrane where they support proliferating and migrating endothelial cells and stabilize the structure of the capillary wall. The ability of HSPG to  
25 interact with ECM macromolecules such as collagen, laminin and fibronectin, and with different attachment sites on plasma membranes suggests a key role for this proteoglycan in the self-assembly and insolubility of ECM components, as well as in cell adhesion and locomotion. Cleavage of the heparan sulfate (HS) chains may therefore  
30 result in degradation of the subendothelial ECM and hence may play a decisive role in extravasation of blood-borne cells. HS catabolism is observed in inflammation, wound repair, diabetes, and cancer metastasis, suggesting that enzymes which degrade HS play important roles in pathologic processes. Heparanase activity has been described in activated

immune system cells and highly metastatic cancer cells (6-8), but research has been handicapped by the lack of biologic tools to explore potential causative roles of heparanase in disease conditions.

***Involvement of Heparanase in Tumor Cell Invasion and***  
5 ***Metastasis:*** Circulating tumor cells arrested in the capillary beds of different organs must invade the endothelial cell lining and degrade its underlying basement membrane (BM) in order to invade into the extravascular tissue(s) where they establish metastasis (9, 10). Metastatic  
10 tumor cells often attach at or near the intercellular junctions between adjacent endothelial cells. Such attachment of the metastatic cells is followed by rupture of the junctions, retraction of the endothelial cell borders and migration through the breach in the endothelium toward the exposed underlying BM (9). Once located between endothelial cells and the BM, the invading cells must degrade the subendothelial glycoproteins and  
15 proteoglycans of the BM in order to migrate out of the vascular compartment. Several cellular enzymes (e.g., collagenase IV, plasminogen activator, cathepsin B, elastase, etc.) are thought to be involved in degradation of BM (10). Among these enzymes is an endo- $\beta$ -D-glucuronidase (heparanase) that cleaves HS at specific intrachain sites (6,  
20 8, 11). Expression of a HS degrading heparanase was found to correlate with the metastatic potential of mouse lymphoma (11), fibrosarcoma and melanoma (8) cells. Moreover, elevated levels of heparanase were detected in sera from metastatic tumor bearing animals and melanoma patients (8) and in tumor biopsies of cancer patients (12).

25 The control of cell proliferation and tumor progression by the local microenvironment, focusing on the interaction of cells with the extracellular matrix (ECM) produced by cultured corneal and vascular endothelial cells, was investigated previously by the present inventors. This cultured ECM closely resembles the subendothelium *in vivo* in its  
30 morphological appearance and molecular composition. It contains collagens (mostly type III and IV, with smaller amounts of types I and V), proteoglycans (mostly heparan sulfate- and dermatan sulfate- proteoglycans, with smaller amounts of chondroitin sulfate proteoglycans), laminin, fibronectin, entactin and elastin (13, 14). The ability of cells to degrade HS  
35 in the cultured ECM was studied by allowing cells to interact with a metabolically sulfate labeled ECM, followed by gel filtration (Sephacrose 6B) analysis of degradation products released into the culture medium (11). While intact HSPG are eluted next to the void volume of the column



( $K_{av} < 0.2$ ,  $M_r \sim 0.5 \times 10^6$ ), labeled degradation fragments of HS side chains are eluted more toward the  $V_t$  of the column ( $0.5 < k_{av} < 0.8$ ,  $M_r = 5-7 \times 10^3$ ) (11).

The heparanase inhibitory effect of various non-anticoagulant species of heparin that might be of potential use in preventing extravasation of blood-borne cells was also investigated by the present inventors. Inhibition of heparanase was best achieved by heparin species containing 16 sugar units or more and having sulfate groups at both the N and O positions. While O-desulfation abolished the heparanase inhibiting effect of heparin, O-sulfated, N-acetylated heparin retained a high inhibitory activity, provided that the N-substituted molecules had a molecular size of about 4,000 daltons or more (7). Treatment of experimental animals with heparanase inhibitors (e.g., non-anticoagulant species of heparin) markedly reduced (>90%) the incidence of lung metastases induced by B16 melanoma, Lewis lung carcinoma and mammary adenocarcinoma cells (7, 8, 16). Heparin fractions with high and low affinity to anti-thrombin III exhibited a comparable high anti-metastatic activity, indicating that the heparanase inhibiting activity of heparin, rather than its anticoagulant activity, plays a role in the anti-metastatic properties of the polysaccharide (7).

**Heparanase activity in the urine of cancer patients:** In an attempt to further elucidate the involvement of heparanase in tumor progression and its relevance to human cancer, urine samples for heparanase activity were screened (16a). Heparanase activity was detected in the urine of some, but not all, cancer patients. High levels of heparanase activity were determined in the urine of patients with an aggressive metastatic disease and there was no detectable activity in the urine of healthy donors.

Heparanase activity was also found in the urine of 20% of normal and microalbuminuric insulin dependent diabetes mellitus (IDDM) patients, most likely due to diabetic nephropathy, the most important single disorder leading to renal failure in adults.

**Possible involvement of heparanase in tumor angiogenesis:** Fibroblast growth factors are a family of structurally related polypeptides characterized by high affinity to heparin (17). They are highly mitogenic for vascular endothelial cells and are among the most potent inducers of neovascularization (17, 18). Basic fibroblast growth factor (bFGF) has been extracted from the subendothelial ECM produced *in vitro* (19) and from basement membranes of the cornea (20), suggesting that ECM may serve as

a reservoir for bFGF. Immunohistochemical staining revealed the localization of bFGF in basement membranes of diverse tissues and blood vessels (21). Despite the ubiquitous presence of bFGF in normal tissues, endothelial cell proliferation in these tissues is usually very low, suggesting that bFGF is somehow sequestered from its site of action. Studies on the interaction of bFGF with ECM revealed that bFGF binds to HSPG in the ECM and can be released in an active form by HS degrading enzymes (15, 20, 22). It was demonstrated that heparanase activity expressed by platelets, mast cells, neutrophils, and lymphoma cells is involved in release of active bFGF from ECM and basement membranes (23), suggesting that heparanase activity may not only function in cell migration and invasion, but may also elicit an indirect neovascular response. These results suggest that the ECM HSPG provides a natural storage depot for bFGF and possibly other heparin-binding growth promoting factors (24, 25). Displacement of bFGF from its storage within basement membranes and ECM may therefore provide a novel mechanism for induction of neovascularization in normal and pathological situations.

Recent studies indicate that heparin and HS are involved in binding of bFGF to high affinity cell surface receptors and in bFGF cell signaling (26, 27). Moreover, the size of HS required for optimal effect was similar to that of HS fragments released by heparanase (28). Similar results were obtained with vascular endothelial cells growth factor (VEGF) (29), suggesting the operation of a dual receptor mechanism involving HS in cell interaction with heparin-binding growth factors. It is therefore proposed that restriction of endothelial cell growth factors in ECM prevents their systemic action on the vascular endothelium, thus maintaining a very low rate of endothelial cells turnover and vessel growth. On the other hand, release of bFGF from storage in ECM as a complex with HS fragment, may elicit localized endothelial cell proliferation and neovascularization in processes such as wound healing, inflammation and tumor development (24, 25).

***Expression of heparanase by cells of the immune system:***  
Heparanase activity correlates with the ability of activated cells of the immune system to leave the circulation and elicit both inflammatory and autoimmune responses. Interaction of platelets, granulocytes, T and B lymphocytes, macrophages and mast cells with the subendothelial ECM is associated with degradation of HS by a specific heparanase activity (6). The enzyme is released from intracellular compartments (e.g., lysosomes,

specific granules, etc.) in response to various activation signals (e.g., thrombin, calcium ionophore, immune complexes, antigens, mitogens, etc.), suggesting its regulated involvement in inflammation and cellular immunity.

5        ***Some of the observations regarding the heparanase enzyme were reviewed in reference No. 6 and are listed hereinbelow:***

First, a proteolytic activity (plasminogen activator) and heparanase participate synergistically in sequential degradation of the ECM HSPG by inflammatory leukocytes and malignant cells.

10        Second, a large proportion of the platelet heparanase exists in a latent form, probably as a complex with chondroitin sulfate. The latent enzyme is activated by tumor cell-derived factor(s) and may then facilitate cell invasion through the vascular endothelium in the process of tumor metastasis.

15        Third, release of the platelet heparanase from  $\alpha$ -granules is induced by a strong stimulant (i.e., thrombin), but not in response to platelet activation on ECM.

Fourth, the neutrophil heparanase is preferentially and readily released in response to a threshold activation and upon incubation of the  
20        cells on ECM.

Fifth, contact of neutrophils with ECM inhibited release of noxious enzymes (proteases, lysozyme) and oxygen radicals, but not of enzymes (heparanase, gelatinase) which may enable diapedesis. This protective role of the subendothelial ECM was observed when the cells were stimulated  
25        with soluble factors but not with phagocytosable stimulants.

Sixth, intracellular heparanase is secreted within minutes after exposure of T cell lines to specific antigens.

Seventh, mitogens (Con A, LPS) induce synthesis and secretion of heparanase by normal T and B lymphocytes maintained *in vitro*. T  
30        lymphocyte heparanase is also induced by immunization with antigen *in vivo*.

Eighth, heparanase activity is expressed by pre-B lymphomas and B-lymphomas, but not by plasmacytomas and resting normal B lymphocytes.

Ninth, heparanase activity is expressed by activated macrophages  
35        during incubation with ECM, but there was little or no release of the enzyme into the incubation medium. Similar results were obtained with human myeloid leukemia cells induced to differentiate to mature macrophages.

Tenth, T-cell mediated delayed type hypersensitivity and experimental autoimmunity are suppressed by low doses of heparanase inhibiting non-anticoagulant species of heparin (30).

Eleventh, heparanase activity expressed by platelets, neutrophils and metastatic tumor cells releases active bFGF from ECM and basement membranes. Release of bFGF from storage in ECM may elicit a localized neovascular response in processes such as wound healing, inflammation and tumor development.

Twelfth, among the breakdown products of the ECM generated by heparanase is a tri-sulfated disaccharide that can inhibit T-cell mediated inflammation *in vivo* (31). This inhibition was associated with an inhibitory effect of the disaccharide on the production of biologically active TNF $\alpha$  by activated T cells *in vitro* (31).

**Other potential therapeutic applications:** Apart from its involvement in tumor cell metastasis, inflammation and autoimmunity, mammalian heparanase may be applied to modulate: bioavailability of heparin-binding growth factors (15); cellular responses to heparin-binding growth factors (e.g., bFGF, VEGF) and cytokines (IL-8) (31a, 29); cell interaction with plasma lipoproteins (32); cellular susceptibility to certain viral and some bacterial and protozoa infections (33, 33a, 33b); and disintegration of amyloid plaques (34). Heparanase may thus prove useful for conditions such as wound healing, angiogenesis, restenosis, atherosclerosis, inflammation, neurodegenerative diseases and viral infections. Mammalian heparanase can be used to neutralize plasma heparin, as a potential replacement of protamine. Anti-heparanase antibodies may be applied for immunodetection and diagnosis of micrometastases, autoimmune lesions and renal failure in biopsy specimens, plasma samples, and body fluids. Common use in basic research is expected.

The identification of the *hpa* gene encoding for heparanase enzyme will enable the production of a recombinant enzyme in heterologous expression systems. Availability of the recombinant protein will pave the way for solving the protein structure function relationship and will provide a tool for developing new inhibitors.

**Viral Infection:** The presence of heparan sulfate on cell surfaces have been shown to be the principal requirement for the binding of Herpes Simplex (33) and Dengue (33a) viruses to cells and for subsequent infection of the cells. Removal of the cell surface heparan sulfate by heparanase may

therefore abolish virus infection. In fact, treatment of cells with bacterial heparitinase (degrading heparan sulfate) or heparinase (degrading heparan) reduced the binding of two related animal herpes viruses to cells and rendered the cells at least partially resistant to virus infection (33). There are some indications that the cell surface heparan sulfate is also involved in HIV infection (33b).

**Neurodegenerative diseases:** Heparan sulfate proteoglycans were identified in the prion protein amyloid plaques of Genstmann-Straussler Syndrome, Creutzfeldt-Jakob disease and Scrape (34). Heparanase may disintegrate these amyloid plaques which are also thought to play a role in the pathogenesis of Alzheimer's disease.

**Restenosis and Atherosclerosis:** Proliferation of arterial smooth muscle cells (SMCs) in response to endothelial injury and accumulation of cholesterol rich lipoproteins are basic events in the pathogenesis of atherosclerosis and restenosis (35). Apart from its involvement in SMC proliferation (i.e., low affinity receptors for heparin-binding growth factors), HS is also involved in lipoprotein binding, retention and uptake (36). It was demonstrated that HSPG and lipoprotein lipase participate in a novel catabolic pathway that may allow substantial cellular and interstitial accumulation of cholesterol rich lipoproteins (32). The latter pathway is expected to be highly atherogenic by promoting accumulation of apoB and apoE rich lipoproteins (i.e. LDL, VLDL, chylomicrons), independent of feed back inhibition by the cellular sterol content. Removal of SMC HS by heparanase is therefore expected to inhibit both SMC proliferation and lipid accumulation and thus may halt the progression of restenosis and atherosclerosis.

#### **Gene therapy:**

The ultimate goal in the management of inherited as well as acquired diseases is a rational therapy with the aim to eliminate the underlying biochemical defects associated with the disease rather than symptomatic treatment. Gene therapy is a promising candidate to meet these objectives. Initially it was developed for treatment of genetic disorders, however, the consensus view today is that it offers the prospect of providing therapy for a variety of acquired diseases, including cancer, viral infections, vascular diseases and neurodegenerative disorders.

The gene-based therapeutic can act either intracellularly, affecting only the cells to which it is delivered, or extracellularly, using the recipient cells as local endogenous factories for the therapeutic product(s). The

application of gene therapy may follow any of the following strategies: (i) prophylactic gene therapy, such as using gene transfer to protect cells against viral infection; (ii) cytotoxic gene therapy, such as cancer therapy, where genes encode cytotoxic products to render the target cells vulnerable to attack by the normal immune response; (iii) biochemical correction, primarily for the treatment of single gene defects, where a normal copy of the gene is added to the affected or other cells.

To allow efficient transfer of the therapeutic genes, a variety of gene delivery techniques have been developed based on viral and non-viral vector systems. The most widely used and most efficient systems for delivering genetic material into target cells are viral vectors. So far, 329 clinical studies (phase I, I/II and II) with over 2,500 patients have been initiated Worldwide since 1989 (50).

The approach of gene addition pose serious barriers. The expression of many genes is tightly regulated and context dependent, so achieving the correct balance and function of expression is challenging. The gene itself is often quite large, containing many exons and introns. The delivery vector is usually a virus, which can infect with a high efficiency but may, on the other hand, induce immunological response and consequently decreases effectiveness, especially upon secondary administration. Most of the current expression vector-based gene therapy protocols fail to achieve clinically significant transgene expression required for treating genetic diseases. Apparently, it is difficult to deliver enough virus to the right cell type to elicit an effective and therapeutic effect (51)

Homologous recombination, which was initially considered to be of limited use for gene therapy because of its low frequency in mammalian cells, has recently emerged as a potential strategy for developing gene therapy. Different approaches have been used to study homologous recombination in mammalian cells; some involve DNA repair mechanisms. These studies aimed at either gene disruption or gene correction and include RNA/DNA chimeric oligonucleotides, small or large homologous DNA fragments, or adeno-associated viral vectors. Most of these studies show a reasonable frequency of homologous recombination, which warrants further *in vivo* testing (52). Homologous recombination-based gene therapy has the potential to develop into a powerful therapeutic modality for genetic diseases. It can offer permanent expression and normal regulation of corrected genes in appropriate cells or organs and probably can be used for treating dominantly inherited diseases such as polycystic kidney disease.

***Genomic sequences function in regulation of gene expression:***

The efficient expression of therapeutic genes in target cells or tissues is an important component of efficient and safe gene therapy. The expression of genes is driven by the promoter region upstream of the coding sequence, although regulation of expression may be supplemented by  
5 farther upstream or downstream DNA sequences or DNA in the introns of the gene. Since this important information is embedded in the DNA, the description of gene structure is crucial to the analysis of gene regulation. Characterization of cell specific or tissue specific promoters, as well as  
10 other tissue specific regulatory elements enables the use of such sequences to direct efficient cell specific, or developmental stage specific gene expression. This information provides the basis for targeting individual genes and for control of their expression by exogenous agents, such as drugs. Identification of transcription factors and other regulatory proteins  
15 required for proper gene expression will point at new potential targets for modulating gene expression, when so desired or required.

Efficient expression of many mammalian genes depends on the presence of at least one intron. The expression of mouse thymidylate synthase (TS) gene, for example, is greatly influenced by intron sequences.  
20 The addition of almost any of the introns from the mouse TS gene to an intronless TS minigene leads to a large increase in expression (42). The involvement of intron 1 in the regulation of expression was demonstrated for many other genes. In human factor IX (hFIX), intron 1 is able to increase the expression level about 3 fold more as compared to that of the  
25 hFIX cDNA (43). The expression enhancing activity of intron 1 is due to efficient functional splicing sequences, present in the precursor mRNA. By being efficiently assembled into spliceosome complexes, transcripts with splicing sequences may be better protected in the nucleus from random degradations, than those without such sequences (44).

30 A forward-inserted intron1-carrying hFIX expression cassette suggested to be useful for directed gene transfer, while for retroviral-mediated gene transfer system, reversely-inserted intron 1-carrying hFIX expression cassette was considered (43).

A highly conserved cis-acting sequence element was identified in the  
35 first intron of the mouse and rat c-Ha-ras, and in the first exon of Ha- and Ki-ras genes of human, mouse and rat. This cis-acting regulatory sequence confers strong transcription enhancer activity that is differentially modulated by steroid hormones in metastatic and nonmetastatic

subpopulations. Perturbations in the regulatory activities of such cis-acting sequences may play an important role in governing oncogenic potency of Ha-ras through transcriptional control mechanisms (45).

Intron sequences affect tissue specific, as well as inducible gene expression. A 182 bp intron 1 DNA segment of the mouse Col2a1 gene contains the necessary information to confer high-level, temporally correct, chondrocyte expression on a reporter gene in intact mouse embryos, while Col2a1 promoter sequences are dispensable for chondrocyte expression (46). In Col1A1 gene the intron plays little or no role in constitutive expression of collagen in the skin, and in cultured cells derived from the skin, however, in the lungs of young mice, intron deletion results in decrease of expression to less than 50 % (47).

A classical enhancer activity was shown in the 2 kb intron fragment in bovine beta-casein gene. The enhancer activity was largely dependent on the lactogenic hormones, especially prolactin. It was suggested that several elements in the intron-1 of the bovine beta-casein gene cooperatively interact not only with each other but also with its promoter for hormonal induction (48).

Identification and characterization of regulatory elements in genomic non-coding sequences, such as introns, provides a tool for designing and constructing novel vectors for tissue specific, hormone regulated or any other defined expression pattern, for gene therapy. Such an expression cassette was developed, utilizing regulatory elements from the human cytokeratin 18 (K18) gene, including 5' genomic sequences and one of its introns. This cassette efficiently expresses reporter genes, as well as the human cystic fibrosis transmembrane conductance regulator (CFTR) gene, in cultured lung epithelial cells (49).

#### ***Alternative splicing:***

Alternative splicing of pre mRNA is a powerful and versatile regulatory mechanism that can effect quantitative control of gene expression and functional diversification of proteins. It contributes to major developmental decisions and also to a fine-tuning of gene function. Genetic and biochemical approaches have identified cis-acting regulatory elements and trans-acting factors that control alternative splicing of specific mRNAs. This mechanism results in the generation of variant isoforms of various proteins from a single gene. These include cell surface molecules such as CD44, receptors, cytokines such as VEGF and enzymes. Products of



alternatively spliced transcripts differ in their expression pattern, substrate specificity and other biological parameters.

The FGF receptor RNA undergoes alternative splicing which results in the production of several isoforms, which exhibit different ligand binding specificities. The alternative splicing is regulated in a cell specific manner (53).

Alternative spliced mRNAs are often correlated with malignancy. An increase in specific splice variant of tyrosinase was identified in murine melanomas (54). Multiple splicing variants of estrogen receptor are present in individual human breast tumors. CD44 has various isoform, some are characteristic of malignant tissues.

Identification of tumor specific alternative splice variants provide new tool for cancer diagnostics. CD44 variants have been used for detection of malignancy in urine samples from patients with urothelial cancer by competitive RT-PCR (55). CD44 exon 6 was suggested as prognostic indicator of metastasis in breast cancer (56).

Different enzymes or polypeptides generated by alternative splicing may have different function or catalytic specificity. The identification and characterization of the enzyme forms, which are involved in pathological processes, is crucial for the design of appropriate and efficient drugs.

***Modulation of gene expression – Antisense technology:***

An antisense oligonucleotide (e.g., antisense oligodeoxyribonucleotide) may bind its target nucleic acid either by Watson-Crick base pairing or Hoogsteen and anti-Hoogsteen base pairing (64). According to the Watson-Crick base pairing, heterocyclic bases of the antisense oligonucleotide form hydrogen bonds with the heterocyclic bases of target single-stranded nucleic acids (RNA or single-stranded DNA), whereas according to the Hoogsteen base pairing, the heterocyclic bases of the target nucleic acid are double-stranded DNA, wherein a third strand is accommodated in the major groove of the B-form DNA duplex by Hoogsteen and anti-Hoogsteen base pairing to form a triple helix structure.

According to both the Watson-Crick and the Hoogsteen base pairing models, antisense oligonucleotides have the potential to regulate gene expression and to disrupt the essential functions of the nucleic acids in cells. Therefore, antisense oligonucleotides have possible uses in modulating a wide range of diseases in which gene expression is altered.

Since the development of effective methods for chemically synthesizing oligonucleotides, these molecules have been extensively used

in biochemistry and biological research and have the potential use in medicine, since carefully devised oligonucleotides can be used to control gene expression by regulating levels of transcription, transcripts and/or translation.

5 Oligodeoxyribonucleotides as long as 100 base pairs (bp) are routinely synthesized by solid phase methods using commercially available, fully automated synthesis machines. The chemical synthesis of oligoribonucleotides, however, is far less routine. Oligoribonucleotides are also much less stable than oligodeoxyribonucleotides, a fact which has  
10 contributed to the more prevalent use of oligodeoxyribonucleotides in medical and biological research, directed at, for example, the regulation of transcription or translation levels.

Gene expression involves few distinct and well regulated steps. The first major step of gene expression involves transcription of a messenger  
15 RNA (mRNA) which is an RNA sequence complementary to the antisense (i.e., -) DNA strand, or, in other words, identical in sequence to the DNA sense (i.e., +) strand, composing the gene. In eukaryotes, transcription occurs in the cell nucleus.

The second major step of gene expression involves translation of a  
20 protein (e.g., enzymes, structural proteins, secreted proteins, gene expression factors, etc.) in which the mRNA interacts with ribosomal RNA complexes (ribosomes) and amino acid activated transfer RNAs (tRNAs) to direct the synthesis of the protein coded for by the mRNA sequence.

Initiation of transcription requires specific recognition of a promoter  
25 DNA sequence located upstream to the coding sequence of a gene by an RNA-synthesizing enzyme -- RNA polymerase. This recognition is preceded by sequence-specific binding of one or more transcription factors to the promoter sequence. Additional proteins which bind at or close to the promoter sequence may trans upregulate transcription via cis elements  
30 known as enhancer sequences. Other proteins which bind to or close to the promoter, but whose binding prohibits the action of RNA polymerase, are known as repressors.

There are also evidence that in some cases gene expression is downregulated by endogenous antisense RNA repressors that bind a  
35 complementary mRNA transcript and thereby prevent its translation into a functional protein.

Thus, gene expression is typically upregulated by transcription factors and enhancers and downregulated by repressors.

However, in many disease situation gene expression is impaired. In many cases, such as different types of cancer, for various reasons the expression of a specific endogenous or exogenous (e.g., of a pathogen such as a virus) gene is upregulated. Furthermore, in infectious diseases caused by pathogens such as parasites, bacteria or viruses, the disease progression depends on expression of the pathogen genes, this phenomenon may also be considered as far as the patient is concerned as upregulation of exogenous genes.

Most conventional drugs function by interaction with and modulation of one or more targeted endogenous or exogenous proteins, e.g., enzymes. Such drugs, however, typically are not specific for targeted proteins but interact with other proteins as well. Thus, a relatively large dose of drug must be used to effectively modulate a targeted protein.

Typical daily doses of drugs are from  $10^{-5}$  -  $10^{-1}$  millimoles per kilogram of body weight or  $10^{-3}$  - 10 millimoles for a 100 kilogram person. If this modulation instead could be effected by interaction with and inactivation of mRNA, a dramatic reduction in the necessary amount of drug could likely be achieved, along with a corresponding reduction in side effects. Further reductions could be effected if such interaction could be rendered site-specific. Given that a functioning gene continually produces mRNA, it would thus be even more advantageous if gene transcription could be arrested in its entirety.

Given these facts, it would be advantageous if gene expression could be arrested or downmodulated at the transcription level.

The ability of chemically synthesizing oligonucleotides and analogs thereof having a selected predetermined sequence offers means for downmodulating gene expression. Three types of gene expression modulation strategies may be considered.

At the transcription level, antisense or sense oligonucleotides or analogs that bind to the genomic DNA by strand displacement or the formation of a triple helix, may prevent transcription (64).

At the transcript level, antisense oligonucleotides or analogs that bind target mRNA molecules lead to the enzymatic cleavage of the hybrid by intracellular RNase H (65). In this case, by hybridizing to the targeted mRNA, the oligonucleotides or oligonucleotide analogs provide a duplex hybrid recognized and destroyed by the RNase H enzyme. Alternatively, such hybrid formation may lead to interference with correct splicing (66).

As a result, in both cases, the number of the target mRNA intact transcripts ready for translation is reduced or eliminated.

At the translation level, antisense oligonucleotides or analogs that bind target mRNA molecules prevent, by steric hindrance, binding of essential translation factors (ribosomes), to the target mRNA, a phenomenon known in the art as hybridization arrest, disabling the translation of such mRNAs (67).

Thus, antisense sequences, which as described hereinabove may arrest the expression of any endogenous and/or exogenous gene depending on their specific sequence, attracted much attention by scientists and pharmacologists who were devoted at developing the antisense approach into a new pharmacological tool (68).

For example, several antisense oligonucleotides have been shown to arrest hematopoietic cell proliferation (69), growth (70), entry into the S phase of the cell cycle (71), reduced survival (72) and prevent receptor mediated responses (73). For use of antisense oligonucleotides as antiviral agents the reader is referred to reference 74.

For efficient *in vivo* inhibition of gene expression using antisense oligonucleotides or analogs, the oligonucleotides or analogs must fulfill the following requirements (i) sufficient specificity in binding to the target sequence; (ii) solubility in water; (iii) stability against intra- and extracellular nucleases; (iv) capability of penetration through the cell membrane; and (v) when used to treat an organism, low toxicity.

Unmodified oligonucleotides are impractical for use as antisense sequences since they have short *in vivo* half-lives, during which they are degraded rapidly by nucleases. Furthermore, they are difficult to prepare in more than milligram quantities. In addition, such oligonucleotides are poor cell membrane penetrators (75).

Thus it is apparent that in order to meet all the above listed requirements, oligonucleotide analogs need to be devised in a suitable manner. Therefore, an extensive search for modified oligonucleotides has been initiated.

For example, problems arising in connection with double-stranded DNA (dsDNA) recognition through triple helix formation have been diminished by a clever "switch back" chemical linking, whereby a sequence of polypurine on one strand is recognized, and by "switching back", a homopurine sequence on the other strand can be recognized. Also, good

helix formation has been obtained by using artificial bases, thereby improving binding conditions with regard to ionic strength and pH.

In addition, in order to improve half-life as well as membrane penetration, a large number of variations in polynucleotide backbones have  
5 been done, nevertheless with little success.

Oligonucleotides can be modified either in the base, the sugar or the phosphate moiety. These modifications include, for example, the use of methylphosphonates, monothiophosphates, dithiophosphates, phosphoramidates, phosphate esters, bridged phosphorothioates, bridged  
10 phosphoramidates, bridged methylenephosphonates, dephospho internucleotide analogs with siloxane bridges, carbonate bridges, carboxymethyl ester bridges, carbonate bridges, carboxymethyl ester bridges, acetamide bridges, carbamate bridges, thioether bridges, sulfoxy bridges, sulfono bridges, various "plastic" DNAs,  $\alpha$ -anomeric bridges and  
15 borane derivatives. For further details the reader is referred to reference 76.

International patent application WO 89/12060 discloses various building blocks for synthesizing oligonucleotide analogs, as well as oligonucleotide analogs formed by joining such building blocks in a defined sequence. The building blocks may be either "rigid" (i.e., containing a ring  
20 structure) or "flexible" (i.e., lacking a ring structure). In both cases, the building blocks contain a hydroxy group and a mercapto group, through which the building blocks are said to join to form oligonucleotide analogs. The linking moiety in the oligonucleotide analogs is selected from the group consisting of sulfide (-S-), sulfoxide (-SO-), and sulfone (-SO<sub>2</sub>-). However,  
25 the application provides no data supporting the specific binding of an oligonucleotide analog to a target oligonucleotide.

International patent application WO 92/20702 describe an acyclic oligonucleotide which includes a peptide backbone on which any selected chemical nucleobases or analogs are stringed and serve as coding characters  
30 as they do in natural DNA or RNA. These new compounds, known as peptide nucleic acids (PNAs), are not only more stable in cells than their natural counterparts, but also bind natural DNA and RNA 50 to 100 times more tightly than the natural nucleic acids cling to each other (77). PNA oligomers can be synthesized from the four protected monomers containing  
35 thymine, cytosine, adenine and guanine by Merrifield solid-phase peptide synthesis. In order to increase solubility in water and to prevent aggregation, a lysine amide group is placed at the C-terminal.

Thus, antisense technology requires pairing of messenger RNA with an oligonucleotide to form a double helix that inhibits translation. The concept of antisense-mediated gene therapy was already introduced in 1978 for cancer therapy. This approach was based on certain genes that are crucial in cell division and growth of cancer cells. Synthetic fragments of genetic substance DNA can achieve this goal. Such molecules bind to the targeted gene molecules in RNA of tumor cells, thereby inhibiting the translation of the genes and resulting in dysfunctional growth of these cells. Other mechanisms has also been proposed. These strategies have been used, with some success in treatment of cancers, as well as other illnesses, including viral and other infectious diseases. Antisense oligonucleotides are typically synthesized in lengths of 13-30 nucleotides. The life span of oligonucleotide molecules in blood is rather short. Thus, they have to be chemically modified to prevent destruction by ubiquitous nucleases present in the body. Phosphorothioates are very widely used modification in antisense oligonucleotide ongoing clinical trials (57). A new generation of antisense molecules consist of hybrid antisense oligonucleotide with a central portion of synthetic DNA while four bases on each end have been modified with 2'O-methyl ribose to resemble RNA. In preclinical studies in laboratory animals, such compounds have demonstrated greater stability to metabolism in body tissues and an improved safety profile when compared with the first-generation unmodified phosphorothioate (Hybridon Inc. news). Dosens of other nucleotide analogs have also been tested in antisense technology.

RNA oligonucleotides may also be used for antisense inhibition as they form a stable RNA-RNA duplex with the target, suggesting efficient inhibition. However, due to their low stability RNA oligonucleotides are typically expressed inside the cells using vectors designed for this purpose. This approach is favored when attempting to target a mRNA that encodes an abundant and long-lived protein (57).

Recent scientific publications have validated the efficacy of antisense compounds in animal models of hepatitis, cancers, coronary artery restenosis and other diseases. The first antisense drug was recently approved by the FDA. This drug Fomivirsen, developed by Isis, is indicated for local treatment of cytomegalovirus in patients with AIDS who are intolerant of or have a contraindication to other treatments for CMV retinitis or who were insufficiently responsive to previous treatments for CMV retinitis (Pharmacotherapy News Network).

Several antisense compounds are now in clinical trials in the United States. These include locally administered antivirals, systemic cancer therapeutics. Antisense therapeutics has the potential to treat many life-threatening diseases with a number of advantages over traditional drugs. Traditional drugs intervene after a disease-causing protein is formed. Antisense therapeutics, however, block mRNA transcription/translation and intervene before a protein is formed, and since antisense therapeutics target only one specific mRNA, they should be more effective with fewer side effects than current protein-inhibiting therapy.

A second option for disrupting gene expression at the level of transcription uses synthetic oligonucleotides capable of hybridizing with double stranded DNA. A triple helix is formed. Such oligonucleotides may prevent binding of transcription factors to the gene's promoter and therefore inhibit transcription. Alternatively, they may prevent duplex unwinding and, therefore, transcription of genes within the triple helical structure.

Another approach is the use of specific nucleic acid sequences to act as decoys for transcription factors. Since transcription factors bind specific DNA sequences it is possible to synthesize oligonucleotides that will effectively compete with the native DNA sequences for available transcription factors *in vivo*. This approach requires the identification of gene specific transcription factor (57).

Indirect inhibition of gene expression was demonstrated for matrix metalloproteinase genes (MMP-1, -3, and -9), which are associated with invasive potential of human cancer cells. E1AF is a transcription activator of MMP genes. Expression of E1AF antisense RNA in HSC3AS cells showed decrease in mRNA and protein levels of MMP-1, -3, and -9. Moreover, HSC3AS showed lower invasive potential *in vitro* and *in vivo*. These results imply that transfection of antisense inhibits tumor invasion by down-regulating MMP genes (58).

#### **Ribozymes:**

Ribozymes are being increasingly used for the sequence-specific inhibition of gene expression by the cleavage of mRNAs encoding proteins of interest. The possibility of designing ribozymes to cleave any specific target RNA has rendered them valuable tools in both basic research and therapeutic applications. In the therapeutics area, ribozymes have been exploited to target viral RNAs in infectious diseases, dominant oncogenes in cancers and specific somatic mutations in genetic disorders. Most notably, several ribozyme gene therapy protocols for HIV patients are

already in Phase 1 trials (62). More recently, ribozymes have been used for transgenic animal research, gene target validation and pathway elucidation. Several ribozymes are in various stages of clinical trials. ANGIOZYME was the first chemically synthesized ribozyme to be studied in human clinical trials. ANGIOZYME specifically inhibits formation of the VEGF-r (Vascular Endothelial Growth Factor receptor), a key component in the angiogenesis pathway. Ribozyme Pharmaceuticals, Inc., as well as other firms have demonstrated the importance of anti-angiogenesis therapeutics in animal models. HEPTAZYME, a ribozyme designed to selectively destroy Hepatitis C Virus (HCV) RNA, was found effective in decreasing Hepatitis C viral RNA in cell culture assays (Ribozyme Pharmaceuticals, Incorporated - WEB home page).

***Gene disruption in animal models:***

The emergence of gene inactivation by homologous recombination methodology in embryonic stem cells has revolutionized the field of mouse genetics. The availability of a rapidly growing number of mouse null mutants has represented an invaluable source of knowledge on mammalian development, cellular biology and physiology, and has provided many models for human inherited diseases. Animal models are required for an effective drug delivery development program and evaluation of gene therapy approach. The improvement of the original knockout strategy, as well as exploitation of exogenous enzymatic systems that are active in the recombination process, has been considerably extended the range of genetic manipulations that can be produced. Additional methods have been developed to provide versatile research tools: Double replacement method, sequential gene targeting, conditional cell type specific gene targeting, single copy integration method, inducible gene targeting, gene disruption by viral delivery, replacing one gene with another, the so called knock-in method and the induction of specific balanced chromosomal translocation. It is now possible to introduce a point mutation as a unique change in the entire genome, therefore allowing very fine dissection of gene function *in vivo*. Furthermore, the advent of methods allowing conditional gene targeting opens the way for analysis of consequence of a particular mutation in a defined organ and at a specific time during the life of the experimental animal (59).

***DNA vaccination:***

Observations in the early 1990s that plasmid DNA could directly transfect animal cells *in vivo* sparked exploration of the use of DNA



plasmids to induce immune response by direct injection into animal of DNA encoding antigenic protein. When a DNA vaccine plasmid enters the eukaryotic cell, the protein it encodes is transcribed and translated within the cell. In the case of pathogens, these proteins are presented to the immune system in their native form, mimicking the presentation of antigens during a natural infection. DNA vaccination is particularly useful for the induction of T cell activation. It was applied for viral and bacterial infectious diseases, as well as for allergy and for cancer. The central hypothesis behind active specific immunotherapy for cancer is that tumor cells express unique antigens that should stimulate the immune system. The first DNA vaccine against tumor was carcino-embryonic antigen (CEA). DNA vaccinated animals expressed immunoprotection and immunotherapy of human CEA-expressing syngeneic mouse colon and breast carcinoma (61). In a mouse model of neuroblastoma, DNA immunization with HuD resulted in tumor growth inhibition with no neurological disease (60). Immunity to the brown locus protein, gp<sup>75</sup> tyrosinase-related protein-1, associated with melanoma, was investigated in a syngeneic mouse model. Priming with human gp<sup>75</sup> DNA broke tolerance to mouse gp<sup>75</sup>. Immunity against mouse gp<sup>75</sup> provided significant tumor protection (60).

***Glycosyl hydrolases:***

Glycosyl hydrolases are a widespread group of enzymes that hydrolyze the o-glycosidic bond between two or more carbohydrates or between a carbohydrate and a noncarbohydrate moiety. The enzymatic hydrolysis of glycosidic bond occurs by using major one or two mechanisms leading to overall retention or inversion of the anomeric configuration. In both mechanisms catalysis involves two residues: a proton donor and a nucleophile. Glycosyl hydrolases have been classified into 58 families based on amino acid similarities. The glycosyl hydrolases from families 1, 2, 5, 10, 17, 30, 35, 39 and 42 act on a large variety of substrates, however, they all hydrolyze the glycosidic bond in a general acid catalysis mechanism, with retention of the anomeric configuration. The mechanism involves two glutamic acid residues, which are the proton donors and the nucleophile, with an asparagine always preceding the proton donor. Analyses of a set of known 3D structures from this group revealed that their catalytic domains, despite the low level of sequence identity, adopt a similar ( $\alpha/\beta$ ) 8 fold with the proton donor and the nucleophile located at the C-terminal ends of strands  $\beta$ 4 and  $\beta$ 7, respectively. Mutations

in the functional conserved amino acids of lysosomal glycosyl hydrolases were identified in lysosomal storage diseases.

Lysosomal glycosyl hydrolases including  $\beta$ -glucuronidase,  $\beta$ -mannosidase,  $\beta$ -glucocerebrosidase,  $\beta$ -galactosidase and  $\alpha$ -L iduronidase, are all exo-glycosyl hydrolases, belong to the GH-A clan and share a similar catalytic site. However, many endo-glucanases from various organisms, such as bacterial and fungal xylanases and cellulases share this catalytic domain.

***Genomic sequence of hpa gene and its implications:***

It is well established that heparanase activity is correlated with cancer metastasis. This correlation was demonstrated at the level of enzymatic activity as well as the levels of protein and *hpa* cDNA expression in highly metastatic cancer cells as compared with non-metastatic cells. As such, inhibition of heparanase activity is desirable, and has been attempted by several means. The genomic region, encoding the *hpa* gene and the surrounding, provides a new powerful tool for regulation of heparanase activity at the level of gene expression. Regulatory sequences may reside in noncoding regions both upstream and downstream the transcribed region as well as in intron sequences. A DNA sequence upstream of the transcription start site contains the promoter region and potential regulatory elements. Regulatory factors, which interact with the promoter region may be identified and be used as potential drugs for inhibition of cancer, metastasis and inflammation. The promoter region can be used to screen for inhibitors of heparanase gene expression. Furthermore, the *hpa* promoter can be used to direct cell specific, particularly cancer cell specific, expression of foreign genes, such as cytotoxic or apoptotic genes, in order to specifically destroy cancer cells.

Cancer and yet unknown related genetic disorders may involve rearrangements and mutations in the heparanase gene, either in coding or non-coding regions. Such mutations may affect expression level or enzymatic activity. The genomic sequence of *hpa* enables the amplification of specific genomic DNA fragments, identification and diagnosis of mutations.

There is thus a widely recognized need for, and it would be highly advantageous to have genomic, cDNA and composite polynucleotides encoding a polypeptide having heparanase activity, vectors including same, genetically modified cells expressing heparanase and a recombinant protein

having heparanase activity, as well as antisense oligonucleotides, constructs and ribozymes which can be used for down regulation heparanase activity.

#### SUMMARY OF THE INVENTION

5 Cloning of the human *hpa* gene which encodes heparanase, and expression of recombinant heparanase by transfected host cells is reported herein, as well as downregulation of heparanase activity by antisense technology.

10 A purified preparation of heparanase isolated from human hepatoma cells was subjected to tryptic digestion and microsequencing. The YGPDVGQPR (SEQ ID NO:8) sequence revealed was used to screen EST databases for homology to the corresponding back translated DNA sequence. Two closely related EST sequences were identified and were thereafter found to be identical. Both clones contained an insert of 1020 bp  
15 which included an open reading frame of 973 bp followed by a 27 bp of 3' untranslated region and a Poly A tail. Translation start site was not identified.

Cloning of the missing 5' end of *hpa* was performed by PCR amplification of DNA from placenta Marathon RACE cDNA composite  
20 using primers selected according to the EST clones sequence and the linkers of the composite. A 900 bp PCR fragment, partially overlapping with the identified 3' encoding EST clones was obtained. The joined cDNA fragment (*hpa*), 1721 bp long (SEQ ID NO:9), contained an open reading frame which encodes a polypeptide of 543 amino acids (SEQ ID NO:10)  
25 with a calculated molecular weight of 61,192 daltons.

Cloning an extended 5' sequence was enabled from the human SK-hep1 cell line by PCR amplification using the Marathon RACE. The 5' extended sequence of the SK-hep1 *hpa* cDNA was assembled with the sequence of the *hpa* cDNA isolated from human placenta (SEQ ID NO:9).  
30 The assembled sequence contained an open reading frame, SEQ ID NOs: 13 and 15, which encodes, as shown in SEQ ID NOs:14 and 15, a polypeptide of 592 amino acids with a calculated molecular weight of 66,407 daltons.

The ability of the *hpa* gene product to catalyze degradation of heparan sulfate in an *in vitro* assay was examined by expressing the entire  
35 open reading frame of *hpa* in insect cells, using the Baculovirus expression system. Extracts and conditioned media of cells infected with virus containing the *hpa* gene, demonstrated a high level of heparan sulfate degradation activity both towards soluble ECM-derived HSPG and intact

ECM. This degradation activity was inhibited by heparin, which is another substrate of heparanase. Cells infected with a similar construct containing no *hpa* gene had no such activity, nor did non-infected cells. The ability of heparanase expressed from the extended 5' clone towards heparin was demonstrated in a mammalian expression system.

The expression pattern of *hpa* RNA in various tissues and cell lines was investigated using RT-PCR. It was found to be expressed only in tissues and cells previously known to have heparanase activity.

A panel of monochromosomal human/CHO and human/mouse somatic cell hybrids was used to localize the human heparanase gene to human chromosome 4. The newly isolated heparanase sequence can be used to identify a chromosome region harboring a human heparanase gene in a chromosome spread.

A human genomic library was screened and the human locus harboring the heparanase gene isolated, sequenced and characterized. Alternatively spliced heparanase mRNAs were identified and characterized. The human heparanase promoter has been isolated, identified and positively tested for activity. The mouse heparanase promoter has been isolated and identified as well. Antisense heparanase constructs were prepared and their influence on cells *in vitro* tested. A predicted heparanase active site was identified. And finally, the presence of sequences hybridizing with human heparanase sequences was demonstrated for a variety of mammals and for an avian.

According to one aspect of the present invention there is provided an isolated nucleic acid comprising a genomic, complementary or composite polynucleotide sequence encoding a polypeptide having heparanase catalytic activity.

According to further features in preferred embodiments of the invention described below, the polynucleotide or a portion thereof is hybridizable with SEQ ID NOs: 9, 13, 42, 43 or a portion thereof at 68 °C in 6 x SSC, 1 % SDS, 5 x Denharts, 10 % dextran sulfate, 100 µg/ml salmon sperm DNA, and <sup>32</sup>p labeled probe and wash at 68 °C with 3 x SSC and 0.1 % SDS.

According to still further features in the described preferred embodiments the polynucleotide or a portion thereof is at least 60 % identical with SEQ ID NOs: 9, 13, 42, 43 or portions thereof as determined using the Bestfit procedure of the DNA sequence analysis software package

developed by the Genetic Computer Group (GCG) at the university of Wisconsin (gap creation penalty - 12, gap extension penalty - 4).

According to still further features in the described preferred embodiments the polypeptide is as set forth in SEQ ID NOs:10, 14, 44 or  
5 portions thereof.

According to still further features in the described preferred embodiments the polypeptide is at least 60 % homologous to SEQ ID NOs:10, 14, 44 or portions thereof as determined with the Smith-Waterman algorithm, using the Bioaccelerator platform developed by Compugene  
10 (gapop: 10.0, gapext: 0.5, matrix: blosum62).

According to additional aspects of the present invention there are provided a nucleic acid construct (vector) comprising the isolated nucleic acid described herein and a host cell comprising the construct.

According to a further aspect of the present invention there is  
15 provided an antisense oligonucleotide comprising a polynucleotide or a polynucleotide analog of at least 10 bases being hybridizable *in vivo*, under physiological conditions, with a portion of a polynucleotide strand encoding a polypeptide having heparanase catalytic activity.

According to an additional aspect of the present invention there is  
20 provided a method of *in vivo* downregulating heparanase activity comprising the step of *in vivo* administering the antisense oligonucleotide herein described.

According to yet an additional aspect of the present invention there is provided a pharmaceutical composition comprising the antisense  
25 oligonucleotide herein described and a pharmaceutically acceptable carrier.

According to still an additional aspect of the present invention there is provided a ribozyme comprising the antisense oligonucleotide described herein and a ribozyme sequence.

According to a further aspect of the present invention there is  
30 provided an antisense nucleic acid construct comprising a promoter sequence and a polynucleotide sequence directing the synthesis of an antisense RNA sequence of at least 10 bases being hybridizable *in vivo*, under physiological conditions, with a portion of a polynucleotide strand encoding a polypeptide having heparanase catalytic activity.

35 According to further features in preferred embodiments of the invention described below, the polynucleotide strand encoding the polypeptide having heparanase catalytic activity is as set forth in SEQ ID NOs: 9, 13, 42 or 43.

According to still further features in the described preferred embodiments the polypeptide having heparanase catalytic activity is as set forth in SEQ ID NOs: 10, 14 or 44.

5 According to still a further aspect of the present invention there is provided a method of *in vivo* downregulating heparanase activity comprising the step of *in vivo* administering the antisense nucleic acid construct herein described.

10 According to yet a further aspect of the present invention there is provided a pharmaceutical composition comprising the antisense nucleic acid construct herein described and a pharmaceutically acceptable carrier.

According to a further aspect of the present invention there is provided a nucleic acid construct comprising a polynucleotide sequence functioning as a promoter, the polynucleotide sequence is derived from SEQ ID NO:42 and includes at least nucleotides 2535-2635 thereof or from  
15 SEQ ID NO:43 and includes at least nucleotides 320-420.

According to a further aspect of the present invention there is provided a method of expressing a polynucleotide sequence comprising the step of ligating the polynucleotide sequence into the nucleic acid construct described above, downstream of the polynucleotide sequence derived from  
20 SEQ ID NOs:42 or 43.

According to a further aspect of the present invention there is provided a recombinant protein comprising a polypeptide having heparanase catalytic activity.

25 According to further features in preferred embodiments of the invention described below, the polypeptide includes at least a portion of SEQ ID NOs:10, 14 or 44.

According to still further features in the described preferred embodiments the protein is encoded by a polynucleotide hybridizable with SEQ ID NOs: 9, 13, 42, 43 or a portion thereof at 68 °C in 6 x SSC, 1 %  
30 SDS, 5 x Denharts, 10 % dextran sulfate, 100 µg/ml salmon sperm DNA, and <sup>32</sup>p labeled probe and wash at 68 °C with 3 x SSC and 0.1 % SDS.

According to still further features in the described preferred embodiments the protein is encoded by a polynucleotide at least 60 % identical with SEQ ID NOs: 9, 13, 42, 43 or portions thereof as determined  
35 using the Bestfit procedure of the DNA sequence analysis software package developed by the Genetic Computer Group (GCG) at the university of Wisconsin (gap creation penalty - 12, gap extension penalty - 4).

According to a further aspect of the present invention there is provided a pharmaceutical composition comprising, as an active ingredient, the recombinant protein herein described.

According to a further aspect of the present invention there is provided a method of identifying a chromosome region harboring a heparanase gene in a chromosome spread comprising the steps of (a) hybridizing the chromosome spread with a tagged polynucleotide probe encoding heparanase; (b) washing the chromosome spread, thereby removing excess of non-hybridized probe; and (c) searching for signals associated with the hybridized tagged polynucleotide probe, wherein detected signals being indicative of a chromosome region harboring a heparanase gene.

According to a further aspect of the present invention there is provided a method of *in vivo* eliciting anti-heparanase antibodies comprising the steps of administering a nucleic acid construct including a polynucleotide segment corresponding to at least a portion of SEQ ID NOs:9, 13 or 43 and a promoter for directing the expression of said polynucleotide segment *in vivo*. Accordingly, there is provided also a DNA vaccine for *in vivo* eliciting anti-heparanase antibodies comprising a nucleic acid construct including a polynucleotide segment corresponding to at least a portion of SEQ ID NOs:9, 13 or 43 and a promoter for directing the expression of said polynucleotide segment *in vivo*.

The present invention can be used to develop new drugs to inhibit tumor cell metastasis, inflammation and autoimmunity. The identification of the *hpa* gene encoding for heparanase enzyme enables the production of a recombinant enzyme in heterologous expression systems. Additional features, advantages, uses and applications of the present invention in biological science and in diagnostic and therapeutic medicine are described hereinafter.

#### BRIEF DESCRIPTION OF THE DRAWINGS

The invention herein described, by way of example only, with reference to the accompanying drawings, wherein:

FIG. 1 presents nucleotide sequence and deduced amino acid sequence of *hpa* cDNA. A single nucleotide difference at position 799 (A to T) between the EST (Expressed Sequence Tag) and the PCR amplified cDNA (reverse transcribed RNA) and the resulting amino acid substitution (Tyr to Phe) are indicated above and below the substituted unit,

respectively. Cysteine residues and the poly adenylation consensus sequence are underlined. The asterisk denotes the stop codon TGA.

FIG. 2 demonstrates degradation of soluble sulfate labeled HSPG substrate by lysates of High Five cells infected with pFhpa2 virus. Lysates of High Five cells that were infected with pFhpa2 virus (●) or control pF2 virus (□) were incubated (18 h, 37 °C) with sulfate labeled ECM-derived soluble HSPG (peak I). The incubation medium was then subjected to gel filtration on Sepharose 6B. Low molecular weight HS degradation fragments (peak II) were produced only during incubation with the pFhpa2 infected cells, but there was no degradation of the HSPG substrate (♦) by lysates of pF2 infected cells.

FIGs. 3a-b demonstrate degradation of soluble sulfate labeled HSPG substrate by the culture medium of pFhpa2 and pFhpa4 infected cells. Culture media of High Five cells infected with pFhpa2 (3a) or pFhpa4 (3b) viruses (●), or with control viruses (□) were incubated (18 h, 37 °C) with sulfate labeled ECM-derived soluble HSPG (peak I, ♦). The incubation media were then subjected to gel filtration on Sepharose 6B. Low molecular weight HS degradation fragments (peak II) were produced only during incubation with the hpa gene containing viruses. There was no degradation of the HSPG substrate by the culture medium of cells infected with control viruses.

FIG. 4 presents size fractionation of heparanase activity expressed by pFhpa2 infected cells. Culture medium of pFhpa2 infected High Five cells was applied onto a 50 kDa cut-off membrane. Heparanase activity (conversion of the peak I substrate, ♦) into peak II HS degradation fragments) was found in the high (> 50 kDa) (●), but not low (< 50 kDa) (○) molecular weight compartment.

FIGs. 5a-b demonstrate the effect of heparin on heparanase activity expressed by pFhpa2 and pFhpa4 infected High Five cells. Culture media of pFhpa2 (5a) and pFhpa4 (5b) infected High Five cells were incubated (18 h, 37 °C) with sulfate labeled ECM-derived soluble HSPG (peak I, ♦) in the absence (●) or presence (Δ) of 10 μg/ml heparin. Production of low molecular weight HS degradation fragments was completely abolished in the presence of heparin, a potent inhibitor of heparanase activity (6, 7).

FIGs. 6a-b demonstrate degradation of sulfate labeled intact ECM by virus infected High Five and Sf21 cells. High Five (6a) and Sf21 (6b) cells were plated on sulfate labeled ECM and infected (48 h, 28 °C) with pFhpa4 (●) or control pF1 (□) viruses. Control non-infected Sf21 cells (R) were



plated on the labeled ECM as well. The pH of the cultured medium was adjusted to 6.0 - 6.2 followed by 24 h incubation at 37 °C. Sulfate labeled material released into the incubation medium was analyzed by gel filtration on Sepharose 6B. HS degradation fragments were produced only by cells  
5 infected with the *hpa* containing virus.

FIG. 7a-b demonstrate degradation of sulfate labeled intact ECM by virus infected cells. High Five (7a) and Sf21 (7b) cells were plated on sulfate labeled ECM and infected (48 h, 28 °C) with pF*hpa*4 (●) or control pF1 (□) viruses. Control non-infected Sf21 cells (R) were plate on labeled  
10 ECM as well. The pH of the cultured medium was adjusted to 6.0 - 6.2, followed by 48 h incubation at 28 °C. Sulfate labeled degradation fragments released into the incubation medium was analyzed by gel filtration on Sepharose 6B. HS degradation fragments were produced only by cells infected with the *hpa* containing virus.

FIGs. 8a-b demonstrate degradation of sulfate labeled intact ECM by the culture medium of pF*hpa*4 infected cells. Culture media of High Five (8a) and Sf21 (8b) cells that were infected with pF*hpa*4 (●) or control pF1 (□) viruses were incubated (48 h, 37 °C, pH 6.0) with intact sulfate labeled ECM. The ECM was also incubated with the culture medium of control  
15 non-infected Sf21 cells (R). Sulfate labeled material released into the reaction mixture was subjected to gel filtration analysis. Heparanase activity was detected only in the culture medium of pF*hpa*4 infected cells.

FIGs. 9a-b demonstrate the effect of heparin on heparanase activity in the culture medium of pF*hpa*4 infected cells. Sulfate labeled ECM was incubated (24 h, 37 °C, pH 6.0) with culture medium of pF*hpa*4 infected High Five (9a) and Sf21 (9b) cells in the absence (●) or presence (V) of 10 µg/ml heparin. Sulfate labeled material released into the incubation  
25 medium was subjected to gel filtration on Sepharose 6B. Heparanase activity (production of peak II HS degradation fragments) was completely inhibited in the presence of heparin.

FIGs. 10a-b demonstrate purification of recombinant heparanase on heparin-Sepharose. Culture medium of Sf21 cells infected with pF*hpa*4 virus was subjected to heparin-Sepharose chromatography. Elution of fractions was performed with 0.35 - 2 M NaCl gradient (◇). Heparanase activity in the eluted fractions is demonstrated in Figure 10a (●). Fractions  
35 15-28 were subjected to 15 % SDS-polyacrylamide gel electrophoresis followed by silver nitrate staining. A correlation is demonstrated between a

major protein band (MW ~ 63,000) in fractions 19 - 24 and heparanase activity.

FIGs. 11a-b demonstrate purification of recombinant heparanase on a Superdex 75 gel filtration column. Active fractions eluted from heparin-Sephadex (Figure 10a) were pooled, concentrated and applied onto Superdex 75 FPLC column. Fractions were collected and aliquots of each fraction were tested for heparanase activity (C, Figure 11a) and analyzed by SDS-polyacrylamide gel electrophoresis followed by silver nitrate staining (Figure 11b). A correlation is seen between the appearance of a major protein band (MW ~ 63,000) in fractions 4 - 7 and heparanase activity.

FIGs. 12a-e demonstrate expression of the *hpa* gene by RT-PCR with total RNA from human embryonal tissues (12a), human extra-embryonal tissues (12b) and cell lines from different origins (12c-e). RT-PCR products using *hpa* specific primers (I), primers for GAPDH housekeeping gene (II), and control reactions without reverse transcriptase demonstrating absence of genomic DNA or other contamination in RNA samples (III). M- DNA molecular weight marker VI (Boehringer Mannheim). For 12a: lane 1 - neutrophil cells (adult), lane 2 - muscle, lane 3 - thymus, lane 4 - heart, lane 5 - adrenal. For 12b: lane 1 - kidney, lane 2 - placenta (8 weeks), lane 3 - placenta (11 weeks), lanes 4-7 - mole (complete hydatidiform mole), lane 8 - cytotrophoblast cells (freshly isolated), lane 9 - cytotrophoblast cells (1.5 h *in vitro*), lane 10 - cytotrophoblast cells (6 h *in vitro*), lane 11 - cytotrophoblast cells (18 h *in vitro*), lane 12 - cytotrophoblast cells (48 h *in vitro*). For 12c: lane 1 - JAR bladder cell line, lane 2 - NCITT testicular tumor cell line, lane 3 - SW-480 human hepatoma cell line, lane 4 - HTR (cytotrophoblasts transformed by SV40), lane 5 - HPTLP-I hepatocellular carcinoma cell line, lane 6 - EJ-28 bladder carcinoma cell line. For 12d: lane 1 - SK-hep-1 human hepatoma cell line, lane 2 - DAMI human megakaryocytic cell line, lane 3 - DAMI cell line + PMA, lane 4 - CHRF cell line + PMA, lane 5 - CHRF cell line. For 12e: lane 1 - ABAE bovine aortic endothelial cells, lane 2 - 1063 human ovarian cell line, lane 3 - human breast carcinoma MDA435 cell line, lane 4 - human breast carcinoma MDA231 cell line.

FIG. 13 presents a comparison between nucleotide sequences of the human *hpa* and a mouse EST cDNA fragment (SEQ ID NO:12) which is 80 % homologous to the 3' end (starting at nucleotide 1066 of SEQ ID NO:9) of the human *hpa*. The aligned termination codons are underlined.

FIG. 14 demonstrates the chromosomal localization of the *hpa* gene. PCR products of DNA derived from somatic cell hybrids and of genomic DNA of hamster, mouse and human of were separated on 0.7 % agarose gel following amplification with *hpa* specific primers. Lane 1 – Lambda DNA digested with *Bst*EII, lane 2 – no DNA control, lanes 3 – 29, PCR amplification products. Lanes 3-5 – human, mouse and hamster genomic DNA, respectively. Lanes 6-29, human monochromosomal somatic cell hybrids representing chromosomes 1-22 and X and Y, respectively. Lane 30 – Lambda DNA digested with *Bst*EII. An amplification product of approximately 2.8 Kb is observed only in lanes 5 and 9, representing human genomic DNA and DNA derived from cell hybrid carrying human chromosome 4, respectively. These results demonstrate that the *hpa* gene is localized in human chromosome 4.

FIG. 15 demonstrates the genomic exon-intron structure of the human *hpa* locus (top) and the relative positions of the lambda clones used as sequencing templates to sequence the locus (below). The vertical rectangles represent exons (E) and the horizontal lines therebetween represent introns (I), upstream (U) and downstream (D) regions. Continuous lines represent DNA fragments, which were used for sequence analysis. The discontinuous line in lambda 6 represent a region, which overlaps with lambda 8 and hence was not analyzed. The plasmid contains a PCR product, which bridges the gap between L3 and L6.

FIG. 16 presents the nucleotide sequence of the genomic region of the *hpa* gene. Exon sequences appear in upper case and intron sequences in lower case. The deduced amino acid sequence of the exons is printed below the nucleotide sequence. Two predicted transcription start sites are shown in bold.

FIG. 17 presents an alignment of the amino acid sequences of human heparanase, mouse and partial sequences of rat homologues. The human and the mouse sequences were determined by sequence analysis of the isolated cDNAs. The rat sequence is derived from two different EST clones, which represent two different regions (5' and 3') of the rat *hpa* cDNA. The human sequence and the amino acids in the mouse and rat homologues, which are identical to the human sequence, appear in bold.

FIG. 18 presents a heparanase Zoo blot. Ten micrograms of genomic DNA from various sources were digested with *Eco*RI and separated on 0.7 % agarose – TBE gel. Following electrophoresis, the was gel treated with HCl and than with NaOH and the DNA fragments were downward

transferred to a nylon membrane (Hybond N+, Amersham) with 0.4 N NaOH. The membrane was hybridized with a 1.6 Kb DNA probe that contained the entire *hpa* cDNA. Lane order: H – Human; M – Mouse; Rt – Rat; P – Pig; Cw – Cow; Hr – Horse; S – Sheep; Rb – Rabbit; D – Dog; Ch – Chicken; F – Fish. Size markers (Lambda *Bst*II) are shown on the left

FIG. 19 demonstrates the secondary structure prediction for heparanase performed using the PHD server – Profile network Prediction Heidelberg. H – helix, E – extended (beta strand), The glutamic acid predicted as the proton donor is marked by asterisk and the possible nucleophiles are underlined.

#### DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention is of a polynucleotide or nucleic acid, referred to hereinbelow interchangeably as *hpa*, *hpa* cDNA or *hpa* gene or identified by its SEQ ID NOs, encoding a polypeptide having heparanase activity, vectors or nucleic acid constructs including same and which are used for over-expression or antisense inhibition of heparanase, genetically modified cells expressing same, recombinant protein having heparanase activity, antisense oligonucleotides and ribozymes for heparanase modulation, and heparanase promoter sequences which can be used to direct the expression of desired genes.

Before explaining at least one embodiment of the invention in detail, it is to be understood that the invention is not limited in its application to the details of construction and the arrangement of the components set forth in the following description or illustrated in the drawings. The invention is capable of other embodiments or of being practiced or carried out in various ways. Also, it is to be understood that the phraseology and terminology employed herein is for the purpose of description and should not be regarded as limiting.

Cloning of the human and mouse *hpa* genes, cDNAs and genomic sequence (for human), encoding heparanase and expressing recombinant heparanase by transfected cells is reported herein. These are the first mammalian heparanase genes to be cloned.

A purified preparation of heparanase isolated from human hepatoma cells was subjected to tryptic digestion and microsequencing.

The YGPDVGQPR (SEQ ID NO:8) sequence revealed was used to screen EST databases for homology to the corresponding back translated

DNA sequences. Two closely related EST sequences were identified and were thereafter found to be identical.

Both clones contained an insert of 1020 bp which includes an open reading frame of 973 bp followed by a 3' untranslated region of 27 bp and a Poly A tail, whereas a translation start site was not identified.

Cloning of the missing 5' end was performed by PCR amplification of DNA from placenta Marathon RACE cDNA composite using primers selected according to the EST clones sequence and the linkers of the composite.

A 900 bp PCR fragment, partially overlapping with the identified 3' encoding EST clones was obtained. The joined cDNA fragment (*hpa*), 1721 bp long (SEQ ID NO:9), contained an open reading frame which encodes, as shown in Figure 1 and SEQ ID NO:11, a polypeptide of 543 amino acids (SEQ ID NO:10) with a calculated molecular weight of 61,192 daltons.

A single nucleotide difference at position 799 (A to T) between the EST clones and the PCR amplified cDNA was observed. This difference results in a single amino acid substitution (Tyr to Phe) (Figure 1). Furthermore, the published EST sequences contained an unidentified nucleotide, which following DNA sequencing of both the EST clones was resolved into two nucleotides (G and C at positions 1630 and 1631 in SEQ ID NO:9, respectively).

The ability of the *hpa* gene product to catalyze degradation of heparan sulfate in an *in vitro* assay was examined by expressing the entire open reading frame in insect cells, using the Baculovirus expression system.

Extracts and conditioned media of cells infected with virus containing the *hpa* gene, demonstrated a high level of heparan sulfate degradation activity both towards soluble ECM-derived HSPG and intact ECM, which was inhibited by heparin, while cells infected with a similar construct containing no *hpa* gene had no such activity, nor did non-infected cells.

The expression pattern of *hpa* RNA in various tissues and cell lines was investigated using RT-PCR. It was found to be expressed only in tissues and cells previously known to have heparanase activity.

Cloning an extended 5' sequence was enabled from the human SK-hep1 cell line by PCR amplification using the Marathon RACE. The 5' extended sequence of the SK-hep1 *hpa* cDNA was assembled with the sequence of the *hpa* cDNA isolated from human placenta (SEQ ID NO:9).

The assembled sequence contained an open reading frame, SEQ ID NOs: 13 and 15, which encodes, as shown in SEQ ID NOs:14 and 15, a polypeptide of 592 amino acids, with a calculated molecular weight of 66,407 daltons. This open reading frame was shown to direct the expression of catalytically active heparanase in a mammalian cell expression system. The expressed heparanase was detectable by anti heparanase antibodies in Western blot analysis.

A panel of monochromosomal human/CHO and human/mouse somatic cell hybrids was used to localize the human heparanase gene to human chromosome 4. The newly isolated heparanase sequence can therefore be used to identify a chromosome region harboring a human heparanase gene in a chromosome spread.

The *hpa* cDNA was then used as a probe to screen a human genomic library. Several phages were positive. These phages were analyzed and were found to cover most of the *hpa* locus, except for a small portion which was recovered by bridging PCR. The *hpa* locus covers about 50,000 bp. The *hpa* gene includes 12 exons separated by 11 introns.

RT-PCR performed on a variety of cells revealed alternatively spliced *hpa* transcripts.

The amino acid sequence of human heparanase was used to search for homologous sequences in the DNA and protein databases. Several human EST's were identified, as well as mouse sequences highly homologous to human heparanase. The following mouse EST's were identified AA177901, AA674378, AA67997, AA047943, AA690179, AI122034, all sharing an identical sequence and correspond to amino acids 336-543 of the human heparanase sequence. The entire mouse heparanase cDNA was cloned, based on the nucleotide sequence of the mouse EST's using Marathon cDNA libraries. The mouse and the human *hpa* genes share an average homology of 78 % between the nucleotide sequences and 81 % similarity between the deduced amino acid sequences. *hpa* homologous sequences from rat were also uncovered (EST's AI060284 and AI237828).

Homology search of heparanase amino acid sequence against the DNA and the protein databases and prediction of its protein secondary structure enabled to identify candidate amino acids that participate in the heparanase active site.

Expression of *hpa* antisense in mammalian cell lines resulted in about five fold decrease in the number of recoverable cells as compared to controls.

Human *Hpa* cDNA was shown to hybridize with genomic DNAs of a variety of mammalian species and with an avian.

The human and mouse *hpa* promoters were identified and the human promoter was tested positive in directing the expression of a reporter gene.

Thus, according to the present invention there is provided an isolated nucleic acid comprising a genomic, complementary or composite polynucleotide sequence encoding a polypeptide having heparanase catalytic activity.

The phrase "composite polynucleotide sequence" refers to a sequence which includes exonal sequences required to encode the polypeptide having heparanase activity, as well as any number of intronal sequences. The intronal sequences can be of any source and typically will include conserved splicing signal sequences. Such intronal sequences may further include cis acting expression regulatory elements.

The term "heparanase catalytic activity" or its equivalent term "heparanase activity" both refer to a mammalian endoglycosidase hydrolyzing activity which is specific for heparan or heparan sulfate proteoglycan substrates, as opposed to the activity of bacterial enzymes (heparinase I, II and III) which degrade heparin or heparan sulfate by means of  $\beta$ -elimination (37).

According to a preferred embodiment of the present invention the polynucleotide or a portion thereof is hybridizable with SEQ ID NOS: 9, 13, 42, 43 or a portion thereof at 68 °C in 6 x SSC, 1 % SDS, 5 x Denharts, 10 % dextran sulfate, 100  $\mu$ g/ml salmon sperm DNA, and  $^{32}$ p labeled probe and wash at 68 °C with 3, 2, 1, 0.5 or 0.1 x SSC and 0.1 % SDS.

According to another preferred embodiment of the present invention the polynucleotide or a portion thereof is at least 60 %, preferably at least 65 %, more preferably at least 70 %, more preferably at least 75 %, more preferably at least 80 %, more preferably at least 85 %, more preferably at least 90 %, most preferably, 95-100 % identical with SEQ ID NOS: 9, 13, 42, 43 or portions thereof as determined using the Bestfit procedure of the DNA sequence analysis software package developed by the Genetic Computer Group (GCG) at the university of Wisconsin (gap creation penalty - 12, gap extension penalty - 4 - which are the default parameters).

According to another preferred embodiment of the present invention the polypeptide encoded by the polynucleotide sequence is as set forth in SEQ ID NOs:10, 14, 44 or portions thereof having heparanase catalytic activity. Such portions are expected to include amino acids Asp-Glu 224-225 (SEQ ID NO:10), which can serve as proton donors and glutamic acid 343 or 396 which can serve as a nucleophile.

According to another preferred embodiment of the present invention the polypeptide encoded by the polynucleotide sequence is at least 60 %, preferably at least 65 %, more preferably at least 70 %, more preferably at least 75 %, more preferably at least 80 %, more preferably at least 85 %, more preferably at least 90 %, most preferably, 95-100 % homologous (both similar and identical acids) to SEQ ID NOs:10, 14, 44 or portions thereof as determined with the Smith-Waterman algorithm, using the Bioaccelerator platform developed by Compugene (gapop: 10.0, gapext: 0.5, matrix: blosum62, see also the description to Figure 17).

Further according to the present invention there is provided a nucleic acid construct comprising the isolated nucleic acid described herein. The construct may and preferably further include an origin of replication and

The construct or vector can be of any type. It may be a phage which infects bacteria or a virus which infects eukaryotic cells. It may also be a plasmid, phagemid, cosmid, bacmid or an artificial chromosome.

Further according to the present invention there is provided a host cell comprising the nucleic acid construct described herein. The host cell can be of any type. It may be a prokaryotic cell, an eukaryotic cell, a cell line, or a cell as a portion of an organism. The polynucleotide encoding heparanase can be permanently or transiently present in the cell. In other words, genetically modified cells obtained following stable or transient transfection, transformation or transduction are all within the scope of the present invention. The polynucleotide can be present in the cell in low copy (say 1-5 copies) or high copy number (say 5-50 copies or more). It may be integrated in one or more chromosomes at any location or be present as an extrachromosomal material.

The present invention is further directed at providing a heparanase over-expression system which includes a cell overexpressing heparanase catalytic activity. The cell may be a genetically modified host cell transiently or stably transfected or transformed with any suitable vector which includes a polynucleotide sequence encoding a polypeptide having



heparanase activity and a suitable promoter and enhancer sequences to direct over-expression of heparanase. However, the overexpressing cell may also be a product of an insertion (e.g., via homologous recombination) of a promoter and/or enhancer sequence downstream to the endogenous  
5 heparanase gene of the expressing cell, which will direct over-expression from the endogenous gene.

The term "over-expression" as used herein in the specification and claims below refers to a level of expression which is higher than a basal level of expression typically characterizing a given cell under otherwise  
10 identical conditions.

According to another aspect the present invention provides an antisense oligonucleotide comprising a polynucleotide or a polynucleotide analog of at least 10, preferably 11-15, more preferably 16-17, more preferably 18, more preferably 19-25, more preferably 26-35, most  
15 preferably 35-100 bases being hybridizable *in vivo*, under physiological conditions, with a portion of a polynucleotide strand encoding a polypeptide having heparanase catalytic activity. The antisense oligonucleotide can be used for downregulating heparanase activity by *in vivo* administration thereof to a patient. As such, the antisense oligonucleotide according to the  
20 present invention can be used to treat types of cancers which are characterized by impaired (over) expression of heparanase, and are dependent on the expression of heparanase for proliferating or forming metastases.

The antisense oligonucleotide can be DNA or RNA or even include  
25 nucleotide analogs, examples of which are provided in the Background section hereinabove. The antisense oligonucleotide according to the present invention can be synthetic and is preferably prepared by solid phase synthesis. In addition, it can be of any desired length which still provides specific base pairing (e.g., 8 or 10, preferably more, nucleotides long) and it  
30 can include mismatches that do not hamper base pairing under physiological conditions.

Further according to the present invention there is provided a pharmaceutical composition comprising the antisense oligonucleotide herein described and a pharmaceutically acceptable carrier. The carrier can  
35 be, for example, a liposome loadable with the antisense oligonucleotide.

According to a preferred embodiment of the present invention the antisense oligonucleotide further includes a ribozyme sequence. The ribozyme sequence serves to cleave a heparanase RNA molecule to which

the antisense oligonucleotide binds, to thereby downregulate heparanase expression.

Further according to the present invention there is provided an antisense nucleic acid construct comprising a promoter sequence and a polynucleotide sequence directing the synthesis of an antisense RNA sequence of at least 10 bases being hybridizable *in vivo*, under physiological conditions, with a portion of a polynucleotide strand encoding a polypeptide having heparanase catalytic activity. Like the antisense oligonucleotide, the antisense construct can be used for downregulating heparanase activity by *in vivo* administration thereof to a patient. As such, the antisense construct, like the antisense oligonucleotide, according to the present invention can be used to treat types of cancers which are characterized by impaired (over) expression of heparanase, and are dependent on the expression of heparanase for proliferating or forming metastases.

Thus, further according to the present invention there is provided a pharmaceutical composition comprising the antisense construct herein described and a pharmaceutically acceptable carrier. The carrier can be, for example, a liposome loadable with the antisense construct.

Formulations for topical administration may include, but are not limited to, lotions, ointments, gels, creams, suppositories, drops, liquids, sprays and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Coated condoms, stents, active pads, and other medical devices may also be useful. Compositions for oral administration include powders or granules, suspensions or solutions in water or non-aqueous media, sachets, capsules or tablets. Thickeners, diluents, flavorings, dispersing aids, emulsifiers or binders may be desirable. Formulations for parenteral administration may include, but are not limited to, sterile aqueous solutions which may also contain buffers, diluents and other suitable additives.

Dosing is dependent on severity and responsiveness of the condition to be treated, but will normally be one or more doses per day, week or month with course of treatment lasting from several days to several months or until a cure is effected or a diminution of disease state is achieved. Persons ordinarily skilled in the art can easily determine optimum dosages, dosing methodologies and repetition rates.

Further according to the present invention there is provided a nucleic acid construct comprising a polynucleotide sequence functioning as a promoter, the polynucleotide sequence is derived from SEQ ID NO:42 and

includes at least nucleotides 2135-2635, preferably 2235-2635, more preferably 2335-2635, more preferably 2435-2635, most preferably 2535-2635 thereof, or SEQ ID NO:43 and includes at least nucleotides 1-420, preferably 120-420, more preferably 220-420, most preferably 320-420, thereof. These nucleotides are shown in the example section that follows to direct the synthesis of a reporter gene in transformed cells. Thus, further according to the present invention there is provided a method of expressing a polynucleotide sequence comprising the step of ligating the polynucleotide sequence downstream to either of the promoter sequences described herein. Heparanase promoters can be isolated from a variety of mammalian and other species by cloning genomic regions present 5' to the coding sequence thereof. This can be readily achievable by one ordinarily skilled in the art using the heparanase polynucleotides described herein, which are shown in the Examples section that follows to participate in efficient cross species hybridization.

Further according to the present invention there is provided a recombinant protein comprising a polypeptide having heparanase catalytic activity. The protein according to the present invention include modifications known as post translational modifications, including, but not limited to, proteolysis (e.g., removal of a signal peptide and of a pro- or preprotein sequence), methionine modification, glycosylation, alkylation (e.g., methylation), acetylation, etc. According to preferred embodiments the polypeptide includes at least a portion of SEQ ID NOs:10, 14 or 44, the portion has heparanase catalytic activity. According to preferred embodiments of the present invention the protein is encoded by any of the above described isolated nucleic acids. Further according to the present invention there is provided a pharmaceutical composition comprising, as an active ingredient, the recombinant protein described herein.

The recombinant protein may be purified by any conventional protein purification procedure close to homogeneity and/or be mixed with additives. The recombinant protein may be manufactured using any of the genetically modified cells described above, which include any of the expression nucleic acid constructs described herein. The recombinant protein may be in any form. It may be in a crystallized form, a dehydrated powder form or in solution. The recombinant protein may be useful in obtaining pure heparanase, which in turn may be useful in eliciting anti-heparanase antibodies, either poly or monoclonal antibodies, and as a

screening active ingredient in an anti-heparanase inhibitors or drugs screening assay or system.

Further according to the present invention there is provided a method of identifying a chromosome region harboring a human heparanase gene in a chromosome spread. the method is executed implementing the following method steps, in which in a first step the chromosome spread (either interphase or metaphase spread) is hybridized with a tagged polynucleotide probe encoding heparanase. The tag is preferably a fluorescent tag. In a second step according to the method the chromosome spread is washed, thereby excess of non-hybridized probe is removed. Finally, signals associated with the hybridized tagged polynucleotide probe are searched for, wherein detected signals being indicative of a chromosome region harboring the human heparanase gene. One ordinarily skilled in the art would know how to use the sequences disclosed herein in suitable labeling reactions and how to use the tagged probes to detect, using *in situ* hybridization, a chromosome region harboring a human heparanase gene.

Further according to the present invention there is provided a method of *in vivo* eliciting anti-heparanase antibodies comprising the steps of administering a nucleic acid construct including a polynucleotide segment corresponding to at least a portion of SEQ ID NOs:9, 13 or 43 and a promoter for directing the expression of said polynucleotide segment *in vivo*. Accordingly, there is provided also a DNA vaccine for *in vivo* eliciting anti-heparanase antibodies comprising a nucleic acid construct including a polynucleotide segment corresponding to at least a portion of SEQ ID NOs:9, 13 or 43 and a promoter for directing the expression of said polynucleotide segment *in vivo*. The vaccine optionally further includes a pharmaceutically acceptable carrier, such as a virus, liposome or an antigen presenting cell. Alternatively, the vaccine is employed as a naked DNA vaccine

The present invention can be used to develop treatments for various diseases, to develop diagnostic assays for these diseases and to provide new tools for basic research especially in the fields of medicine and biology.

Specifically, the present invention can be used to develop new drugs to inhibit tumor cell metastasis, inflammation and autoimmunity. The identification of the *hpa* gene encoding for the heparanase enzyme enables the production of a recombinant enzyme in heterologous expression systems.

Furthermore, the present invention can be used to modulate bioavailability of heparin-binding growth factors, cellular responses to heparin-binding growth factors (e.g., bFGF, VEGF) and cytokines (e.g., IL-8), cell interaction with plasma lipoproteins, cellular susceptibility to viral, protozoa and some bacterial infections, and disintegration of neurodegenerative plaques. Recombinant heparanase offers a potential treatment for wound healing, angiogenesis, restenosis, atherosclerosis, inflammation, neurodegenerative diseases (such as, for example, Genstmann-Straussler Syndrome, Creutzfeldt-Jakob disease, Scrape and Alzheimer's disease) and certain viral and some bacterial and protozoa infections. Recombinant heparanase can be used to neutralize plasma heparin, as a potential replacement of protamine.

As used herein, the term "modulate" includes substantially inhibiting, slowing or reversing the progression of a disease, substantially ameliorating clinical symptoms of a disease or condition, or substantially preventing the appearance of clinical symptoms of a disease or condition. A "modulator" therefore includes an agent which may modulate a disease or condition. Modulation of viral, protozoa and bacterial infections includes any effect which substantially interrupts, prevents or reduces any viral, bacterial or protozoa activity and/or stage of the virus, bacterium or protozoon life cycle, or which reduces or prevents infection by the virus, bacterium or protozoon in a subject, such as a human or lower animal.

As used herein, the term "wound" includes any injury to any portion of the body of a subject including, but not limited to, acute conditions such as thermal burns, chemical burns, radiation burns, burns caused by excess exposure to ultraviolet radiation such as sunburn, damage to bodily tissues such as the perineum as a result of labor and childbirth, including injuries sustained during medical procedures such as episiotomies, trauma-induced injuries including cuts, those injuries sustained in automobile and other mechanical accidents, and those caused by bullets, knives and other weapons, and post-surgical injuries, as well as chronic conditions such as pressure sores, bedsores, conditions related to diabetes and poor circulation, and all types of acne, etc.

Anti-heparanase antibodies, raised against the recombinant enzyme, would be useful for immunodetection and diagnosis of micrometastases, autoimmune lesions and renal failure in biopsy specimens, plasma samples, and body fluids. Such antibodies may also serve as neutralizing agents for heparanase activity.

The genomic heparanase sequences described herein can be used to construct knock-in and knock-out constructs. Such constructs include a fragment of 10-20 Kb of a heparanase locus and a negative and a positive selection markers and can be used to provide heparanase knock-in and knock-out animal models by methods known to the skilled artisan. Such animal models can be used for studying the function of heparanase in developmental processes, and in normal as well as pathological processes. They can also serve as an experimental model for testing drugs and gene therapy protocols. The complementary heparanase sequence (cDNA) can be used to derive transgenic animals, overexpressing heparanase for same. Alternatively, if cloned in the antisense orientation, the complementary heparanase sequence (cDNA) can be used to derive transgenic animals under-expressing heparanase for same.

The heparanase promoter sequences described herein and other cis regulatory elements linked to the heparanase locus can be used to regulated the expression of genes. For example, these promoters can be used to direct the expression of a cytotoxic protein, such as TNF, in tumor cells. It will be appreciated that heparanase itself is abnormally expressed under the control of its own promoter and other cis acting elements in a variety of tumors, and its expression is correlated with metastasis. It is also abnormally highly expressed in inflammatory cells. The introns of the heparanase gene can be used for the same purpose, as it is known that introns, especially upstream introns include cis acting element which affect expression. A heparanase promoter fused to a reporter protein can be used to study/monitor its activity.

The polynucleotide sequences described herein can also be used to provide DNA vaccines which will elicit in vivo anti heparanase antibodies. Such vaccines can therefore be used to combat inflammatory and cancer.

Antisense oligonucleotides derived according to the heparanase sequences described herein, especially such oligonucleotides supplemented with ribozyme activity, can be used to modulate heparanase expression. Such oligonucleotides can be from the coding region, from the introns or promoter specific. Antisense heparanase nucleic acid constructs can similarly function, as well known in the art.

The heparanase sequences described herein can be used to study the catalytic mechanism of heparanase. Carefully selected site directed mutagenesis can be employed to provide modified heparanase proteins

having modified characteristics in terms of, for example, substrate specificity, sensitivity to inhibitors, etc.

While studying heparanase expression in a variety of cell types alternatively spliced transcripts were identified. Such transcripts if found  
5 characteristic of certain pathological conditions can be used as markers for such conditions. Such transcripts are expected to direct the synthesis of heparanases with altered functions.

Additional objects, advantages, and novel features of the present  
10 invention will become apparent to one ordinarily skilled in the art upon examination of the following examples, which are not intended to be limiting. Additionally, each of the various embodiments and aspects of the present invention as delineated hereinabove and as claimed in the claims section below finds experimental support in the following examples.

#### EXAMPLES

Generally, the nomenclature used herein and the laboratory  
procedures in recombinant DNA technology described below are those well known and commonly employed in the art. Standard techniques are used for  
20 cloning, DNA and RNA isolation, amplification and purification. Generally enzymatic reactions involving DNA ligase, DNA polymerase, restriction endonucleases and the like are performed according to the manufacturers' specifications. These techniques and various other techniques are generally performed according to Sambrook et al., Molecular Cloning--A Laboratory  
25 Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y. (1989), which is incorporated herein by reference. Other general references are provided throughout this document. The procedures therein are believed to be well known in the art and are provided for the convenience of the reader. All the information contained therein is incorporated herein by reference.

30 The following protocols and experimental details are referenced in the Examples that follow:

***Purification and characterization of heparanase from a human  
35 hepatoma cell line and human placenta:*** A human hepatoma cell line (Sk-hep-1) was chosen as a source for purification of a human tumor-derived heparanase. Purification was essentially as described in U.S. Pat. No. 5,362,641 to Fuks, which is incorporated by reference as if fully set forth

herein. Briefly, 500 liter,  $5 \times 10^{11}$  cells were grown in suspension and the heparanase enzyme was purified about 240,000 fold by applying the following steps: (i) cation exchange (CM-Sephadex) chromatography performed at pH 6.0, 0.3-1.4 M NaCl gradient; (ii) cation exchange (CM-Sephadex) chromatography performed at pH 7.4 in the presence of 0.1% CHAPS, 0.3-1.1 M NaCl gradient; (iii) heparin-Sepharose chromatography performed at pH 7.4 in the presence of 0.1% CHAPS, 0.35-1.1 M NaCl gradient; (iv) ConA-Sepharose chromatography performed at pH 6.0 in buffer containing 0.1 % CHAPS and 1 M NaCl, elution with 0.25 M  $\alpha$ -methyl mannoside; and (v) HPLC cation exchange (Mono-S) chromatography performed at pH 7.4 in the presence of 0.1 % CHAPS, 0.25-1 M NaCl gradient.

Active fractions were pooled, precipitated with TCA and the precipitate subjected to SDS polyacrylamide gel electrophoresis and/or tryptic digestion and reverse phase HPLC. Tryptic peptides of the purified protein were separated by reverse phase HPLC (C8 column) and homogeneous peaks were subjected to amino acid sequence analysis.

The purified enzyme was applied to reverse phase HPLC and subjected to N-terminal amino acid sequencing using the amino acid sequencer (Applied Biosystems).

**Cells:** Cultures of bovine corneal endothelial cells (BCECs) were established from steer eyes as previously described (19, 38). Stock cultures were maintained in DMEM (1 g glucose/liter) supplemented with 10 % newborn calf serum and 5 % FCS. bFGF (1 ng/ml) was added every other day during the phase of active cell growth (13, 14).

**Preparation of dishes coated with ECM:** BCECs (second to fifth passage) were plated into 4-well plates at an initial density of  $2 \times 10^5$  cells/ml, and cultured in sulfate-free Fisher medium plus 5 % dextran T-40 for 12 days.  $\text{Na}_2^{35}\text{SO}_4$  (25  $\mu\text{Ci/ml}$ ) was added on day 1 and 5 after seeding and the cultures were incubated with the label without medium change. The subendothelial ECM was exposed by dissolving (5 min., room temperature) the cell layer with PBS containing 0.5 % Triton X-100 and 20 mM  $\text{NH}_4\text{OH}$ , followed by four washes with PBS. The ECM remained intact, free of cellular debris and firmly attached to the entire area of the tissue culture dish (19, 22).

To prepare soluble sulfate labeled proteoglycans (peak I material), the ECM was digested with trypsin (25  $\mu\text{g/ml}$ , 6 h, 37 °C), the digest was concentrated by reverse dialysis and the concentrated material was applied



onto a Sepharose 6B gel filtration column. The resulting high molecular weight material ( $K_{av} < 0.2$ , peak I) was collected. More than 80 % of the labeled material was shown to be composed of heparan sulfate proteoglycans (11, 39).

5       **Heparanase activity:** Cells ( $1 \times 10^6$ /35-mm dish), cell lysates or conditioned media were incubated on top of  $^{35}\text{S}$ -labeled ECM (18 h,  $37^\circ\text{C}$ ) in the presence of 20 mM phosphate buffer (pH 6.2). Cell lysates and conditioned media were also incubated with sulfate labeled peak I material (10-20  $\mu\text{l}$ ). The incubation medium was collected, centrifuged (18,000  $\times g$ ,  
10     $4^\circ\text{C}$ , 3 min.), and sulfate labeled material analyzed by gel filtration on a Sepharose CL-6B column (0.9  $\times$  30 cm). Fractions (0.2 ml) were eluted with PBS at a flow rate of 5 ml/h and counted for radioactivity using Bio-fluor scintillation fluid. The excluded volume ( $V_o$ ) was marked by blue dextran and the total included volume ( $V_t$ ) by phenol red. The latter was  
15    shown to comigrate with free sulfate (7, 11, 23). Degradation fragments of HS side chains were eluted from Sepharose 6B at  $0.5 < K_{av} < 0.8$  (peak II) (7, 11, 23). A nearly intact HSPG released from ECM by trypsin - and, to a lower extent, during incubation with PBS alone - was eluted next to  $V_o$  ( $K_{av} < 0.2$ , peak I). Recoveries of labeled material applied on the columns  
20    ranged from 85 to 95 % in different experiments (11). Each experiment was performed at least three times and the variation of elution positions ( $K_{av}$  values) did not exceed  $\pm 15\%$ .

**Cloning of hpa cDNA:** cDNA clones 257548 and 260138 were obtained from the I.M.A.G.E Consortium (2130 Memorial Parkway SW,  
25    Huntsville, AL 35801). The cDNAs were originally cloned in *EcoRI* and *NotI* cloning sites in the plasmid vector pT3T7D-Pac. Although these clones are reported to be somewhat different, DNA sequencing demonstrated that these clones are identical to one another. Marathon RACE (rapid amplification of cDNA ends) human placenta (poly-A) cDNA  
30    composite was a gift of Prof. Yossi Shiloh of Tel Aviv University. This composite is vector free, as it includes reverse transcribed cDNA fragments to which double, partially single stranded adapters are attached on both sides. The construction of the specific composite employed is described in reference 39a.

35       Amplification of hp3 PCR fragment was performed according to the protocol provided by Clontech laboratories. The template used for amplification was a sample taken from the above composite. The primers used for amplification were:

First step: 5'-primer: AP1: 5'-CCATCCTAATACGACTCACT  
ATAGGGC-3', SEQ ID NO:1; 3'-primer: HPL229: 5'-GTAGTGATGCCA  
TGTAAGTGAATC-3', SEQ ID NO:2.

Second step: nested 5'-primer: AP2: 5'-ACTCACTATAGGGCTCG  
5 AGCGGC-3', SEQ ID NO:3; nested 3'- primer: HPL171: 5'-  
GCATCTTAGCCGTCTTTCTTCG-3', SEQ ID NO:4. The HPL229 and  
HPL171 were selected according to the sequence of the EST clones. They  
include nucleotides 933-956 and 876-897 of SEQ ID NO:9, respectively.

PCR program was 94 °C - 4 min., followed by 30 cycles of 94 °C -  
10 40 sec., 62 °C - 1 min., 72 °C - 2.5 min. Amplification was performed with  
Expand High Fidelity (Boehringer Mannheim). The resulting ca. 900 bp  
hp3 PCR product was digested with *Bfr*I and *Pvu*II. Clone 257548 (*phpa*1)  
was digested with *Eco*RI, followed by end filling and was then further  
digested with *Bfr*I. Thereafter the *Pvu*II - *Bfr*I fragment of the hp3 PCR  
15 product was cloned into the blunt end - *Bfr*I end of clone *phpa*1 which  
resulted in having the entire cDNA cloned in pT3T7-pac vector, designated  
*phpa*2.

**RT-PCR:** RNA was prepared using TRI-Reagent (Molecular  
research center Inc.) according to the manufacturer instructions. 1.25 µg  
20 were taken for reverse transcription reaction using MuMLV Reverse  
transcriptase (Gibco BRL) and Oligo (dT)<sub>15</sub> primer, SEQ ID NO:5,  
(Promega). Amplification of the resultant first strand cDNA was  
performed with *Taq* polymerase (Promega). The following primers were  
used:

25 HPU-355: 5'-TTCGATCCCAAGAAGGAATCAAC-3', SEQ ID NO:6,  
nucleotides 372-394 in SEQ ID NOs:9 or 11.  
HPL-229: 5'-GTAGTGATGCCATGTAAGTGAATC-3', SEQ ID NO:7,  
nucleotides 933-956 in SEQ ID NOs:9 or 11.

PCR program: 94 °C - 4 min., followed by 30 cycles of 94 °C - 40  
30 sec., 62 °C - 1 min., 72 °C - 1 min.

Alternatively, total RNA was prepared from cell cultures using Tri-  
reagent (Molecular Research Center, Inc.) according to the manufacturer  
recommendation. Poly A+ RNA was isolated from total RNA using mRNA  
separator (Clontech). Reverse transcription was performed with total RNA  
35 using Superscript II (GibcoBRL). PCR was performed with Expand high  
fidelity (Boehringer Mannheim). Primers used for amplification were as  
follows:

Hpu-685, 5'-GAGCAGCCAGGTGAGCCCAAGAT-3', SEQ ID NO:24

Hpu-355, 5'-TTCGATCCCAAGAAGGAATCAAC-3', SEQ ID NO:25

Hpu 565, 5'-AGCTCTGTAGATGTGCTATACAC-3', SEQ ID NO:26

Hpl 967, 5'-TCAGATGCAAGCAGCAACTTTGGC-3', SEQ ID NO:27

Hpl 171, 5'-GCATCTTAGCCGTCTTTCTTCG-3', SEQ ID NO:28

5 Hpl 229, 5'-GTAGTGATGCCATGTAAGTGAATC-3', SEQ ID NO:29

PCR reaction was performed as follows: 94 °C 3 minutes, followed by 32 cycles of 94 °C 40 seconds, 64 °C 1 minute, 72 °C 3 minutes, and one cycle 72 °C, 7 minutes.

**Expression of recombinant heparanase in insect cells:** Cells, High Five and Sf21 insect cell lines were maintained as monolayer cultures in SF900II-SFM medium (GibcoBRL).

**Recombinant Baculovirus:** Recombinant virus containing the *hpa* gene was constructed using the Bac to Bac system (GibcoBRL). The transfer vector pFastBac was digested with *SaI* and *NotI* and ligated with a 1.7 kb fragment of *phpa2* digested with *XhoI* and *NotI*. The resulting plasmid was designated pFast*hpa2*. An identical plasmid designated pFast*hpa4* was prepared as a duplicate and both independently served for further experimentations. Recombinant bacmid was generated according to the instructions of the manufacturer with pFast*hpa2*, pFast*hpa4* and with pFastBac. The latter served as a negative control. Recombinant bacmid DNAs were transfected into Sf21 insect cells. Five days after transfection recombinant viruses were harvested and used to infect High Five insect cells,  $3 \times 10^6$  cells in T-25 flasks. Cells were harvested 2 - 3 days after infection.  $4 \times 10^6$  cells were centrifuged and resuspended in a reaction buffer containing 20 mM phosphate citrate buffer, 50 mM NaCl. Cells underwent three cycles of freeze and thaw and lysates were stored at -80 °C. Conditioned medium was stored at 4 °C.

**Partial purification of recombinant heparanase:** Partial purification of recombinant heparanase was performed by heparin-Sepharose column chromatography followed by Superdex 75 column gel filtration. Culture medium (150 ml) of Sf21 cells infected with pFhpa4 virus was subjected to heparin-Sepharose chromatography. Elution of 1 ml fractions was performed with 0.35 - 2 M NaCl gradient in presence of 0.1 % CHAPS and 1 mM DTT in 10 mM sodium acetate buffer, pH 5.0. A 25 µl sample of each fraction was tested for heparanase activity. Heparanase activity was eluted at the range of 0.65 - 1.1 M NaCl (fractions 18-26, Figure 10a). 5 µl of each fraction was subjected to 15 % SDS-polyacrylamide gel electrophoresis followed by silver nitrate staining.

Active fractions eluted from heparin-Sepharose (Figure 10a) were pooled and concentrated (x 6) on YM3 cut-off membrane. 0.5 ml of the concentrated material was applied onto 30 ml Superdex 75 FPLC column equilibrated with 10 mM sodium acetate buffer, pH 5.0, containing 0.8 M NaCl, 1 mM DTT and 0.1 % CHAPS. Fractions (0.56 ml) were collected at a flow rate of 0.75 ml/min. Aliquots of each fraction were tested for heparanase activity and were subjected to SDS-polyacrylamide gel electrophoresis followed by silver nitrate staining (Figure 11b).

**PCR amplification of genomic DNA:** 94 °C 3 minutes, followed by 32 cycles of 94 °C 45 seconds, 64 °C 1 minute, 68 °C 5 minutes, and one cycle at 72 °C, 7 minutes. Primers used for amplification of genomic DNA included:

GHpu-L3 5'-AGGCACCCTAGAGATGTTCCAG-3', SEQ ID NO:30

GHpl-L6 5'-GAAGATTTCTGTTTCCATGACGTG-3', SEQ ID NO:31.

**Screening of genomic libraries:** A human genomic library in Lambda phage EMBLE3 SP6/T7 (Clontech, Paulo Alto, CA) was screened. 5 x 10<sup>5</sup> plaques were plated at 5 x 10<sup>4</sup> pfu/plate on NZCYM agar/top agarose plates. Phages were absorbed on nylon membranes in duplicates (Qiagen). Hybridization was performed at 65 °C in 5 x SSC, 5 x Denhart's, 10 % dextran sulfate, 100 µg/ml Salmon sperm, <sup>32</sup>p labeled probe (10<sup>6</sup> cpm/ml). A 1.6 kb fragment, containing the entire *hpa* cDNA was labeled by random priming (Boehringer Mannheim). Following hybridization membranes were washed once with 2 x SSC, 0.1 % SDS at 65 °C for 20 minutes, and twice with 0.2 x SSC, 0.1 % SDS at 65 °C for 15 minutes. Hybridizing plaques were picked, and plated at 100 pfu/plate. Hybridization was performed as above and single isolated positive plaques were picked.

Phage DNA was extracted using a Lambda DNA extraction kit (Qiagen). DNA was digested with *Xho*I and *Eco*RI, separated on 0.7 % agarose gel and transferred to nylon membrane Hybond N+ (Amersham). Hybridization and washes were performed as above.

**cDNA Sequence analysis:** Sequence determinations were performed with vector specific and gene specific primers, using an automated DNA sequencer (Applied Biosystems, model 373A). Each nucleotide was read from at least two independent primers.

**Genomic sequence analysis:** Large-scale sequencing was performed by Commonwealth Biotechnology Incorporation.

**Isolation of mouse *hpa*:** Mouse *hpa* cDNA was amplified from either Marathon ready cDNA library of mouse embryo or from mRNA isolated from mouse melanoma cell line BL6, using the Marathon RACE kit from Clontech. Both procedures were performed according to the manufacturer's recommendation.

**Primers used for PCR amplification of mouse *hpa*:**

- Mhpl773 5'-CCACACTGAATGTAATACTGAAGTG-3', SEQ ID NO:32  
 MHpl736 5'-CGAAGCTCTGGAACCTCGGCAAG-3', SEQ ID NO:33  
 MHpl83 5'-GCCAGCTGCAAAGGTGTTGGAC-3', SEQ ID NO:34  
 10 Mhpl152 5'-AACACCTGCCTCATCACGACTTC-3', SEQ ID NO:35  
 Mhpl114 5'-GCCAGGCTGGCGTCGATGGTGA-3', SEQ ID NO:36  
 MHpl103 5'-GTCGATGGTGATGGACAGGAAC-3', SEQ ID NO:37  
 Apl 5'-GTAATACGACTCACTATAGGGC-3', SEQ ID NO:38 -  
 (Genome walker)  
 15 Ap2 5'-ACTATAGGGCACGCGTGGT-3', SEQ ID NO:39 -  
 (Genome walker)  
 Apl 5'-CCATCCTAATACGACTCACTATAGGGC-3', SEQ ID NO:40 -  
 (Marathon RACE)  
 Ap2 5'-ACTCACTATAGGGCTCGAGCGGC-3', SEQ ID NO:41 -  
 20 (Marathon RACE)

**Southern analysis of genomic DNA:** Genomic DNA was extracted from animal or from human blood using Blood and cell culture DNA maxi kit (Qiagene). DNA was digested with *EcoRI*, separated by gel electrophoresis and transferred to a nylon membrane Hybond N+ (Amersham). Hybridization was performed at 68 °C in 6 x SSC, 1 % SDS, 5 x Denharts, 10 % dextran sulfate, 100 µg/ml salmon sperm DNA, and <sup>32</sup>p labeled probe. A 1.6 kb fragment, containing the entire *hpa* cDNA was used as a probe. Following hybridization, the membrane was washed with 3 x SSC, 0.1 % SDS, at 68 °C and exposed to X-ray film for 3 days. Membranes were then washed with 1 x SSC, 0.1 % SDS, at 68 °C and were reexposed for 5 days.

**Construction of *hpa* promoter-GFP expression vector:** Lambda DNA of phage L3, was digested with *SacI* and *BglII*, resulting in a 1712 bp fragment which contained the *hpa* promoter (877-2688 of SEQ ID NO:42). The pEGFP-1 plasmid (Clontech) was digested with *BglII* and *SacI* and ligated with the 1712 bp fragment of the *hpa* promoter sequence. The resulting plasmid was designated phpEGL. A second *hpa* promoter-GFP plasmid was constructed containing a shorter fragment of the *hpa* promoter

region: phpEGL was digested with *HindIII*, and the resulting 1095 bp fragment (nucleotides 1593-2688 of SEQ ID NO:42) was ligated with *HindIII* digested pEGFP-1. The resulting plasmid was designated phpEGS.

**Computer analysis of sequences:** Homology searches were performed using several computer servers, and various databases. Blast 2.0 service, at the NCBI server was used to screen the protein database swplus and DNA databases such as GenBank, EMBL, and the EST databases. Blast 2.0 search was performed using the basic search option of the NCBI server. Sequence analysis and alignments were done using the DNA sequence analysis software package developed by the Genetic Computer Group (GCG) at the university of Wisconsin. Alignments of two sequences were performed using Bestfit (gap creation penalty - 12, gap extension penalty - 4). Protein homology search was performed with the Smith-Waterman algorithm, using the Bioaccelerator platform developed by Compugene. The protein database swplus was searched using the following parameters: gapop: 10.0, gapext: 0.5, matrix: blosum62. Blocks homology was performed using the Blocks WWW server developed at Fred Hutchinson Cancer Research Center in Seattle, Washington, USA. Secondary structure prediction was performed using the PHD server - Profile network Prediction Heidelberg. Fold recognition (threading) was performed using the UCLA-DOE structure prediction server. The method used for prediction was gonnet+predss. Alignment of three sequences was performed using the pileup application (gap creation penalty - 5, gap extension penalty - 1). Promoter analysis was performed using TSSW and TSSG programs (BCM Search Launcher Human Genome Center, Baylor College of Medicine, Houston TX).

## EXAMPLE 1

### *Cloning of human hpa cDNA*

Purified fraction of heparanase isolated from human hepatoma cells (SK-hep-1) was subjected to tryptic digestion and microsequencing. EST (Expressed Sequence Tag) databases were screened for homology to the back translated DNA sequences corresponding to the obtained peptides. Two EST sequences (accession Nos. N41349 and N45367) contained a DNA sequence encoding the peptide YGPDVGQPR (SEQ ID NO:8). These two sequences were derived from clones 257548 and 260138 (I.M.A.G.E Consortium) prepared from 8 to 9 weeks placenta cDNA library (Soares). Both clones which were found to be identical contained an insert

of 1020 bp which included an open reading frame (ORF) of 973 bp followed by a 3' untranslated region of 27 bp and a Poly A tail. No translation start site (AUG) was identified at the 5' end of these clones.

Cloning of the missing 5' end was performed by PCR amplification of DNA from a placenta Marathon RACE cDNA composite. A 900 bp fragment (designated hp3), partially overlapping with the identified 3' encoding EST clones was obtained.

The joined cDNA fragment, 1721 bp long (SEQ ID NO:9), contained an open reading frame which encodes, as shown in Figure 1 and SEQ ID NO:11, a polypeptide of 543 amino acids (SEQ ID NO:10) with a calculated molecular weight of 61,192 daltons. The 3' end of the partial cDNA inserts contained in clones 257548 and 260138 started at nucleotide G<sup>721</sup> of SEQ ID NO:9 and Figure 1.

As further shown in Figure 1, there was a single sequence discrepancy between the EST clones and the PCR amplified sequence, which led to an amino acid substitution from Tyr<sup>246</sup> in the EST to Phe<sup>246</sup> in the amplified cDNA. The nucleotide sequence of the PCR amplified cDNA fragment was verified from two independent amplification products. The new gene was designated *hpa*.

As stated above, the 3' end of the partial cDNA inserts contained in EST clones 257548 and 260138 started at nucleotide 721 of *hpa* (SEQ ID NO:9). The ability of the *hpa* cDNA to form stable secondary structures, such as stem and loop structures involving nucleotide stretches in the vicinity of position 721 was investigated using computer modeling. It was found that stable stem and loop structures are likely to be formed involving nucleotides 698-724 (SEQ ID NO:9). In addition, a high GC content, up to 70 %, characterizes the 5' end region of the *hpa* gene, as compared to about only 40 % in the 3' region. These findings may explain the immature termination and therefore lack of 5' ends in the EST clones.

To examine the ability of the *hpa* gene product to catalyze degradation of heparan sulfate in an *in vitro* assay the entire open reading frame was expressed in insect cells, using the Baculovirus expression system. Extracts of cells, infected with virus containing the *hpa* gene, demonstrated a high level of heparan sulfate degradation activity, while cells infected with a similar construct containing no *hpa* gene had no such activity, nor did non-infected cells. These results are further demonstrated in the following Examples.

**EXAMPLE 2*****Degradation of soluble ECM-derived HSPG***

Monolayer cultures of High Five cells were infected (72 h, 28 °C) with recombinant Baculovirus containing the pFast*hpa* plasmid or with control virus containing an insert free plasmid. The cells were harvested and lysed in heparanase reaction buffer by three cycles of freezing and thawing. The cell lysates were then incubated (18 h, 37 °C) with sulfate labeled, ECM-derived HSPG (peak I), followed by gel filtration analysis (Sephacrose 6B) of the reaction mixture.

As shown in Figure 2, the substrate alone included almost entirely high molecular weight (Mr) material eluted next to V<sub>0</sub> (peak I, fractions 5-20, Kav < 0.35). A similar elution pattern was obtained when the HSPG substrate was incubated with lysates of cells that were infected with control virus. In contrast, incubation of the HSPG substrate with lysates of cells infected with the *hpa* containing virus resulted in a complete conversion of the high Mr substrate into low Mr labeled degradation fragments (peak II, fractions 22-35, 0.5 < Kav < 0.75).

Fragments eluted in peak II were shown to be degradation products of heparan sulfate, as they were (i) 5- to 6-fold smaller than intact heparan sulfate side chains (Kav approx. 0.33) released from ECM by treatment with either alkaline borohydride or papain; and (ii) resistant to further digestion with papain or chondroitinase ABC, and susceptible to deamination by nitrous acid (6, 11). Similar results (not shown) were obtained with Sf21 cells. Again, heparanase activity was detected in cells infected with the *hpa* containing virus (pF*hpa*), but not with control virus (pF). This result was obtained with two independently generated recombinant viruses. Lysates of control not infected High Five cells failed to degrade the HSPG substrate.

In subsequent experiments, the labeled HSPG substrate was incubated with medium conditioned by infected High Five or Sf21 cells.

As shown in Figures 3a-b, heparanase activity, reflected by the conversion of the high Mr peak I substrate into the low Mr peak II which represents HS degradation fragments, was found in the culture medium of cells infected with the pF*hpa2* or pF*hpa4* viruses, but not with the control pF1 or pF2 viruses. No heparanase activity was detected in the culture medium of control non-infected High Five or Sf21 cells.

The medium of cells infected with the pF*hpa4* virus was passed through a 50 kDa cut off membrane to obtain a crude estimation of the



molecular weight of the recombinant heparanase enzyme. As demonstrated in Figure 4, all the enzymatic activity was retained in the upper compartment and there was no activity in the flow through (<50 kDa) material. This result is consistent with the expected molecular weight of the *hpa* gene product.

In order to further characterize the *hpa* product the inhibitory effect of heparin, a potent inhibitor of heparanase mediated HS degradation (40) was examined.

As demonstrated in Figures 5a-b, conversion of the peak I substrate into peak II HS degradation fragments was completely abolished in the presence of heparin.

Altogether, these results indicate that the heparanase enzyme is expressed in an active form by insect cells infected with Baculovirus containing the newly identified human *hpa* gene.

### EXAMPLE 3

#### *Degradation of HSPG in intact ECM*

Next, the ability of intact infected insect cells to degrade HS in intact, naturally produced ECM was investigated. For this purpose, High Five or Sf21 cells were seeded on metabolically sulfate labeled ECM followed by infection (48 h, 28 °C) with either the pF*hpa4* or control pF2 viruses. The pH of the medium was then adjusted to pH 6.2-6.4 and the cells further incubated with the labeled ECM for another 48 h at 28 °C or 24 h at 37 °C. Sulfate labeled material released into the incubation medium was analyzed by gel filtration on Sepharose 6B.

As shown in Figures 6a-b and 7a-b, incubation of the ECM with cells infected with the control pF2 virus resulted in a constant release of labeled material that consisted almost entirely (>90%) of high Mr fragments (peak I) eluted with or next to  $V_0$ . It was previously shown that a proteolytic activity residing in the ECM itself and/or expressed by cells is responsible for release of the high Mr material (6). This nearly intact HSPG provides a soluble substrate for subsequent degradation by heparanase, as also indicated by the relatively large amount of peak I material accumulating when the heparanase enzyme is inhibited by heparin (6, 7, 12, Figure 9). On the other hand, incubation of the labeled ECM with cells infected with the pF*hpa4* virus resulted in release of 60-70% of the ECM-associated radioactivity in the form of low Mr sulfate-labeled fragments (peak II,  $0.5 < K_{av} < 0.75$ ), regardless of whether the infected cells were incubated with

the ECM at 28 °C or 37 °C. Control intact non-infected Sf21 or High Five cells failed to degrade the ECM HS side chains.

In subsequent experiments, as demonstrated in Figures 8a-b, High Five and Sf21 cells were infected (96 h, 28 °C) with pFhpa4 or control pF1 viruses and the culture medium incubated with sulfate-labeled ECM. Low Mr HS degradation fragments were released from the ECM only upon incubation with medium conditioned by pFhpa4 infected cells. As shown in Figure 9, production of these fragments was abolished in the presence of heparin. No heparanase activity was detected in the culture medium of control, non-infected cells. These results indicate that the heparanase enzyme expressed by cells infected with the pFhpa4 virus is capable of degrading HS when complexed to other macromolecular constituents (i.e. fibronectin, laminin, collagen) of a naturally produced intact ECM, in a manner similar to that reported for highly metastatic tumor cells or activated cells of the immune system (6, 7).

#### EXAMPLE 4

##### *Purification of recombinant human heparanase*

The recombinant heparanase was partially purified from medium of pFhpa4 infected Sf21 cells by Heparin-Sepharose chromatography (Figure 10a) followed by gel filtration of the pooled active fractions over an FPLC Superdex 75 column (Figure 11a). A ~ 63 kDa protein was observed, whose quantity, as was detected by silver stained SDS-polyacrylamide gel electrophoresis, correlated with heparanase activity in the relevant column fractions (Figures 10b and 11b, respectively). This protein was not detected in the culture medium of cells infected with the control pF1 virus and was subjected to a similar fractionation on heparin-Sepharose (not shown).

#### EXAMPLE 5

##### *Expression of the human hpa cDNA in various cell types, organs and tissues*

Referring now to Figures 12a-e, RT-PCR was applied to evaluate the expression of the hpa gene by various cell types and tissues. For this purpose, total RNA was reverse transcribed and amplified. The expected 585 bp long cDNA was clearly demonstrated in human kidney, placenta (8 and 11 weeks) and mole tissues, as well as in freshly isolated and short termed (1.5-48 h) cultured human placental cytotrophoblastic cells (Figure 12a), all known to express a high heparanase activity (41). The hpa

transcript was also expressed by normal human neutrophils (Figure 12b). In contrast, there was no detectable expression of the *hpa* mRNA in embryonic human muscle tissue, thymus, heart and adrenal (Figure 12b). The *hpa* gene was expressed by several, but not all, human bladder carcinoma cell lines (Figure 12c), SK hepatoma (SK-hep-1), ovarian carcinoma (OV 1063), breast carcinoma (435, 231), melanoma and megakaryocytic (DAMI, CHRF) human cell lines (Figures 12d-e).

The above described expression pattern of the *hpa* transcript was determined to be in a very good correlation with heparanase activity levels determined in various tissues and cell types (not shown).

### EXAMPLE 6

#### *Isolation of an extended 5' end of hpa cDNA from human SK-hep1 cell line*

The 5' end of *hpa* cDNA was isolated from human SK-hep1 cell line by PCR amplification using the Marathon RACE (rapid amplification of cDNA ends) kit (Clontech). Total RNA was prepared from SK-hep1 cells using the TRI-Reagent (Molecular research center Inc.) according to the manufacturer instructions. Poly A<sup>+</sup> RNA was isolated using the mRNA separator kit (Clonetech).

The Marathon RACE SK-hep1 cDNA composite was constructed according to the manufacturer recommendations. First round of amplification was performed using an adaptor specific primer AP1: 5'-CCATCCTAATACG ACTCACTATAGGGC-3', SEQ ID NO:1, and a *hpa* specific antisense primer hpl-629: 5'-CCCCAGGAGCAGCAGCATCAG-3', SEQ ID NO:17, corresponding to nucleotides 119-99 of SEQ ID NO:9. The resulting PCR product was subjected to a second round of amplification using an adaptor specific nested primer AP2: 5'-ACTCACTATAGGGCTCGAGCGGC-3', SEQ ID NO:3, and a *hpa* specific antisense nested primer hpl-666 5'-AGGCTTCGAGCGCAGCAGCAT-3', SEQ ID NO:18, corresponding to nucleotides 83-63 of SEQ ID NO:9. The PCR program was as follows: a hot start of 94 °C for 1 minute, followed by 30 cycles of 90 °C - 30 seconds, 68 °C - 4 minutes. The resulting 300 bp DNA fragment was extracted from an agarose gel and cloned into the vector pGEM-T Easy (Promega). The resulting recombinant plasmid was designated pHPSK1.

The nucleotide sequence of the pHPSK1 insert was determined and it was found to contain 62 nucleotides of the 5' end of the placenta *hpa* cDNA

(SEQ ID NO:9) and additional 178 nucleotides upstream, the first 178 nucleotides of SEQ ID NOs:13 and 15.

A single nucleotide discrepancy was identified between the SK-hep1 cDNA and the placenta cDNA. The "T" derivative at position 9 of the placenta cDNA (SEQ ID NO:9), is replaced by a "C" derivative at the corresponding position 187 of the SK-hep1 cDNA (SEQ ID NO:13).

The discrepancy is likely to be due to a mutation at the 5' end of the placenta cDNA clone as confirmed by sequence analysis of several additional cDNA clones isolated from placenta, which like the SK-hep1 cDNA contained C at position 9 of SEQ ID NO:9.

The 5' extended sequence of the SK-hep1 *hpa* cDNA was assembled with the sequence of the *hpa* cDNA isolated from human placenta (SEQ ID NO:9). The assembled sequence contained an open reading frame which encodes, as shown in SEQ ID NOs:14 and 15, a polypeptide of 592 amino acids with a calculated molecular weight of 66,407 daltons. The open reading frame is flanked by 93 bp 5' untranslated region (UTR).

#### EXAMPLE 7

##### *Isolation of the upstream genomic region of the hpa gene*

The upstream region of the *hpa* gene was isolated using the Genome Walker kit (Clontech) according to the manufacturer recommendations. The kit includes five human genomic DNA samples each digested with a different restriction endonuclease creating blunt ends: *EcoRV*, *ScaI*, *DraI*, *PvuII* and *SspI*.

The blunt ended DNA fragments are ligated to partially single stranded adaptors. The Genomic DNA samples were subjected to PCR amplification using the adaptor specific primer and a gene specific primer. Amplification was performed with Expand High Fidelity (Boehringer Mannheim).

A first round of amplification was performed using the ap1 primer: 5'-G TAATACGACTCACTATAGGGC-3', SEQ ID NO:19, and the *hpa* specific antisense primer hpl-666: 5'-AGGCTTCGAGCGCAGCAGCAT-3', SEQ ID NO:18, corresponding to nucleotides 83 – 63 of SEQ ID NO:9. The PCR program was as follows: a hot start of 94 °C - 3 minutes, followed by 36 cycles of 94 °C - 40 seconds, 67 °C - 4 minutes.

The PCR products of the first amplification were diluted 1:50. One µl of the diluted sample was used as a template for a second amplification using a nested adaptor specific primer ap2: 5'-

ACTATAGGGGCACGCGTGGT-3', SEQ ID NO:20, and a *hpa* specific antisense primer hpl-690, 5'-CTTGGGCTCACC TGGCTGCTC-3', SEQ ID NO:21, corresponding to nucleotides 62-42 of SEQ ID NO:9. The resulting amplification products were analyzed using agarose gel electrophoresis. Five different PCR products were obtained from the five amplification reactions. A DNA fragment of approximately 750 bp which was obtained from the *SspI* digested DNA sample was gel extracted. The purified fragment was ligated into the plasmid vector pGEM-T Easy (Promega). The resulting recombinant plasmid was designated pGHP6905 and the nucleotide sequence of the *hpa* insert was determined.

A partial sequence of 594 nucleotides is shown in SEQ ID NO:16. The last nucleotide in SEQ ID NO:13 corresponds to nucleotide 93 in SEQ ID:13. The DNA sequence in SEQ ID NO:16 contains the 5' region of the *hpa* cDNA and 501 nucleotides of the genomic upstream region which are predicted to contain the promoter region of the *hpa* gene.

#### EXAMPLE 8

##### *Expression of the 592 amino acids HPA polypeptide in a human 293 cell line*

The 592 amino acids open reading frame (SEQ ID NOs:13 and 15) was constructed by ligation of the 110 bp corresponding to the 5' end of the SK-hep1 *hpa* cDNA with the placenta cDNA. More specifically the Marathon RACE - PCR amplification product of the placenta *hpa* DNA was digested with *SacI* and an approximately 1 kb fragment was ligated into a *SacI*-digested pGHP6905 plasmid. The resulting plasmid was digested with *EarI* and *AatII*. The *EarI* sticky ends were blunted and an approximately 280 bp *EarI*/blunt-*AatII* fragment was isolated. This fragment was ligated with pFast*hpa* digested with *EcoRI* which was blunt ended using Klenow fragment and further digested with *AatII*. The resulting plasmid contained a 1827 bp insert which includes an open reading frame of 1776 bp, 31 bp of 3' UTR and 21 bp of 5' UTR. This plasmid was designated pFast*Lhpa*.

A mammalian expression vector was constructed to drive the expression of the 592 amino acids heparanase polypeptide in human cells. The *hpa* cDNA was excised from pFast*Lhpa* with *BssHII* and *NotI*. The resulting 1850 bp *BssHII*-*NotI* fragment was ligated to a mammalian expression vector pSI (Promega) digested with *MluI* and *NotI*. The resulting recombinant plasmid, pSI*hpa*Met2 was transfected into a human 293 embryonic kidney cell line.

Transient expression of the 592 amino-acids heparanase was examined by western blot analysis and the enzymatic activity was tested using the gel shift assay. Both these procedures are described in length in U.S. Pat. application No. 09/071,739, filed May 1, 1998, which is incorporated by reference as if fully set forth herein. Cells were harvested 3 days following transfection. Harvested cells were re-suspended in lysis buffer containing 150 mM NaCl, 50 mM Tris pH 7.5, 1% Triton X-100, 1 mM PMSF and protease inhibitor cocktail (Boehringer Mannheim). 40 µg protein extract samples were used for separation on a SDS-PAGE. Proteins were transferred onto a PVDF Hybond-P membrane (Amersham). The membrane was incubated with an affinity purified polyclonal anti heparanase antibody, as described in U.S. Pat. application No. 09/071,739. A major band of approximately 50 kDa was observed in the transfected cells as well as a minor band of approximately 65 kDa. A similar pattern was observed in extracts of cells transfected with the pShpa as demonstrated in U.S. Pat. application No. 09/071,739. These two bands probably represent two forms of the recombinant heparanase protein produced by the transfected cells. The 65 kDa protein probably represents a heparanase precursor, while the 50 kDa protein is suggested herein to be the processed or mature form.

The catalytic activity of the recombinant protein expressed in the pShpaMet2 transfected cells was tested by gel shift assay. Cell extracts of transfected and of mock transfected cells were incubated overnight with heparin (6 µg in each reaction) at 37 °C, in the presence of 20 mM phosphate citrate buffer pH 5.4, 1 mM CaCl<sub>2</sub>, 1 mM DTT and 50 mM NaCl. Reaction mixtures were then separated on a 10 % polyacrylamide gel. The catalytic activity of the recombinant heparanase was clearly demonstrated by a faster migration of the heparin molecules incubated with the transfected cell extract as compared to the control. Faster migration indicates the disappearance of high molecular weight heparin molecules and the generation of low molecular weight degradation products.

## EXAMPLE 9

### *Chromosomal localization of the hpa gene*

Chromosomal mapping of the *hpa* gene was performed utilizing a panel of monochromosomal human/CHO and human/mouse somatic cell hybrids, obtained from the UK HGMP Resource Center (Cambridge, England).

40 ng of each of the somatic cell hybrid DNA samples were subjected to PCR amplification using the *hpa* primers: hpu565 5'-AGCTCTGTAGATGTGC TATACAC-3', SEQ ID NO:22, corresponding to nucleotides 564-586 of SEQ ID NO:9 and an antisense primer hpl171 5'-GCATCTTAGCCGTCTTTCTTCG-3', SEQ ID NO:23, corresponding to nucleotides 897-876 of SEQ ID NO:9.

The PCR program was as follows: a hot start of 94 °C – 3 minutes, followed by 7 cycles of 94 °C – 45 seconds, 66 °C – 1 minute, 68 °C – 5 minutes, followed by 30 cycles of 94 °C – 45 seconds, 62 °C – 1 minute, 68 °C – 5 minutes, and a 10 minutes final extension at 72 °C.

The reactions were performed with Expand long PCR (Boehringer Mannheim). The resulting amplification products were analyzed using agarose gel electrophoresis. As demonstrated in Figure 14, a single band of approximately 2.8 Kb was obtained from chromosome 4, as well as from the control human genomic DNA. A 2.8 kb amplification product is expected based on amplification of the genomic *hpa* clone (data not shown). No amplification products were obtained neither in the control DNA samples of hamster and mouse nor in somatic hybrids of other human chromosome.

## EXAMPLE 10

### *Human genomic clone encoding heparanase*

Five plaques were isolated following screening of a human genomic library and were designated L3-1, L5-1, L8-1, L10-1 and L6-1. The phage DNAs were analyzed by Southern hybridization and by PCR with *hpa* specific and vector specific primers. Southern analysis was performed with three fragments of *hpa* cDNA: a *PvuII-BamHI* fragment (nucleotides 32-450, SEQ ID NO:9), a *BamHI-NdeI* fragment (nucleotides 451-1102, SEQ ID NO:9) and an *NdeI-XhoI* fragment (nucleotides 1103-1721, SEQ ID NO:9).

Following Southern analysis, phages L3, L6, L8 were selected for further analysis. A scheme of the genomic region and the relative position of the three phage clones is depicted in Figure 15. A 2 kb DNA fragment containing the gap between phages L6 and L3 was PCR amplified from human genomic DNA with two gene specific primers GHpuL3 and GHplL6. The PCR product was cloned into the plasmid vector pGEM-T-easy (Promega).

Large scale DNA sequencing of the three Lambda clones and the amplified fragment was performed with Lambda purified DNA by primer walking. A nucleotide sequence of 44,898 bp was analyzed (Figure 16, SEQ ID NO:42). Comparison of the genomic sequence with that of *hpa* cDNA revealed 12 exons separated by 11 introns (Figures 15 and 16). The genomic organization of the *hpa* gene is depicted in Figure 15 (top). The sequence includes the coding region from the first ATG to the stop codon which spans 39,113 nucleotides, 2742 nucleotides upstream of the first ATG and 3043 nucleotides downstream of the stop codon. Splice site consensus sequences were identified at exon/intron junctions.

### EXAMPLE 11

#### *Alternative splicing*

Several minor RT-PCR products were obtained from various cell types, following amplification with *hpa* specific primers. Each one found to contain a deletion of one or two exons. Some of these PCR products contain ORFs, which encode potential shorter proteins.

Table 1 below summarizes the alternative spliced products isolated from various cell lines.

Fragments of similar sizes were obtained following amplification with two cell lines, placenta and platelets.

Cell type	Nucleotides deleted	Exons deleted	ORF
Platelets	1047-1267	8, 9	+
Platelets	1154-1267	9	-
Platelets	289-435, 562-735	2, 4	-
Sk-hep1, platelets, Zr75	562-735	4	+
Sk-hep1 (hepatoma)	561-904	4, 5	-
Zr75 (breast carcinoma)	96-203	1 (partial)	+

### EXAMPLE 12

#### *Mouse and rat hpa*

EST databases were screened for sequences homologous to the *hpa* gene. Three mouse EST's were identified (accession No. Aa177901, from mouse spleen, Aa067997 from mouse skin, Aa47943 from mouse embryo), assembled into a 824 bp cDNA fragment which contains a partial open reading frame (lacking a 5' end) of 629 bp and a 3' untranslated region of



195 bp (SEQ ID NO:12). As shown in Figure 13, the coding region is 80 % similar to the 3' end of the *hpa* cDNA sequence. These EST's are probably cDNA fragments of the mouse *hpa* homolog that encodes for the mouse heparanase.

5 Searching for consensus protein domains revealed an amino terminal homology between the heparanase and several precursor proteins such as Procollagen Alpha 1 precursor, Tyrosine-protein kinase-RYK, Fibulin-1, Insulin-like growth factor binding protein and several others. The amino terminus is highly hydrophobic and contains a potential trans-membrane  
10 domain. The homology to known signal peptide sequences suggests that it could function as a signal peptide for protein localization.

The amino acid sequence of human heparanase was used to search for homologous sequences in the DNA and protein databases. Several human EST's were identified, as well as mouse sequences highly  
15 homologous to human heparanase. The following mouse EST's were identified AA177901, AA674378, AA67997, AA047943, AA690179, AI122034, all sharing an identical sequence and correspond to amino acids 336-543 of the human heparanase sequence. The entire mouse heparanase cDNA was cloned, based on the nucleotide sequence of the mouse EST's.  
20 PCR primers were designed and a Marathon RACE was performed using a Marathon cDNA library from 15 days mouse embryo (Clontech) and from BL6 mouse melanoma cell line. The mouse *hpa* homologous cDNA was isolated following several amplification steps. A 1.1 kb fragment was amplified from mouse embryo Marathon cDNA library. The first cycle of  
25 amplification was performed with primers mhpl773 and Ap1 and the second cycle with primers mhpl736 and AP2. A 1.1 kb fragment was then amplified from BL6 Marathon cDNA library. The first cycle of amplification was performed with the primers mhpl152 and Ap1, and the second with mhpl83 and AP2. The combined sequence was homologous to  
30 nucleotides 157 - 1702 of the human *hpa* cDNA, which encode amino acids 33-543. The 5' end of the mouse *hpa* gene was isolated from a mouse genomic DNA library using the Genome Walker kit (Clontech). An 0.9 kb fragment was amplified from a *DraI* digested Genome walker DNA library. The first cycle of amplification was performed with primers mhpl114 and  
35 Ap1 and the second with primers mhpl103 and AP2. The assembled sequence (SEQ ID NOs:43, 45) is 2396 nucleotides long. It contains an open reading frame of 1605 nucleotides, which encode a polypeptide of 535 amino acids (SEQ ID NOs:44, 45), 196 nucleotides of 3' untranslated

region (UTR), and anupstream sequence which includes the promoter region and the 5'-UTR of the mouse *hpa* cDNA.. According to two promoter predicting programs TSSW and TSSG, the transcription start site is localized to nucleotide 431 of SEQ ID NOs:43, 45, 163 nucleotides upstream of the first ATG codon. The 431 upstream genomic sequence contains the promoter region. A TATA box is predicted at position 394 of SEQ ID NOs:43, 45. The mouse and the human *hpa* genes share an average homology of 78 % between the nucleotide sequences and 81 % similarity between the deduced amino acid sequences.

Search for *hpa* homologous sequences, using the Blast 2.0 server revealed two EST's from rat: AI060284 (385 nucleotides, SEQ ID NO:46) which is homologous to the amino terminus (68 % similarity to amino acids 12-136) of human heparanase and AI237828 (541 nucleotides, SEQ ID NO:47) which is homologous to the carboxyl terminus (81 % similarity to amino acids 500-543) of human heparanase, and contains a 3'-UTR. A comparison between the human heparanase and the mouse and rat homologous sequences is demonstrated in Figure 17.

### EXAMPLE 13

#### *Prediction of heparanase active site*

Homology search of heparanase amino acid sequence against the DNA and the protein databases revealed no significant homologies. The protein secondary structure as predicted by the PHD program consists of alternating alpha helices and beta sheets. The fold recognition server of UCLA predicted alpha/beta barrel structure, with under-threshold confidence.

Five of 15 proteins, which were predicted to have most similar folds, were glycosyl hydrolases from various organisms: 1xyza – xylanase from *Clostridium Thermocellum*, 1pbga – 6-phospho-beta- $\delta$ -galactosidase from *Lactococcus Lactis*, 1amy – alpha-amylase from Barley, 1ecea – endocellulase from *Acidothermus Cellulolyticus* and 1qbc – hexosaminidase alpha chain, glycosyl hydrolase.

Protein homology search using the bioaccelerator pulled out several proteins, including glycosyl hydrolyses such as beta-fructofuranosidase from *Vicia faba* (broad bean) and from potato, lactase phlorizin hydrolase from human, xylanases from *Clostridium thermocellum* and from *Streptomyces halstedii* and cellulase from *Clostridium thermocellum*. Blocks 9.3 database pulled out the active site of glycosyl hydrolases family

five, which includes cellulases from various bacteria and fungi. Similar active site motif is shared by several lysosomal acid hydrolases (63) and other glycosyl hydrolases. The common mechanism shared by these enzymes involves two glutamic acid residues, a proton donor and a nucleophile.

Despite the lack of an overall homology between the heparanase and other glycosyl hydrolases, the amino acid couple Asp-Glu (NE), which is characteristic of the proton donor of glycosyl hydrolyses of the GH-A clan, was found at positions 224-225 of the human heparanase protein sequence. As in other clan members, this NE couple is located at the end of a  $\beta$  sheet.

Considering the relative location of the proton donor and the predicted secondary structure, the glutamic acid that functions as nucleophile is most likely located at position 343, or at position 396. Identification of the active site and the amino acids directly involved in hydrolysis opens the way for expression of the defined catalytic domain. In addition, it will provide the tools for rational design of enzyme activity either by modification of the microenvironment or catalytic site itself.

#### EXAMPLE 14

##### *Expression of hpa antisense in mammalian cell lines*

A mammalian expression vector Hpa2Kepcdna3 was constructed in order to express *hpa* antisense in mammalian cells. *hpa* cDNA (1.7 kb *EcoRI* fragment) was cloned into the plasmid pCDNA3 in 3'>5' (antisense) orientation. The construct was used to transfect MBT2-T50 and T24P cell lines. 2 x 10<sup>5</sup> cells in 35 mm plates were transfected using the Fugene protocol (Boehringer Mannheim). 48 hours after transfection cells were trypsinized and seeded in six well plates. 24 hours later G418 was added to initiate selection. The number of colonies per 35 mm plate following 3 weeks:

	Antisense	No insert
T24P	15	60
MBT-T50	1	6

The lower number of colonies obtained after transfection with *hpa* antisense, as compared with the control plasmid suggests that the introduction of *hpa* antisense interfere with cell growth. This experiment demonstrates the use of complementary antisense *hpa* DNA sequence to

control heparanase expression in cells. This approach may be used to inhibit expression of heparanase *in vivo*, in, for example, cancer cells and in other pathological processes in which heparanase is involved.

5

### EXAMPLE 15

#### *Zoo blot*

*Hpa* cDNA was used as a probe to detect homologous sequences in human DNA and in DNA of various animals. The autoradiogram of the Southern analysis is presented in Figure 18. Several bands were detected in human DNA, which correlated with the accepted pattern according to the genomic *hpa* sequence. Several intense bands were detected in all mammals, while faint bands were detected in chicken. This correlates with the phylogenetic relation between human and the tested animals. The intense bands indicate that *hpa* is conserved among mammals as well as in more genetically distant organisms. The multiple bands patterns suggest that in all animals, like in human, the *hpa* locus occupy large genomic region. Alternatively, the various bands could represent homologous sequences and suggest the existence of a gene family, which can be isolated based on their homology to the human *hpa* reported herein. This conservation was actually found, between the isolated human *hpa* cDNA and the mouse homologue.

25

### EXAMPLE 16

#### *Characterization of the hpa promoter*

The DNA sequence upstream of the *hpa* first ATG was subjected to computational analysis in order to localize the predicted transcription start site and to identify potential transcription factors binding sites. Recognition of human PolII promoter region and start of transcription were predicted using the TSSW and TSSG programs. Both programs identified a promoter region upstream of the coding region. TSSW pointed at nucleotide 2644 and TSSG at 2635 of SEQ ID NO:42. These two predicted transcription start sites are located 4 and 13 nucleotides upstream of the longest *hpa* cDNA isolated by RACE.

A *hpa* promoter-GFP reporter vector was constructed in order to investigate the regulation of *hpa* transcription. Two constructs were made, containing 1.8 kb and 1.1 kb of the *hpa* promoter region. The reporter vector was transfected into T50-mouse bladder carcinoma cells. Cells transfected with both constructs exhibited green fluorescence, which

indicated the promoter activity of the genomic sequence upstream of the *hpa*-coding region. This reporter vector, enables the monitoring of *hpa* promoter activity, at various conditions and in different cell types and to characterize the factors involved regulation of *hpa* expression.

5

Although the invention has been described in conjunction with specific embodiments thereof, it is evident that many alternatives, modifications and variations will be apparent to those skilled in the art. Accordingly, it is intended to embrace all such alternatives, modifications  
10 and variations that fall within the spirit and broad scope of the appended claims.

LIST OF REFERENCES

1. Wight, T.N., Kinsella, M.G., and Qwarnstromn, E.E. (1992). The role of proteoglycans in cell adhesion, migration and proliferation. *Curr. Opin. Cell Biol.*, 4, 793-801.
2. Jackson, R.L., Busch, S.J., and Cardin, A.L. (1991). Glycosaminoglycans: Molecular properties, protein interactions and role in physiological processes. *Physiol. Rev.*, 71, 481-539.
3. Wight, T.N. (1989). Cell biology of arterial proteoglycans. *Arteriosclerosis*, 9, 1-20.
4. Kjellen, L., and Lindahl, U. (1991). Proteoglycans: structures and interactions. *Annu. Rev. Biochem.*, 60, 443-475.
5. Ruoslahti, E., and Yamaguchi, Y. (1991). Proteoglycans as modulators of growth factor activities. *Cell*, 64, 867-869.
6. Vlodavsky, I., Eldor, A., Haimovitz-Friedman, A., Matzner, Y., Ishai-Michaeli, R., Levi, E., Bashkin, P., Lider, O., Naparstek, Y., Cohen, I.R., and Fuks, Z. (1992). Expression of heparanase by platelets and circulating cells of the immune system: Possible involvement in diapedesis and extravasation. *Invasion & Metastasis*, 12, 112-127.
7. Vlodavsky, I., Mohsen, M., Lider, O., Ishai-Michaeli, R., Ekre, H.-P., Svahn, C.M., Vigoda, M., and Peretz, T. (1995). Inhibition of tumor metastasis by heparanase inhibiting species of heparin. *Invasion & Metastasis*, 14, 290-302.
8. Nakajima, M., Irimura, T., and Nicolson, G.L. (1988). Heparanase and tumor metastasis. *J. Cell. Biochem.*, 36, 157-167.
9. Nicolson, G.L. (1988). Organ specificity of tumor metastasis: Role of preferential adhesion, invasion and growth of malignant cells at specific secondary sites. *Cancer Met. Rev.*, 7, 143-188.

10. Liotta, L.A., Rao, C.N., and Barsky, S.H. (1983). Tumor invasion and the extracellular matrix. *Lab. Invest.*, 49, 639-649.
11. Vlodavsky, I., Fuks, Z., Bar-Ner, M., Ariav, Y., and Schirmacher, V. (1983). Lymphoma cell mediated degradation of sulfated proteoglycans in the subendothelial extracellular matrix: Relationship to tumor cell metastasis. *Cancer Res.*, 43, 2704-2711.
12. Vlodavsky, I., Ishai-Michaeli, R., Bar-Ner, M., Fridman, R., Horowitz, A.T., Fuks, Z. and Biran, S. (1988). Involvement of heparanase in tumor metastasis and angiogenesis. *Is. J. Med.*, 24, 464-470.
13. Vlodavsky, I., Liu, G.M., and Gospodarowicz, D. (1980). Morphological appearance, growth behavior and migratory activity of human tumor cells maintained on extracellular matrix vs. plastic. *Cell*, 19, 607-616.
14. Gospodarowicz, D., Delgado, D., and Vlodavsky, I. (1980). Permissive effect of the extracellular matrix on cell proliferation in-vitro. *Proc. Natl. Acad. Sci. USA.*, 77, 4094-4098.
15. Bashkin, P., Doctrow, S., Klagsbrun, M., Svahn, C.M., Folkman, J., and Vlodavsky, I. (1989). Basic fibroblast growth factor binds to subendothelial extracellular matrix and is released by heparitinase and heparin-like molecules. *Biochemistry*, 28, 1737-1743.
16. Parish, C.R., Coombe, D.R., Jakobsen, K.B., and Underwood, P.A. (1987). Evidence that sulphated polysaccharides inhibit tumor metastasis by blocking tumor cell-derived heparanase. *Int. J. Cancer*, 40, 511-517.
- 16a. Vlodavsky, I., Hua-Quan Miao., Benezra, M., Lider, O., Bar-Shavit, R., Schmidt, A., and Peretz, T. (1997). Involvement of the extracellular matrix, heparan sulfate proteoglycans and heparan sulfate degrading enzymes in angiogenesis and metastasis. In: *Tumor Angiogenesis*. Eds. C.E. Lewis, R. Bicknell & N. Ferrara. Oxford University Press, Oxford UK, pp. 125-140.

17. Burgess, W.H., and Maciag, T. (1989). The heparin-binding (fibroblast) growth factor family of proteins. *Annu. Rev. Biochem.*, 58, 575-606.
18. Folkman, J., and Klagsbrun, M. (1987). Angiogenic factors. *Science*, 235, 442-447.
19. Vlodavsky, I., Folkman, J., Sullivan, R., Fridman, R., Ishai-Michaeli, R., Sasse, J., and Klagsbrun, M. (1987). Endothelial cell-derived basic fibroblast growth factor: Synthesis and deposition into subendothelial extracellular matrix. *Proc. Natl. Acad. Sci. USA*, 84, 2292-2296.
20. Folkman, J., Klagsbrun, M., Sasse, J., Wadzinski, M., Ingber, D., and Vlodavsky, I. (1980). A heparin-binding angiogenic protein - basic fibroblast growth factor - is stored within basement membrane. *Am. J. Pathol.*, 130, 393-400.
21. Cardon-Cardo, C., Vlodavsky, I., Haimovitz-Friedman, A., Hicklin, D., and Fuks, Z. (1990). Expression of basic fibroblast growth factor in normal human tissues. *Lab. Invest.*, 63, 832-840.
22. Ishai-Michaeli, R., Svahn, C.-M., Chajek-Shaul, T., Korner, G., Ekre, H.-P., and Vlodavsky, I. (1992). Importance of size and sulfation of heparin in release of basic fibroblast factor from the vascular endothelium and extracellular matrix. *Biochemistry*, 31, 2080-2088.
23. Ishai-Michaeli, R., Eldor, A., and Vlodavsky, I. (1990). Heparanase activity expressed by platelets, neutrophils and lymphoma cells releases active fibroblast growth factor from extracellular matrix. *Cell Reg.*, 1, 833-842.
24. Vlodavsky, I., Bar-Shavit, R., Ishai-Michaeli, R., Bashkin, P., and Fuks, Z. (1991). Extracellular sequestration and release of fibroblast growth factor: a regulatory mechanism? *Trends Biochem. Sci.*, 16, 268-271.
25. Vlodavsky, I., Bar-Shavit, R., Korner, G., and Fuks, Z. (1993). Extracellular matrix-bound growth factors, enzymes and plasma proteins. In *Basement membranes: Cellular and molecular aspects* (eds.



D.H. Rohrbach and R. Timpl), pp327-343. Academic press Inc., Orlando, Fl.

26. Yayon, A., Klagsbrun, M., Esko, J.D., Leder, P., and Ornitz, D.M. (1991). Cell surface, heparin-like molecules are required for binding of basic fibroblast growth factor to its high affinity receptor. *Cell*, 64, 841-848.

27. Spivak-Kroizman, T., Lemmon, M.A., Dikic, I., Ladbury, J.E., Pinchasi, D., Huang, J., Jaye, M., Crumley, G., Schlessinger, J., and Lax, I. (1994). Heparin-induced oligomerization of FGF molecules is responsible for FGF receptor dimerization, activation, and cell proliferation. *Cell*, 79, 1015-1024.

28. Ornitz, D.M., Herr, A.B., Nilsson, M., West, a., J., Svahn, C.-M., and Waksman, G. (1995). FGF binding and FGF receptor activation by synthetic heparan-derived di- and trisaccharides. *Science*, 268, 432-436.

29. Gitay-Goren, H., Soker, S., Vlodavsky, I., and Neufeld, G. (1992). Cell surface associated heparin-like molecules are required for the binding of vascular endothelial growth factor (VEGF) to its cell surface receptors. *J. Biol. Chem.*, 267, 6093-6098.

30. Lider, O., Baharav, E., Mekori, Y., Miller, T., Naparstek, Y., Vlodavsky, I., and Cohen, I.R. (1989). Suppression of experimental autoimmune diseases and prolongation of allograft survival by treatment of animals with heparinoid inhibitors of T lymphocyte heparanase. *J. Clin. Invest.*, 83, 752-756.

31. Lider, O., Cahalon, L., Gilat, D., HersHKovitz, R., Siegel, D., Margalit, R., Shoseyov, O., and Cohn, I.R. (1995). A disaccharide that inhibits tumor necrosis factor  $\alpha$  is formed from the extracellular matrix by the enzyme heparanase. *Proc. Natl. Acad. Sci. USA.*, 92, 5037-5041.

31a. Rapraeger, A., Krufka, A., and Olwin, B.R. (1991). Requirement of heparan sulfate for bFGF-mediated fibroblast growth and myoblast differentiation. *Science*, 252, 1705-1708.

32. Eisenberg, S., Schayek, E., Olivecrona, T., and Vlodavsky, I. (1992). Lipoprotein lipase enhances binding of lipoproteins to heparan sulfate on cell surfaces and extracellular matrix. *J. Clin. Invest.*, 90, 2013-2021.
33. Shieh, M-T., Wundunn, D., Montgomery, R.I., Esko, J.D., and Spear, P.G. J. (1992). Cell surface receptors for herpes simplex virus are heparan sulfate proteoglycans. *J Cell Biol.*, 116, 1273-1281.
- 33a. Chen, Y., Maguire, T., Hileman, R.E., Fromm, J.R., Esko, J.D., Linhardt, R.J., and Marks, R.M. (1997). Dengue virus infectivity depends on envelope protein binding to target cell heparan sulfate. *Nature Medicine* 3, 866-871.
- 33b. Putnak, J.R., Kanasa-Thanan, N., and Innis, B.L. (1997). A putative cellular receptor for dengue viruses. *Nature Medicine* 3, 828-829.
34. Narindrasorasak, S., Lowery, D., Gonzalez-DeWhitt, P., Poorman, R.A., Greenberg, B., Kisilevsky, R. (1991). High affinity interactions between the Alzheimer's beta-amyloid precursor protein and the basement membrane form of the heparan sulfate proteoglycan. *J. Biol. Chem.*, 266, 12878-83.
35. Ross, R. (1993). The pathogenesis of atherosclerosis: a perspective for the 1990s. *Nature (Lond.)*, 362:801-809.
36. Zhong-Sheng, J., Walter, J., Brecht, R., Miranda, D., Mahmood Hussain, M., Innerarity, T.L. and Mahley, W.R. (1993). Role of heparan sulfate proteoglycans in the binding and uptake of apolipoprotein E-enriched remnant lipoproteins by cultured cells. *J. Biol. Chem.*, 268, 10160-10167.
37. Ernst, S., Langer, R., Cooney, Ch.L., and Sasisekharan, R. (1995). Enzymatic degradation of glycosaminoglycans. *Critical Reviews in Biochemistry and Molecular Biology*, 30(5), 387-444.

38. Gospodarowicz, D., Mescher, AL., Birdwell, CR. (1977). Stimulation of corneal endothelial cell proliferation in vitro by fibroblast and epidermal growth factors. *Exp Eye Res* 25, 75-89.
39. Haimovitz-Friedman, A., Falcone, D.J., Eldor, A., Schirmacher, V., Vlodavsky, I., and Fuks, Z. (1991) Activation of platelet heparitinase by tumor cell-derived factors. *Blood*, 78, 789-796.
- 39a. Savitsky, K., Platzer, M., Uziel, T., Gilad, S., Sartiel, A., Rosental, A., Elroy-Stein, O., Siloh, Y. and Rotman, G. (1997). Ataxia-telangiectasia: structural diversity of untranslated sequences suggests complex post-translational regulation of ATM gene expression. *Nucleic Acids Res.* 25(9), 1678-1684.
40. Bar-Ner, M., Eldor, A., Wasserman, L., Matzner, Y., and Vlodavsky, I. (1987). Inhibition of heparanase mediated degradation of extracellular matrix heparan sulfate by modified and non-anticoagulant heparin species. *Blood*, 70, 551-557.
41. Goshen, R., Hochberg, A., Korner, G., Levi, E., Ishai-Michaeli, R., Elkin, M., de Grot, N., and Vlodavsky, I. (1996). Purification and characterization of placental heparanase and its expression by cultured cytotrophoblasts. *Mol. Human Reprod.*, 2, 679-684.
42. Korb M., Ke Y. and Johnson L.F. (1993) Stimulation of gene expression by introns: conversion of an inhibitory intron to a stimulatory intron by alteration of the splice donor sequence. *Nucleic Acids Res.*, 21(25):5901-8.
43. Zheng B., Qiu X.Y., Tan M., Xing Y.N., Lo D., Xue J.L. and Qiu X.F. (1997) Increment of hFIX expression with endogenous intron 1 in vitro. *Cell Res.*, 7(1):21-29.
44. Kurachi S., Hitomi Y., Furukawa M. and Kurachi K. (1995) Role of intron I in expression of the human factor IX gene. *J. Biol. Chem.* 270(10):5276-5281.

45. Shekhar P.V. and Miller F.R. (1994-5) Correlation of differences in modulation of ras expression with metastatic competence of mouse mammary tumor subpopulations. *Invasion Metastasis*, 14(1-6):27-37.
46. Zhou G., Garofalo S., Mukhopadhyay K., Lefebvre V., Smith C.N., Eberspaecher H. and de Crombrughe B. (1995) A 182 bp fragment of the mouse pro alpha 1(II) collagen gene is sufficient to direct chondrocyte expression in transgenic mice. *J. Cell Sci.*, 108 ( Pt 12):3677-3684.
47. Hormuzdi S.G., Penttinen R., Jaenisch R. and Bornstein P. (1998) A gene-targeting approach identifies a function for the first intron in expression of the alpha1(I) collagen gene. *Mol. Cell*, 18(6):3368-3375.
48. Kang Y.K., Lee C.S., Chung A.S. and Lee K.K. (1998) Prolactin-inducible enhancer activity of the first intron of the bovine beta-casein gene. *Mol. Cells*, 30;8(3):259-265.
49. Chow Y.H., O'Brodovich H., Plumb J., Wen Y., Sohn K.J., Lu Z., Zhang F., Lukacs G.L., Tanswell A.K., Hui C.C., Buchwald M. and Hu J. (1997) Development of an epithelium-specific expression cassette with human DNA regulatory elements for transgene expression in lung airways. *Proc. Natl. Acad. Sci. USA*, 23;94(26):14695-14700.
50. Gottschalk U. and Chan S. (1998) Somatic gene therapy. Present situation and future perspective. *Arzneimittelforschung*, 48(11):1111-1120.
51. Ye S., Cole-Strauss A.C., Frank B. and Kmiec E.B. (1998) Targeted gene correction: a new strategy for molecular medicine. *Mol. Med. Today*, 4(10):431-437.
52. Lai L., and Lien Y. (1999) Homologous recombination based gene therapy. *Exp. Nephrol.*, 7(1):11-14.

53. Yazaki N., Fujita H., Ohta M., Kawasaki T. and Itoh N. (1993) The structure and expression of the FGF receptor-1 mRNA isoforms in rat tissues. *Biochim. Biophys. Acta.*, 20;1172(1-2):37-42.
54. Le Fur N., Kelsall S.R., Silvers W.K. and Mintz B. (1997) Selective increase in specific alternative splice variants of tyrosinase in murine melanomas: a projected basis for immunotherapy. *Proc. Natl. Acad. Sci. USA*, 13;94(10):5332-5337.
55. Miyake H., Okamoto I., Hara I., Gohji K., Yamanaka K., Arakawa S., Kamidono S. and Saya H. (1998) Highly specific and sensitive detection of malignancy in urine samples from patients with urothelial cancer by CD44v8-10/CD44v10 competitive RT-PCR. *Int. J. Cancer*, 18;79(6):560-564.
56. Guriec N., Marcellin L., Gairard B., Calderoli H., Wilk A., Renaud R., Bergerat J.P. and Oberling F. (1996) CD44 exon 6 expression as a possible early prognostic factor in primary node negative breast carcinoma. *Clin. Exp. Metastasis*, 14(5):434-439.
57. Gewirtz A.M., Sokol D.L. and Ratajczak M.Z. (1998) Nucleic acid therapeutics: state of the art and future prospects. *Blood*, 1;92(3):712-736.
58. Hida K., Shindoh M., Yasuda M., Hanzawa M., Funaoka K., Kohgo T., Amemiya A., Totsuka Y., Yoshida K. and Fujinaga K (1997) Antisense E1AF transfection restrains oral cancer invasion by reducing matrix metalloproteinase activities. *Am. J. Pathol.* 150(6):2125-2132.
59. Shastry B.S. (1998) Gene disruption in mice: models of development and disease. *Mol. Cell. Biochem.* 1998 Apr;181(1-2):163-179.
60. Carpentier A.F., Rosenfeld M.R., Delattre J.Y., Whalen R.G., Posner J.B. and Dalmau J. (1998) DNA vaccination with HuD inhibits growth of a neuroblastoma in mice. *Clin. Cancer Res.*, 4(11):2819-2824.
61. Lai W.C. and Bennett M. (1998) DNA vaccines. *Crit. Rev. Immunol.*, 18(5):449-484.

62. Welch P.J., Barber J.R., and Wong-Staal F. (1998) Expression of ribozymes in gene transfer systems to modulate target RNA levels. *Curr. Opin. Biotechnol.*, 9(5):486-496.
63. Durand P., Lehn P., Callebaunt I., Fabrega S., Henrissat B. and Mornon J.P. (1997) Active-site motifs of lysosomal acid hydrolyses: invariant features of clan GH-A glycosyl hydrolases deduced from hydrophobic cluster analysis. *Glycobiology*, 7(2):277-284.
64. Thuong and Helene (1993) Sequence specific recognition and modification of double helical DNA by oligonucleotides *Angev. Chem. Int. Ed. Engl.* 32:666
65. Dash P., Lotan I., Knapp M., Kandel E.R. and Goelet P. (1987) Selective elimination of mRNAs in vivo: complementary oligodeoxynucleotides promote RNA degradation by an RNase H-like activity. *Proc. Natl. Acad. Sci. USA*, 84:7896.
66. Chiang M.Y., Chan H., Zounes M.A., Freier S.M., Lima W.F. and Bennett C.F. (1991) Antisense oligonucleotides inhibit intercellular adhesion molecule 1 expression by two distinct mechanisms. *J. Biol. Chem.* 266:18162-71.
- 5 67. Paterson Paterson B.M., Roberts B.E and Kuff EL . (1977) Structural gene identification and mapping by DNA-mRNA hybrid-arrested cell-free translation. *Proc. Natl. Acad. Sci. USA*, 74:4370.
68. Cohen (1992) Oligonucleotide therapeutics. *Trends in Biotechnology*, 10:87.
69. Szczylik et al (1991) Selective inhibition of leukemia cell proliferation by BCR-ABL antisense oligodeoxynucleotides. *Science* 253:562.
70. Calabretta et al. (1991) Normal and leukemic hematopoietic cell manifest differential sensitivity to inhibitory effects of c-myc antisense

oligodeoxynucleotides: an in vitro study relevant to bone marrow purging. Proc. Natl. Acad. Sci. USA 88:2351.

71. Heikhila et al. (1987) A c-myc antisense oligodeoxynucleotide inhibits entry into S phase but not progress from G(0) to G(1). Nature, 328:445.

72. Reed et al. (1990) Antisense mediated inhibition of BCL2 protooncogene expression and leukemic cell growth and survival: comparison of phosphodiester and phosphorothioate oligodeoxynucleotides. Cancer Res. 50:6565.

73. Burch and Mahan (1991) Oligodeoxynucleotides antisense to the interleukin I receptor mRNA block the effects of interleukin I in cultured murine and human fibroblasts and in mice. J. Clin. Invest. 88:1190.

74. Agrawal (1992) Antisense oligonucleotides as antiviral agents. TIBTECH 10:152.

75. Uhlmann et al. (1990) Chem. Rev. 90:544.

76. Cook (1991) Medicinal chemistry of antisense oligonucleotides - future opportunities. Anti-Cancer Drug Design 6:585.

77. Biotechnology research news (1993) Can DNA mimics improve on the real thing? Science 262:1647.

## WHAT IS CLAIMED IS:

1. An isolated nucleic acid comprising a genomic, complementary or composite polynucleotide sequence encoding a polypeptide having heparanase catalytic activity.
2. The isolated nucleic acid of claim 1, wherein said polynucleotide or a portion thereof is hybridizable with SEQ ID NOs: 9, 13, 42, 43 or a portion thereof at 68 °C in 6 x SSC, 1 % SDS, 5 x Denharts, 10 % dextran sulfate, 100 µg/ml salmon sperm DNA, and <sup>32</sup>p labeled probe and wash at 68 °C with 3 x SSC and 0.1 % SDS.
3. The isolated nucleic acid of claim 1, wherein said polynucleotide or a portion thereof is at least 60 % identical with SEQ ID NOs: 9, 13, 42, 43 or portions thereof as determined using the Bestfit procedure of the DNA sequence analysis software package developed by the Genetic Computer Group (GCG) at the university of Wisconsin (gap creation penalty - 12, gap extension penalty - 4).
4. The isolated nucleic acid of claim 1, wherein said polypeptide is as set forth in SEQ ID NOs:10, 14, 44 or portions thereof.
5. The isolated nucleic acid of claim 1, wherein said polypeptide is at least 60 % homologous to SEQ ID NOs:10, 14, 44 or portions thereof as determined with the Smith-Waterman algorithm, using the Bioaccelerator platform developed by Compugene (gapop: 10.0, gapext: 0.5, matrix: blosum62).
6. A nucleic acid construct comprising the isolated nucleic acid of claim 1.
7. A host cell comprising the nucleic acid construct of claim 6.
8. An antisense oligonucleotide comprising a polynucleotide or a polynucleotide analog of at least 10 bases being hybridizable *in vivo*, under physiological conditions, with a portion of a polynucleotide strand encoding a polypeptide having heparanase catalytic activity.



9. The antisense oligonucleotide of claim 8, wherein said polynucleotide strand encoding said polypeptide having heparanase catalytic activity is as set forth in SEQ ID NOs: 9, 13, 42, or 43.

10. The antisense oligonucleotide of claim 8, wherein said polypeptide having heparanase catalytic activity is as set forth in SEQ ID NOs: 10, 14 and 44.

11. A method of *in vivo* downregulating heparanase activity comprising the step of *in vivo* administering the antisense oligonucleotide of claim 8.

12. A pharmaceutical composition comprising the antisense oligonucleotide of claim 8 and a pharmaceutically acceptable carrier.

13. A ribozyme comprising the antisense oligonucleotide of claim 8 and a ribozyme sequence.

14. An antisense nucleic acid construct comprising a promoter sequence and a polynucleotide sequence directing the synthesis of an antisense RNA sequence of at least 10 bases being hybridizable *in vivo*, under physiological conditions, with a portion of a polynucleotide strand encoding a polypeptide having heparanase catalytic activity.

15. The antisense nucleic acid construct of claim 14, wherein said polynucleotide strand encoding said polypeptide having heparanase catalytic activity is as set forth in SEQ ID NOs: 9, 13, 42 or 43.

16. The antisense nucleic acid construct of claim 14, wherein said polypeptide having heparanase catalytic activity is as set forth in SEQ ID NOs: 10, 14 or 44.

17. A method of *in vivo* downregulating heparanase activity comprising the step of *in vivo* administering the antisense nucleic acid construct of claim 14.

18. A pharmaceutical composition comprising the antisense nucleic acid construct of claim 14 and a pharmaceutically acceptable carrier.

19. A nucleic acid construct comprising a polynucleotide sequence functioning as a promoter, said polynucleotide sequence is derived from SEQ ID NO:42 and includes at least nucleotides 2535-2635 thereof or from SEQ ID NO:43 and includes at least nucleotides 320-420.

20. A method of expressing a polynucleotide sequence comprising the step of ligating the polynucleotide sequence into the nucleic acid construct of claim 19, downstream of said polynucleotide sequence derived from SEQ ID NOs:42 or 43.

21. A recombinant protein comprising a polypeptide having heparanase catalytic activity.

22. The recombinant protein of claim 21, wherein said polypeptide includes at least a portion of SEQ ID NOs:10, 14 or 44.

23. The recombinant protein of claim 21, wherein the protein is encoded by a polynucleotide hybridizable with SEQ ID NOs: 9, 13, 42, 43 or a portion thereof at 68 °C in 6 x SSC, 1 % SDS, 5 x Denharts, 10 % dextran sulfate, 100 µg/ml salmon sperm DNA, and <sup>32</sup>p labeled probe and wash at 68 °C with 3 x SSC and 0.1 % SDS.

24. The recombinant protein of claim 21, wherein the protein is encoded by a polynucleotide at least 60 % identical with SEQ ID NOs: 9, 13, 42, 43 or portions thereof as determined using the Bestfit procedure of the DNA sequence analysis software package developed by the Genetic Computer Group (GCG) at the university of Wisconsin (gap creation penalty - 12, gap extension penalty - 4).

25. A pharmaceutical composition comprising, as an active ingredient, the recombinant protein of claim 21.

26. A method of identifying a chromosome region harboring a heparanase gene in a chromosome spread comprising the steps of:

- (a) hybridizing the chromosome spread with a tagged polynucleotide probe encoding heparanase;
- (b) washing the chromosome spread, thereby removing excess of non-hybridized probe; and
- (c) searching for signals associated with said hybridized tagged polynucleotide probe, wherein detected signals being indicative of a chromosome region harboring a heparanase gene.

27. A method of *in vivo* eliciting anti-heparanase antibodies comprising the steps of administering a nucleic acid construct including a polynucleotide segment corresponding to at least a portion of SEQ ID NOs:9, 13 or 43 and a promoter for directing the expression of said polynucleotide segment *in vivo*.

28. A DNA vaccine for *in vivo* eliciting anti-heparanase antibodies comprising a nucleic acid construct including a polynucleotide segment corresponding to at least a portion of SEQ ID NOs:9, 13 or 43 and a promoter for directing the expression of said polynucleotide segment *in vivo*.


1/34

Fig 1

1 CTAGAGCTTTCGACTCTCCGCTGCGCGGCAGCTGGCGGGGGAGCAGCCAGGTGAGCCCA  
61 AGATGCTGCTGCGCTCGAAGCTTTCGCTGCGCGCGCGCTGATGCTGCTGCTCTGCGGC  
M L L R S K P A L P P P L M L L L L G P  
121 CGCTGGGTCCCCCTCTCCCCCTGGCGCCCTGCCCCGACCTGCGCAAGCACAGGACGTCGTGG  
L G P L S P G A L P R P A Q A Q D V V D  
181 ACCTGGACTTCTTTCACCCAGGAGCGCTGACCTGGTGAGCCCTCGTTCTCTGTCCTCA  
L D F F T Q E P L H L V S P S P L S V T  
241 CCATTGACGCCAACCTGCCACGGACCGCGGTTCTCTCATCTCTGCGGTTCCTCAAAGC  
I D A N L A T D P R P L I L L G S P K L  
301 TTCTTACCTTGGCCAGGCTTGTCTCTGCTACCTGAGGTTTGGTGGCACCAGACAG  
R T L A R G L S P A Y L R F G G T K T D  
361 ACTTCTTAATTTTCGATCCCAAGGAATCAACCTTGAAGAGAGAAGTTACTGCAAT  
F L I F D P K K E S T F E E R S Y W Q S  
421 CTCAAGTCAACCAAGATATTTCAAATATGGATCCATCCCTCTCATGTTGGAGGAGAAGT  
Q V N Q D I C K Y G S I P P D V E E K L  
481 TACGGTTGGAATGGCCCTACCAGGAGCAATTGCTACTCCGAGAACACTACCAGAAAAGT  
R L E W P Y Q E Q L L L R E H Y Q K K F  
541 TCAAGAACAGCACCTACTCAAGAAGCTCTGTAGATGTGCTATACACTTTTGCAAACCTGCT  
K N S T Y S R S S V D V L Y T F A N C S  
601 CAGGACTGGACTTGATCTTTGGCCTAAATGCGTTATTAGAACAGCAGATTTGCACTGGA  
G L D L I F G L N A L L R T A D L Q W N  
661 ACAGTTCTAATGCTCAGTTGCTCTGACTACTGCTCTTCCAAGGGGTATAACATTTCTT  
S S N A Q L L L D Y C S S K G Y N I S W  
721 GGGAACTAGGCAATGAACCTAACAGTTTCTTAAAGAAGGCTGATATTTTCATCAATGGGT  
E L G N E P N S P L K K A D I F I N G S  
781 CGCACTTAGGAGAAGATTATTTCAATTGCATAAATCTTCAAGAAAGTCCACCTTCAAAA  
(T)  
Q L G E D Y I Q L H K L L R K S T F K N  
(F)  
841 ATGCAAAACTCTATGGTCTGATGTTGGTCAGCCTCGAAGAAAGACCGCTAAGATGCTGA  
A K L Y G P D V G Q P R R K T A K M L K  
901 AGAGCTTCTGAAAGGCTGTTGAGAAAGTGAATGATTGATTGATTGATTGATTGATTGATT  
S F L K A G G E V I D S V T W H H Y Y L  
961 TGAATGGACGGACTGCTACCAGGGAAGATTTTCTAAACCTGATGATTGGACATTTTAA  
N G R T A T R E D F L N P D V L D I F I  
1021 TTTTCATCTGTGCAAAAAGTTTTCAGTGGTTGAGAGCACCAGGCTGGCAAGAAGTCT  
S S V Q K V F Q V V E S T R P G K K V N  
1081 GGTTAGGAGAAAACAGCTCTGCATATGGAGGGGGAGCGCCCTTCTATCGACACCTTTG  
L G E T S S A Y G G G A P L L S D T F A  
1141 CAGCTGGCTTTATGTGGCTGGATAAATTTGGGCTGTGAGCCCGAATGGGAATAGAAAGTGG  
A G F M W L D K L G L S A R M G I E V V  
1201 TGATGAGGCAAGTATTTCTTGGAGCAGCAAACTACCATTAGTGGATGAAAACCTTCGATC  
M R Q V F P G A G N Y H L V D E N P D P  
1261 CTTTACCTGATTATTGGCTATCTCTTCTGTTCAAGAAATGGTGGGCACCAAGGTGTTAA  
L P D Y W L S L L F K K L V G T K V L M  
1321 TGGCAAGCGTGCAAGGTTCAAAGAGAAGGAAGCTTCGAGTATACCTTCATTGCACAAACA  
A S V Q G S K R R K L R V Y L H C T N T  
1381 CTGACAATCCAAGGTATAAAGAAGGAGATTAACTCTGTATGCCATAAACCTCCATAACG  
D N P R Y K E G D L T L Y A I N L H N V  
1441 TCACCAAGTACTTGGGTTACCTATCTTTTCTAACAAGCAAGTGGATAAATACCTTC  
T K Y L R L P Y P F S N K Q V D K Y L L  
1501 TAAGACCTTTGGGACCTCATGGATTACTTTCCAAATCTGTCCAACCTCAATGGTCTAACTC  
R P L G P H G L L S K S V Q L N G L T L  
1561 TAAAGATGGTGGATGATCAACCTTGGCACCTTTAATGGAAAAACCTCTCCGGCCAGGAA  
K M V D D Q T L P P L M E K P L R P G S  
1621 GTTCACTGGGCTTGGCAGCTTTCTCATATAGTTTTTTTGTGATAAGAAATGCCAAAGTTC  
S L G L P A F S Y S F F V I R N A K V A  
1681 CTGCTTGCATCTGAAAAATAAATACTAGTCTGCACTG  
A C I

2/34

FIG. 2



3/34

FIG. 3A




FIG. 3B





FIG. 4



5/34

FIG. 5A





FIG. 5B





6/34

FIG. 6A





FIG. 6B



7/34

FIG. 7A





FIG. 7B



8/34

FIG. 8A





FIG. 8B



9/34

FIG. 9A





FIG. 9B



10/34

FIG. 10A





FIG. 10B



11/34

FIG. 11A










FIG. 11B





mouse	CTGGCAAGAAGGCTCTGGTTGGGAGAGACGAGCTCAGCTTACGGTGGCGGT	50
human	CTGGCAAGAAGGCTCTGGTTAGGAGAAACAAGCTCTGCATATGGAGGCGGA	115
mouse	GCACCCTTGCTGTCCAACACCTTTGCAGCTGGCTTTATGTGGCTGGATAA	100
human	GCGCCCTTGCTATCCGACACCTTTGCAGCTGGCTTTATGTGGCTGGATAA	1165
mouse	ATTGGGCCTGTCTAGCCAGATGGGCATAGAAGTCGTGATGAGGCAGGTGT	150
human	ATTGGGCCTGTCTAGCCCGAATGGGAATAGAAGTGGTGTGATGAGGCAAGTAT	1215
mouse	TCTTCGGAGCAGGCAACTACCACTTAGTGGATGAAAACCTTTGAGCCTTTA	200
human	TCTTTGGAGCAGGAAACTACCACTTAGTGGATGAAAACCTTCGATCCTTTA	1265
mouse	CCTGATTACTGGCTCTCTCTTCTGTTCAAGAACTGGTAGGTCCCAGGGT	250
human	CCTGATTATTGGCTATCTCTTCTGTTCAAGAAATTGGTGGGCACCAAGGT	1315
mouse	GTTACTGTCAAGAGTGAAAGGCCAGACAGGAGCAAACTCCGAGTGTATC	300
human	GTTAATGGCAAGCGTGCAAGGTTCAAAGAGAAGGAAGCTTCGAGTATACC	1365
mouse	TCCACTGCACCTAACGTCTATACCCACGATATCAGGAAGGAGATCTAACT	350
human	TTCAATGCACAAACACTGACAATCCAAGGTATAAAGAAGGAGATTAACT	1415
mouse	CTGTATGTCTCGAACCTCCATAATGTCAACCAAGCACTTGAAGGTACCGCC	400
human	CTGTATGCCATAAACCTCCATAACGTCAACCAAGTACTTGCGGTTACCCCTA	1465
mouse	TCCGTTGTTTCAGGAAACCAAGTGGATACGTACCTTCTGAAGCCTTCGGGGC	450
human	TCCTTTTCTAACAAGCAAGTGGATAAATACCTTCTAAGACCTTTGGGAC	1515
mouse	CGGATGGATTACTTTCCAAATCTGTCCAACCTGAACGGTCAAATCTGAAG	500
human	CTCATGGATTACTTTCCAAATCTGTCCAACCTCAATGGTCTAACTCTAAAG	1565
mouse	ATGGTGGATGAGCAGACCCTGCCAGCTTTGACAGAAAAACCTCTCCCCGC	550
human	ATGGTGGATGATCAAACCTTGCCACCTTTAATGGAAAAACCTCTCCGGCC	1615
mouse	AGGAAGTGCACTAAGCCTGCCTGCCTTTTCTATGGTTTTTTGTGCATAA	600
human	AGGAAGTTCAGTGGCTTGCCAGCTTTCTCATATAGTTTTTTGTGATAA	1665
mouse	GAAATGCCAAATCGCTGCTTGTATATGAAAAATAAAA	637
human	GAAATGCCAAAGTTGCTGCTTGCATCTGAAAAATAAAA	1702



FIG. 14





Figure 15



16/34

Figure 16

ggatcttggctcactgcaatctctgcctcccatgcaattcttatgcatca	50
gcctcctgagtagcttggattataggtctgcgccaccactcctggctaca	100
ccatgttggccaggtggtcttgaactcttgggctctagtgatccacccg	150
ccttggcctcccaagtgcctgggattacaggtgtgagccatcacacccg	200
cccccgcttccatattagtaactcacatgtagaccacaaggatgcaacta	250
tttagaaaacttgcattggtccacttttcaaatcacccaaacatgttaaa	300
gaaattggtatgactgggcatggcacagtggctcatgcctgcaatcctag	350
cattttgtgaggtgagacgggcagatcacgaggtcaggagattgagacc	400
atcctgacagacatggtgaaatcccatctctactaaaaatacaaaacaat	450
tagccgggggtgatggcaggccctgtagtccagctactcgggagggtg	500
aggcaggagaattggcgtgaatccaggaggcagagcttgagtgagccgag	550
atggtgccactgcaactccagcctgggcagacagagcgagactccgtctcaa	600
aaaaaaaaaaaaaagaaagaaattggtatgactgttgactcacacaggag	650
tcaggggcatggggtgggtgtaagattaatgtcatgacaaatgtggaaa	700
agaaacttctgttttccaaactccacgtctgctaccataattattacactc	750
ttctggtagtgtggtgtttatgtgtgaatttttttcatatgtatacagt	800
aattgtaggatatgaacctgattctagtgtgcaaaactcactatgagctta	850
gcttttaagttgcttaagaataggtagatctatgcaataatgataatta	900
ttattattattttaagagaggggtctcactttgtcaccagggtggagtc	950
agtgtgtgtgattaagggctcactgcaacctccacctccagggtcaaataa	1000
acctccacctcagcctcccgtagctggaaccacaggcagggccacc	1050
acgcctggctaaattttttgtattttttgtagagatggggtttcatcatgt	1100
tgcccagggtgttcttgaattcctcgggtcaagcaatcctccaccttg	1150
cctcccaaaatgctggcatcacaggcatgatggcatcactggcatcacat	1200
accatgcctggcctgatttatgcaaatagatatgcatttcaaaataatc	1250
tatttttattttgttccttattggtggtacaatctcaagtggaaaaatct	1300
aagggttttggtgttattttgcttactcaaccaatatttattagactctta	1350
ctaagcaccaatgatcacatgcctgagctatggctagcatagcgtgtg	1400
agacaaacttaactctgttttggtggagcatataatctagttagatgaag	1450
ccaatgttgagcaacatcacataactaacaattgaggatgtctacgagag	1500
tgtctaacaaattgaggatgctacgagagtggtctaacaaattgaggatgc	1550
tatgagagtggtgtcatggagagctgcctggagattgagagaaagcttcc	1600
tgagggaagtacatttcagctgaaacacactgccatctgctcgagggttt	1650
tgtaaactgcattcacatcccgattctgacacttcacatcccgattctgac	1700
acttcacccagttactgtctcagagcttgggtccgcatgtgtataaacag	1750
gacagtatgcacttggcagggttgtgagaagggaagagaacacaagtaaa	1800
gcacctgtatcaggcatcacagtaggcactaagcgtgcgatgtctgtatg	1850
attatacatcagtgtaagcatcaaggaaaagctgaagaaaagcttgacca	1900
acagcgaaagataaatgcgcagaggagaaatttggcaaggctccaaatt	1950
caggggcagctccgtactctacactttgtatgggggcttcaggtcctgagt	2000
tccagacattggagcaactaaccctttaagattgctaaatattgtcttaa	2050
tgagaagttgataaagaattttgggtggttgatctctttccagctgcagt	2100
ttagcgtatgctgagggccagattttttcaagcaaaagtaaaatcctgag	2150
aaactgcctggccagaggacaatcagatttttggtggtcgaagtgaacag	2200
caagtgtttataagctagatgggagagggaaggatgaatactccattgga	2250
ggttttactcagggtcagagggataaccggcgccatcagaatgggatct	2300
gggagtcggaaacgctgggttcccacgagagcgcgagaacacgctgcgtc	2350
aggaagcctggtccgggatgccagcgctgctcccggggcgtcctcccc	2400
gggcgctcctccccaggcctcccgggcgcttggtcccggccatctccgc	2450
acccttcaagtgggtgtgggtgatctcgttaagtgaacgtgaccgccaccg	2500
aggggaaagcgagcaaggaagtaggagagagccgggcaggcggggcgggg	2550
ttgattggggagcagtgaggagggatgcagaagaggagtgaggagggatgga	2600
gggcgcagtgaggaggggtgaggagggcgtaacgggGCGGAGGAAAGGAGAA	2650
AAGGGCGCTGGGGCTCGGCGGGAGGAAGTGCTAGAGCTCTCGACTCTCCG	2700
CTGCCGGCAGCTGGCGGGGGAGCAGCCAGGTGAGCCCAAGATGCTGCT	2750
M L L	
CGCTCGAAGCCTGCCTGCCGCCGCCGCTGATGCTGCTGCTCTGGGGC	2800
R S K P A L P P P L M L L L L G	
CGCTGGGTCCCTCTCCCTGGCGCCCTGCCCGACCTGCACAAGCACAG	2850

Fig. 16 (continued)

17/34

P L G P L S P G A L P R P A Q A Q  
GACGTCGTGGACCTGGACTTCTTACCCAGGAGCCGCTGCACCTGGTGAG 2900  
D V V D L D F F T Q E P L H L V S  
CCCCTCGTTCCTGTCCGTACCATTTGACGCCAACCTGGCCACGGACCCGC 2950  
P S F L S V T I D A N L A T D P  
GGTTCCTCATCCTCTCTGGGgtaagcgccagcctcctgtgctcctcctt  
R F L I L L G  
tctgtcctcctgacacctatgtctgccccgcccagcggtctccttcttt 3050  
tgccggaagaaacacttcacaccggaacctccccgcctgtctctccccacc 3100  
ccacttccccgcctctcatttctccctctcctccttactctcagacccca 3150  
aaccgctttttgggggtatcatcttaaaaaatagatttaggggttacaag 3200  
tgcagttctgttccatgggtatattgcattgtgggtggcatctgggctctt 3250  
agtgttaactgtcacccgaatgttgatcattgtatctaataaggtaatttct 3300  
catccctcatccctctccaccctccacccttttgaggtctccagtgtct 3350  
actattccactaagtccatgtgtacacattgttttagcgccactctaaat 3400  
gagcctttttgtttcattcattctgttaagtgttgaataggcaccaccta 3450  
ggtcaggtataagtggaaatttgaaaaaagaaactgccacttgccccagt 3500  
acttccctagccaagaggaggaaaccaggcaggtgcacctgaaggcctg 3550  
tgagtgtctgatttgctgtgtaggttaggacaagtaagattgtgcatagc 3600  
cttctgtatttaagactgtgttaggaagatttctctttctttcttttct 3650  
ttttctttttctttcttttttttttttttaggcagatgaaaaggcgctca 3700  
cagaacaggaataaaaaatctaaatattcaataaatgagacctaggagact 3750  
actgcagtgaacttacaagtcttaataaaaaagatgtctctccaaatggg 3800  
gctgcaaaatgtggtgctgccttatcagctctaaagtttttctctacctg 3850  
agaaagaaggaacctgatgcaggttcagggtcctgccccatgaatgcag 3900  
gctgactccaagatggggagctacagggaacaatcccaggtcttctaggcc 3950  
tcttatttaggcctgggagcctccagagatggccacatcttgaccagcc 4000  
cagatagagggaagatcacatttatctcacctctgtgtcaaatacctag 4050  
atgctgtctcctccctgagccacactatagttgcccagcgctaatttaagg 4100  
gtagtgtactggtaagagatggacagaccatcctggcttgactctcagc 4150  
tctggcaaaagatgagtgcattgggttttccatattctctggccacaccaa 4200  
ccttgatttcttcagctgtagaatggaaatttctcaagcttgccctcaagga 4250  
ttattggccgaggtattgatgatattgtaagagcttctcagtggttgacc 4300  
catagtaagtgtttgacgtttcaaacgaattgtttctttcttaggacatgg 4350  
tgagcatttggtagccattcacgggttttctgtttctttggatcatagtt 4400  
aacctctccttttcttcttctggcactacaattttctgggtggggaagaatcc 4450  
ttaactttctgccccttcccttaaggataggaagctgatactaggcagcaa 4500  
ctagttgggggataggaagattgttccagagaaatgctgaaccatagggc 4550  
tccagatcacaggacccagctcttagctgtggtgggtggtgggtggggg 4600  
ggcggttactgaacatgggtatgaagtagatgtccatttactgaaatgt 4650  
gaggacctgaggcctcttctattgtctgtagccagcatattccccaacctc 4700  
tccccaaagaaaggacagatgggggttccccctggagtaacaggtccaaa 4750  
agaaaaaacatacagtgaggacttccaggatctgggctgatcaccagca 4800  
gtcaagctccccgaattgactaaacccccctaaacagtagaaattcca 4850  
atctgcaatttagtgaggatgatacctttattcttcttaatacatctct 4900  
tcatttcccagagcaccttttttccctcctctgcacctttttgttaaa 4950  
gactggagtataatgaaataccaagagagcataacatgtgatacataaaa 5000  
ctttttttctggttttcaaaaacagttcattcttgtccatacgtgcttctc 5050  
tccaaggctggtgctgtctgttccagcccgttctgcttgagaggccat 5100  
ctgccatacctgctccccagacgcatcgacaagcaccccagagtggtat 5150  
ctgctaagacctaaaagaggagggaacccccctctcctcatctaagacct 5200  
gcttctaaattagagtgtgagggtccatctccccaggagggggacaggggc 5250  
ccaaacagcccagccatctcagaagacaacactaagctttgtagggggtcc 5300  
acagttagaggagagtaagacgcctgttgttttaattttattacagttcctca 5350  
aaagtgaagatgtgtggcgggatggcaagagctgagcagacgaaagctg 5400  
aagggaataaggaaagagaggaggacacaaacagctgacacttccctcagtt 5450  
cttgtcatttgcctggccctgttctaagcacttctaggtattaatccat 5500  
ttagtttggtcatacaacactgtgagtaactagttttgtcacccttcttt 5550  
aaaaatgaagaaagtgaggtcagggaggttaagtaacttgccacagtt 5600  
tgaaactagactctgatcacatgagataatagtgccataaaaaagggaag 5650  
gcagatttatatttttaaggaaagagagtaggatattggtagaaaaagat 5700

Fig. 16 (continued)

18/34

tgtttggaagggaattgagagattgatataatgaaaagaagcattcacat 5750  
gagagttaacagtatcagggcccaaaccttcatctaaggtacttcaaagag 5800  
gcctaagcaaaccttagtcactggcgtggttctagtctccatgatggcaaa 5850  
tacattgtgtacagcccaactccacacaaaacttaaataccaatgataga 5900  
gcaatctaaaatttgaagaaaaaatctttcaatttgcgtcttcccaga 5950  
gggacttaatcaagaaaccaatcaaaaacttccctaagcctaactgtgtg 6000  
cagaactccaaagagagccagccctaaatcaacactgtccaatggaaat 6050  
ataatataatgtgggcctcatatgcaaggtcatatgtaattttaaatttt 6100  
ctagttagccatatataaaaggtaaaaagaaacaagtgaatttaatttta 6150  
taattttatttagttcaatagatccaaaatgttttctcagcatgtaatca 6200  
atataaaaatatataatgaggtatttattatttctctctcaaaccaagtc 6250  
tattctataatctggcgtgtattatttacagcacttctcagactatattt 6300  
ctttctttcttttttttccgagacaatttgcgtctgtcacccaagct 6350  
agagtacaatggcgtttacctcggctcactgcaacctccgctcccgggtt 6400  
caagttatttctcctgcctcagtcctcccaagtagctgggactagaggcatg 6450  
caccaccgcgctggcctaattgtgtatttttagtagagacagggtttcac 6500  
catgttggccaggcctaattctcaaacctcctgagctcaggtgatatgccac 6550  
ctcggcctcccaaagtgttgggattacagcgtgagccactgcacccggc 6600  
ctcagattaactatatttcaagcgttcagtagccacatgtagctagtgtct 6650  
atggtagtggacagtagacagatctgcatttcaattaaagacagtatacaag 6700  
catagttcactaatgacaggttaaaaaaagtatagtgctgagtcgggtgt 6750  
agaaatccataactgcagagcaaaagtggtagaacagcaatctcagtc 6800  
gataatgcaaccatgcttgcctttcattgcaatttgccttatttctctca 6850  
gcaagtttcatccatttttggcaattcaataaatatttactgataaaaaac 6900  
tttcaatattagattcttgcacatcttcatagacagagttgcttttcacatt 6950  
tagaaaattacttatcaatgttaaacacacgcttttgataaccagtggttg 7000  
aaagaggtgcagactccccatgtgcctattgatggcagaaaatttcacag 7050  
ccaaaggggaaacaaagggctggggacaatcacacacctcatgtctcctaa 7100  
ctcctgggaagtgtctgctccctctgattgagctcttattattgccttccc 7150  
actaacctgtccactgtgccttgaggccctttgcagggttacctgtctct 7200  
gtcctcctcagagaatatctcctctacctccttgcagcagctacaacttg 7250  
gctattctctgatgacactgtcttccctgtagcccttttgagtaattggct 7300  
gcataattctcccatagtccagttcttttctgttctccagtcgtggtctct 7350  
ggatgacagcccaactagtttgaactccactgctatagttcaagtcctct 7400  
tttgactgtttaccttgggcaaatcacctccttttgttcagggttccttgt 7450  
ttgtaaaatgacgataataatgccatttgccttcagtggttattttgaaa 7500  
ttgagtgaagaaggcgggtagcttccctacacgctcagtgtagactagc 7550  
ctgagtgtgcattacgggtgatgccatgactcagtggttttctctcatctc 7600  
cacatctggctctcatccagtgctcctgcttacggcactctgtccccctc 7650  
ttacttactcccccttataactgaagactggcactgatctcacagtttct 7700  
ctctccacttcttagtctcaccatcatcctagatgacttcaagtcaccta 7750  
gataaaactgtctcagtttcttcaactcacattttttataacagataatgt 7800  
tacactcaagttgtaacagaaccagcttatccagctcatgaaatgtatgc 7850  
atttcatctcaactctgtattcagtgacatcctgtgggtatctggaaatc 7900  
agccatggtgagaatatttaccatggaaattggcaataactaaaaagcag 7950  
agcaccttttttctgagagccagaccatagctcttctactccatagcac 8000  
ccatcataacaatttttaaatacctccactgaacagcttcttctctctc 8050  
tacttcttccatatctgatttgagcttcttaatttatcatgtgaaccact 8100  
cttgtaataataaccaccaaatccctgttccattgttcttctctgtaaaat 8150  
actaaacctgggttagtccaaacctattttctctcttttggaatctacagg 8200  
gtggcccaaaaacctggaaatggaaaaatattacttattaatttttaagt 8250  
atattaataagccatttttaagtcttcatctccagtcctcagtgccacct 8300  
gtatagctgggctattgagctcttgcgggaggaggagtgaggacagtctcc 8350  
cagccacacagactgatgttgaccaaaccatttttttagcttccagacttc 8400  
cctggcccttaggttacccttaactctccatttctctgcctttcacatt 8450  
ctctacttttaaaaatctctgactccaccttccacttatcattcttagc 8500  
acatgaccatacttctgcttccaaagaaaatgagcaattacttcttctt 8550  
ccttttctctctgctcatcaaatctgcagacatgtcatgcctaagtccagc 8600  
tttctctcttctctgatctcagtcgtcttcttccatttctgccttgaat 8650  
ccgctccctccccaaccccccaaggacttcgctctatcagtcacctcttc 8700  
cctctctgtatcttcaactcctccatttttactggcttcttctctcaagc 8750

Fig. 16 (continued)

19/34

ctttccccaagcctttcccatctcaattacctcctcgacatgcctctgc 8800  
agaaaccaccccggtttcttccctccctcggcagcctgttcttccctgttc 8850  
tgccctcatgatggcaccatcattgtgtcactaaaatcaatctctccgac 8900  
atcatcaatggccttcccttgggtgggaaacctaataaacactttatctta 8950  
tttgggtctttgttatgggttgatgaggttaccgccaaatccatattaga 9000  
agtccaaacccccagctacctcagaatgtgactttatttgggaatagggtc 9050  
attgcagacgttattagttaggatgaggtcactggaatgtgatgggct 9100  
gcttatctaataatgactgatgtccttataacaaggagaaatttggagaca 9150  
gacacgcacataggggagaataccatgtgatgacaggagttatggagtgg 9200  
agtcaaaaagctatgggaacttgggagaaagacctggaaacaaatcccttc 9250  
ctgcgccttagagagggagtatggccctgccactaccttgaattcaacgtt 9300  
tcggcttttcaaaactgttaagacaatacatttctgtgttcaaaccaatt 9350  
agtttgcagtactctgcgactgcagccctaacaactaatacagtcctct 9400  
ggaggcattttggcaaggttgacaatggaagcactttcttacccttttagg 9450  
tctgtcgcctttcttgttggggggtgttttctaacaattcctctccatct 9500  
ctctctctctagtttcttcttaaacattgtgttcttcagacttctgacct 9550  
aggccttcttttcacttcacatattccctgggtgtctcaccacacttcc 9600  
agaaattacttaaatctctgctcatgcagtactgtgctggaaactgttta 9650  
acaactggctctctgggaagaggggagactgggtgatgggttttgtgtat 9700  
ttctgtggtgttaataactccctccatggccaattccaaactgccaaact 9750  
ttaacaactggctcacaattttctccaaatttaacatttggctttcaca 9800  
ggccaacaactgggtcagccaactccagcacacctctgcttttgtgtca 9850  
gagagaagtaacttattttgtacaaaaggtaaaaaaaaacacctgcag 9900  
gcccccttttttcttaacaactgctctagaatagaatagctgaagc 9950  
ttcttttatgcattctctgttatttccatgtcactgtgtgtgtgggatt 10000  
atttttcccttatttttcttgtatatggttgaaatactgtacctttgatc 10050  
agttttagttttttaggcattgttttgaccatattaaatctagtttttgt 10100  
cagagggcgtcaatatttttctcaaaaacagaaaatatttcatgtgcaa 10150  
aggagacaaaacaaaagggtccttaataacaaaactttgaaatgtgatctc 10200  
ttgtacttggcagtgctccaagtggtaaaccccaaacagtattgggttttca 10250  
ttttgttcaggaaaagtctttgtctggcagcgacttacccttacatcaggc 10300  
gggccttgcctcattcattcacttaagtatttattaaacaccagcggtgtg 10350  
ccaagtacttatctaggtatcgggtagattctgataagtcagtcaggctcc 10400  
ctgctctcagggagcttgagcagagatgggggctgcaatagagagtaag 10450  
ccaaggaaaatgaaaagggaagtgtattttagagagtgatgaatgctatga 10500  
agaaaatgaaggcagcgagtgatggagagtgaccgaaggtgtgacag 10550  
tttgtacctctaaggaccagactgtgacccaggtcactcacagatgcccg 10600  
tcagtgtagccacagcaacttttccaggtgctcgtttcctcccaacttcc 10650  
cagtctcttgcacgcccgcgactgcttacaatacagctagagggaatcta 10700  
aatgaggttccctctatcatcaaacccaatcaaaatgccaagggaacagaat 10750  
cagtgcctggctgaaggcagtggaacagggccagcctggagtggttctct 10800  
ctgagggaagttccctcatcttgggttttagggccataccttgtgacctgtga 10850  
gctaggggttgccagtccttgacatttctactgaggactcgcctgtctat 10900  
attcccgccctgtatgtgtctcctgagttccagacacacagggcgaagcg 10950  
cctgatggatggaaagtatttttgggtgttccattggtatctcaaatc 11000  
tacaacaaacttagtgccttctctcctcctgttccctcccatcttcagttc 11050  
atcacctgttccctcatccagcaaatgatattaccatcttccaaggagctt 11100  
cccaggagtaatccttgactcctcctcaacatccaattaataatcaaatc 11150  
taggccaggtacaatagctcacgcctataatcccagcactttgggaggct 11200  
gaggcaggtggatcattttagggccaggagtcaagaccagcctggccaac 11250  
aaggtgaaacctgtctcatttataaaaaagttatttataaaactcaaatct 11300  
attatttctacctctaagtgtgtcttgaatttatccatctctctccatct 11350  
ctgagctgttaccttaacctcagtcacatcacgttttgtctacgttaacatg 11400  
accagagctcttgttcttagtctgggtgaggtcactccagctgcttcagatc 11450  
cttccatggctcaccgttgccctcatataaagttggcactcctggacatg 11500  
tggcttacggggccctccgtgatgtggccctatttgcctctccattctgt 11550  
tctctccagcctctctgccccatctctaggcaccacacacacccctct 11600  
gctcgtcaatgggtgccagcttctctctctatctctgttcttggacagact 11650  
ttcccttcacctggaatgctttcttcaatccctaccccaactctcttaat 11700  
ctagataaggtttattcttttgaatgtctagcagtgaaccatttcccc 11750  
tgaaaaaccttctctaaccaacccctaccctcagcccaaggctctagatt 11800

Fig. 16 (continued)

20/34

aggagtcctctgaaatgtttccatagcatttttaagaattgcctattta 11850  
cttggttcgtatctatcactaaactacaaattgtatgagaacagccactat 11900  
ctctgcctgggttcaccattcatctccagcaactagcataatgcctggcag 11950  
agtcagcctgcaacaaatatttgttgaataaattaacagatggctttatc 12000  
tccttaagtaaattcttgctttttcacctattaaaacagacgcacaggcc 12050  
agggtgtggggcccatgcctgtaatcccagcactttggcaggctgaggtg 12100  
ggcggatcacctgagggtcaggagttcaagaccagcctggccaacatggtg 12150  
aaaccccatctctataaaaaatacaaaaattagctgggcatggtggtggg 12200  
tgcgatatagtcacagctactagggaggctgaggcaagagaatcgcttgaa 12250  
ccagagggcagaggtggcagtgagccgagatcatgccactgtaçtccag 12300  
cctggatgacagagaccctgtctcaaaacacacacacacacacacacaca 12350  
ca 12400  
taacgtgcttgttatggaacacttgaataacaggaagtaataaaaaa 12450  
gtctaccatctagctcaccacataatgaccattgctatcatcctggcata 12500  
attctctcctgtatataaataataattcttttattgttataaattacacta 12550  
tgagtactattttatttttactgtggcaaaatgcgcaaaacataaaat 12600  
cttgccattttaaggatgagcttgggtgacattcaccacactcacattgt 12650  
tgtgcaaatatcaccactatctatctcagaacttctcgtcttcccaaac 12700  
tgaaactctgtacccattaaacaatagtgatcctctgcttttccctccc 12750  
tacaattttatttttatttgggtttgtacaaactgaaaatagctgcttct 12800  
tccttacttagttcagattagcatttccatttattttagccgtggttttga 12850  
ggatgccatgacagatgccatccttccctagagctcttggggctgtcagg 12900  
tatttcagtcagggtgaattcgggttgataacattttaaaatctcacttt 12950  
attctgaggttccatgtgtcagagcccacgtatttttagggactcccaa 13000  
gttacaacaaaaaatatggtgagggaatcactgaagtttaacacaag 13050  
agacttacattttgttcaatttctatcttttagtttatttccctaaacata 13100  
aagaaatactttgaaaattttacatagcattatacatatttaattaaagca 13150  
tgagcacatcttaaaacttttaatttttagatcagatctttaattccctagg 13200  
atataaagggttactggcaatttggccagggtgtggtggttcacgcctata 13250  
atcccaacactttgggagggtgaagtggggaattgctagagcccaggag 13300  
gtggagggtgcaatggcctgagatcacgccatcgactccagcctggatg 13350  
atgagaatgaaatcctgtctcaaaaaaaaaaaaaaaaaaaaaaagaagaa 13400  
gaagaagtattggcaatcagtgctccaggaataatttccctgacttgaaat 13450  
aaacctacatgtagacaaactaattaggccattccaagagttgctagcat 13500  
tggtttaatatgttttcagagcattccaggaagcagtggtggccagcattg 13550  
catgtttgatacttcagaaatgtatgacaggtgtttctcttaccagggtc 13600  
ttctgttttcttagttttgctcatgtataatatttgaacatcctcatct 13650  
ttttgaggggaaggattatagatcattctaatccattttctagcatttg 13700  
gtaccattctaaacacatgataggcacccatttggagcatttttggcttg 13750  
acagaatatgcaatttagaattgttcaaatagaggtgtcagtgatgggaa 13800  
tagaataactatataattctaagtcatttgacttaaatcaaaaagaatga 13850  
ttttccttggtgggaatgggtgaaggaggaggaggttaagaaggaggaga 13900  
agagatcctaagtcattttataaaacttctctggaaagacaggtgtgtgaag 13950  
actttttaaaaagtcatttcaccaaattgtgtgtgtgtgtgtgtgtgtgt 14000  
ttaaatagactttatttttttagagcagtttttaggttcacagcaaaattga 14050  
atgcaaggacagagatttccataaaacccctgccacacacatgcatag 14100  
cctccctcattatcaacatccccaccagagaggtgtttgttctagttagat 14150  
gaacctacactgacacatcattatcacccaaagtccatagttcacggcag 14200  
ggttcactgtcggtgtacattctatgggtttgagcaaatgtataatgaca 14250  
tgtatccaccattatagtaacatacagagtattttcagtgccctgcaaat 14300  
cccctgttctccacctattcatccctccctctctgcatttccacccccag 14350  
cccctggtaacccgtgatctttttactgtcccatagtttcggacgatcta 14400  
tttttcagacagacacagagctgtcttcccttagtttctattctatcat 14450  
ttctttctccccatccataaaaaggctatgagtttttttaagtgttg 14500  
aacaccatcctactgttcaagttaaaaacataagctcctggctgggtacag 14550  
tggctcatgctgttaattctcagcattttgggagggtgtggcagaagcatc 14600  
acttgaagccagaagtttgagaccagcctgggcaacatagcaagacccca 14650  
tccctccacacacaaacacacacacacacacacacacacacacacacaca 14700  
cacacacacacacaaaaaacaagctcttgccagaattagagctacaaattg 14750  
ccctcaggttccatagaagatcagtccttcaattagattcagattgagatg 14800  
cttctcttttaacaatgattccctttctatcatgcccaataagaaaac 14850

Fig. 16(continued)

21/34

aaataaaaattaaacaatactgcctgtaattctcagctacccaggaggcag 14900  
 aagcagaactgcttcaaccggcaagcagaagttgcagtgaaagtgcagatc 14950  
 gcgccactgcactccagcctgggaaacagagcaagattctgtctcaaaaa 15000  
 caaaacaatgtgatttctcctcctaagtcctgcacagggaatgttaaga 15050  
 aataggtccaccaggaaagaaggaagtaagaatgttgactagattgtct 15100  
 tggaaaaaatagttatactttcttgcctgtcttcttaacagTTCTCCAAA 15150  
 S P K  
 GCTTCGTACCTTGGCCAGAGGCTTGTCTCCTGCGTACCTGAGGTTTGGTG 15200  
 L R T L A R G L S P A Y L R F G  
 GCACCAAGACAGACTTCTTAATTTTCGATCCCAAGAAGGAATCAACCTTT 15250  
 G T K T D F L I F D P K K E S T F  
 GAAGAGAGAAGTTACTGGCAATCTCAAGTCAACCAGGgtgaaaattttta 15300  
 E E R S Y W Q S Q V N Q  
 aagattcactctatatattttaattaaacgtcagtcctcatgagaatgcttt 15350  
 gagaaaactgttattttctcacacctaacaattaatgagattaacttctc 15400  
 tccctcatctgcactgtggaggaatctgaacaaggaggaggagcagtg 15450  
 gcagggttctctatcatgatgtttgtcatgttcagtgtaggagcctcaca 15500  
 aaaaaaaataaaaaaaaggcgtcctggatataactgagagctcattg 15550  
 tacagtaaatattaataaaaacagtgattgttagctgaaggatagaactgct 15600  
 tggagggagcaagtggttagaatcgcgctcaaaactaaagagcatttctagc 15650  
 caaagacacaatgatagattgaaggatatttattctaaatatagaatatg 15700  
 ggtgaacgagatctgtggacttctgggctccaacgcttagattctgatttt 15750  
 agcaagcttctcaggggattctgatattgaaaggctgtggccttcacctg 15800  
 agaaacctgcccagggggcatgaaaaatttgcctgtcttcagaagtg 15850  
 ctatcacacatcaaatggaaagttaaatcgtatcttaacaattactaggat 15900  
 gggcgagtgactcacacctgtaatcccaacactttgggaggctgaggca 15950  
 ggaggatcacttgagcccaggagttcgggaccagcctgggcaacatagag 16000  
 agacgttgtctctattttttaataatttaaagagaaaaaatactgaaaa 16050  
 tattgtatacacacctgaattataataatgtgtatataatgtatatattc 16100  
 attatgaggaatatttgattatttcatatattatatttcttctctgtt 16150  
 tattttatccagttatgaagtattttagaacaattcatcagtaattggggc 16200  
 taaattgacagaaatagtaatacagagaaaaatagaaaaagacagatgggtta 16250  
 tctttgaataaccagggttgagttgtttatgggtttgtttttgttttggg 16300  
 ggcggttttttagacagagtcacctctgttgccaggctggagtgagtg 16350  
 ggcacaagcatggcccactgcactccttgacctcttgggctcaagcaatct 16400  
 tcccaccttagcctcctgagtagctgggaccacaggtgcatgtcaccaca 16450  
 cccagctaattttttttttttttttagagacagtcctttctatgttatcca 16500  
 ggctgatctcaaaactcctgcactcaagtgatccccctgccttggcgctccc 16550  
 aaagtattgggattatagccatagccaccacaccccaacctagtttctatt 16600  
 tagacttggccctttcccaccagtcatttgtgtccaaaagatctcataaa 16650  
 tgtagacaggaaactgtcctttgtctcatcagttttcttcatcctgtgtct 16700  
 agggggatgggtgggtgggggaaactgggggttatgcaagttcctctgaaac 16750  
 atcctctgtgagcccagggtggatgaggcaccagccgagcgagtcag 16800  
 tgtgcagctttccagaaaggaagtcacagccagtcagccggccctggca 16850  
 gccagcaccgggcaaccctgctgtcttgtgataaagaaatggtctgcctg 16900  
 acaggatgggtggtgatttttcttttttcttttttttttttgagacagg 16950  
 gtcgtgctctgtcgcaccaggctggagtgcaatggcggtatcttggtcac 17000  
 tgcagcctctgcctcccaggctcaaggcatcctcccacctcggctctccg 17050  
 agtagctgggaccacaggcacacacaccacgcaccaactaagttttcgta 17100  
 ttttagtagaggcagggttttactatgttgcaggctagtctcaaaact 17150  
 cctgagctcaagctatccatctgccttggcctcccaaagagctggaatta 17200  
 caagcgtgagccactgtgcctgaccagggtggattttttcaagtgcacat 17250  
 gttgtggtccagaagctctgatggtaccaaattccaagcgaaaaaaagt 17300  
 caatgggtccaccacctacacctcccatgatggcaagaggaaatcacca 17350  
 cactgcagatacagtcacatgtaaaacaaattgctatggattttgaaagt 17400  
 aaccttaagagaactgcactatgttttcttcattagagttctctggtaat 17450  
 tccagcttttttttttttttttttagacagtgctcgccttgtcgccc 17500  
 agtgtcaccaggtggagtgacgtgacgtgatctcggtcactgcaacc 17550  
 tccgcctcgtgggtgaagtgattctcctgcctcagcctcctgagtagct 17600  
 gtatttttagtagagacgaggtttcaccaatttggccaggctggtctcgaa 17650  
 tccctgacctcaagtgtattccgcccatctcagcctcccaaagtgctgggatt 17700



Fig. 16 (continued)

22/34

acagggtgtgagccactgcacccggccagtaatttcaagcttctgaggagc 17750  
 cctttgaattgtttaaataaactttagctatgtccaacatatccatgttca 17800  
 gtgtatgttcgatatttcttaggaaacctgccccttggtgttttctttgt 17850  
 ggtaattcatgagccggcaaatgtgacatgtgttacagaatataacctttt 17900  
 ctctgctctcctacctcataaccagaacttaattatcctgcttttagtcac 17950  
 ataaatagctaactaaataaataatgatgagatttcagtcctcactgtga 18000  
 aaatagaccttctaaatgatctcttccacttgcagATATTGCAAAATATG 18050  
 D I C K Y  
 GATCCATCCCTCCTGATGTGGAGGAGAAGTTACGGTTGGAATGGCCCTAC 18100  
 G S I P P D V E E K L R L E W P Y  
 CAGGAGCAATTGCTACTCCGAGAACACTACCAGAAAAAGTTCAAGAACAG 18150  
 Q E Q L L L R E H Y Q K K F K N S  
 CACCTACTCAAGtaagaaatgaaaggcaccctagagatgttccagcccca 18200  
 T Y S  
 aagatatttgaatagggttgactcgggacccaatctagcaagtcctacgg 18250  
 aagttgtataaagctgaaaaactgaagcatttcccaaatgggaaatcct 18300  
 aaactcaaaacttgcttttttggttttttggttgtttgttttcttcat 18350  
 ctgacattgcttagtagtcacagaatgaaagataaatcaatcattcatga 18400  
 tctaacaatgaccttcagtgctctctaaaaaactacggagtcagggaaca 18450  
 tgaatatattcctcatgtaaaattaaaatacagacataaaagggaacaaa 18500  
 catgaacatcattcataccttgagggtccgtccccctccagaaataaccc 18550  
 ccagatgctcttggttagagcattaagcaggaggccctgagtcactcc 18600  
 agacagtccttgaccaccaagcagcattctcttttggtttctctgtggct 18650  
 tttgcaaacacagggctagctcagctacccattagtagtgtttcagtcac 18700  
 taaaacagtcctccagtcctcaaattaggatgacattgtcacatggggct 18750  
 ttaaagcaagtgaacaaggaaccccttttttttttttttgagatgga 18800  
 atctcactcttgctgcccagcctggagtgcaatggcgcaatcttggtca 18850  
 ctgcaacctccacctccaggttcaagagattctcctgccttagcctcct 18900  
 attcattatgaggaatatttgattattcagttcctgtagggttaaagata 18950  
 taccoccatcatattattgattattgagtagctgagattacaggtgcct 19000  
 gccaccacgacccggctaattttttgtatttttttagtagagacagggttc 19050  
 accatgttgccaggtctccaggtcctgctcgaactcctgacctcagtgga 19100  
 tccaccacctcagcctcccaagttctgggattacaggcgtgagccacc 19150  
 actcctggccacaatctcttttttaactatgaaataatatttttatctgaag 19200  
 tttgatgtttatacccaactgagggatgatgttcccatatctcagttaaa 19250  
 gaaataacctgctcagatacttcaagctcttcttttgacttttgaaata 19300  
 aatgatcttgaagttactatactttgtttgggttagttaacattatttaa 19350  
 agtatattattttaatttaattatctttgttaagattttactgtatactacc 19400  
 tggagttcaatgtatcagatggatttcaaatatgtacattttttatgt 19450  
 atatggtacagaaaaaatgtgatccataagaaatcagaaaaatagcgcat 19500  
 atgctaatagtctaattgtgtcctctaaaaaacttatttttgcatttttaa 19550  
 gagggggatatactctgacactttaataagtgttaattatttgactgg 19600  
 aatttggcatgaggcagggccatttcagatcccatataaggaatgacaca 19650  
 taccagagaaccacagaagtaaggccacatttgaataaatcattatagc 19700  
 tctgctaggagaagacccagttgtattaggttaattaatggatttgctctt 19750  
 aaaacacatgtcccgaagatataggtgagtccttggggggccgcattaaa 19800  
 cattataccaatgtatcttacatttctaagaaagttttactactttacag 19850  
 gatcttttctgttaccaaaatggaaggtttccaactccaggacttggcttt 19900  
 catagttcctacaccaggggaaatgccttcttttgcatactatgcaacca 19950  
 ggttagtttagtgtaagtccagccacctgttggcaatgctaaaagggtaca 20000  
 acaaacacagaattttatttgcatgtgttaaacatttctgtgctcga 20050  
 aattttcagttttcatgggcacgtcatggaacagaaatcttctgtgttt 20100  
 agtttgggcacactactcattgtagtacaaaatatttcagaagccaatagg 20150  
 ggattccacaaattgttctgaacctgtggctgagactggtaattggctgag 20200  
 tgacatggggacataccacaaaagaagaggttagcaaaaggctgctgagat 20250  
 aaggacatgttcttagcttagctagtggtgaccttcaaacacatgt 20300  
 cccaggctgggtgctgtggctcacgcctgtaatcccagcacttggggagg 20350  
 ctgaggcgggtggattacctgaggtcaggagttcgagaccaacctggcca 20400  
 acatagtgaacctcatttctactaaaaatacaaaaaattagccaggcatg 20450  
 gtggcggggcctgtagtcccagctactcaggaggcaggcaggagaatta 20500  
 cttgaatctgggaggcagaggttgggtgagccgagattgcgccaccgca 20550

Fig. 16 (continued)

23/34

cgctagcctgggcgacaaagtgagactctgtctcaaaaaacaaaaacaa 20600  
aaaaacaaacaaacaaaaacaaacaaacaaaaacgggtatcccagaa 20650  
gatacaggtaagtttttctaacacaggtcctctgtatggtgcgttccact 20700  
taagtagaagatgacaaaaacatttgtcatgagaatatagactcacattt 20750  
taaacctgtttgagcaggaaaaggaagcaatgttacagatgtaattctgg 20800  
gtgtgactgcagaaaggtgactcccttattaaagtagtcatcctgagtg 20850  
agctaactcttgtacttctctctctctctctctctctccctcatcaccca 20900  
ttcttccgttgctacacccaggccacattggatgctgacatagactta 20950  
catgggtacagtcacagggaaagatctgccatttttttcaatgtgtcatct 21000  
tggttatcttcattccaaggatctctccactttttatacagtaagagatg 21050  
agagtctggaaggatttgggaataagataatgaattgtaagttttaatt 21100  
gttcttcgtatttttggggaaggagtaggctagggtggtccttctgttttt 21150  
ttttgttttttttttaaaagtagatgtggccagacgtggtggctcacgcc 21200  
tgtaatcccagcactttgagaggctgaggcagggtggatcacttgatgtca 21250  
ggagttcaagaccagcctggccaacacagtgaaaccccgctcttactaaa 21300  
aatacaaaaaactagccgggttgggtggcgtccacctgtagtcccagctac 21350  
tgcagaggtggaggcaggagaatcacttgaacccgggaggtggaggttgc 21400  
agtggaccgaagatcatgccattgtactccagcctgggagacagaaacata 21450  
ctctgtctcaaaaaaaagagaaaaagaaaaagaaaaaagattggtttga 21500  
actcagtcgtcaatagcctctattccaggagatgttacagttgattatgt 21550  
tatagggggtgtataatagaatttcgagctatgtaaattccaagtgcatt 21600  
tggaagaatgaagaaatggaggaagggtaaagtatgagtgaagcattcc 21650  
agggtttttgaaaaatgctataatctttgttcagggttagtacaagtgct 21700  
atthagctgtaagggtttttgtgatttacagacagttttcacatgtgtc 21750  
atttcaaccttgggtttttatggcgaaaggcatgtgatggtgcttgtcccagg 21800  
actttagatccatatctgaggttccgtgctgggcaaaagataattaccctga 21850  
tcatattatagctctataagtgggagagttgtgcctggagctcaagtctta 21900  
tgattttctgatccagggcacttccatacaacatgattttgcaatataaaaag 21950  
cctataatgtgtgactaaaagcagggtcactcaccccttgtaacagactcta 22000  
gtaatggtactgccaccaaaggctgcgtgatattgggcaaaagacttacc 22050  
ttatttgaatctcagtttccctcctagaaaaatgaggggtggaggttaagca 22100  
taggctgatgatcctaaagcctccatactgccctaaactgtggctctaag 22150  
atccagtagaatgtgtgggtcacaggactctagggagcttttcaaacccaa 22200  
atgtctgtcattccttgatggtaggcagcagtttatggaagtgggagaca 22250  
cagcaaatatcaaaatccctaaagcagcttgcaagagttgttctgccta 22300  
gtgtcttttatagttataattaaatagtttaatttttttttttttgagac 22350  
agagtcctgtctgttaccaggctgcagtgagtgccacaatctcggct 22400  
cactgcaacctccacctcccggtttgagcaattctgtctcagcctccca 22450  
agtagctgggactacaggtgcatgccactgcacccagctaatttttgtat 22500  
tttagtagagacgggtttcaccatattgggcaggctggtctcgaaactc 22550  
ttgacctcaggtgatccacctgcctcagcctcccaaagtgtgggattac 22600  
aggcatgagccactgacccagcttaaatagctaataatttaattatttc 22650  
tatagttattcaagtaattcaggccaaagacttagaaacaaaaacaaaag 22700  
ccacttttaaggagaaaagggtgtaagtttgccagatagatagagatcttt 22750  
cttttttaactacaagagttcaggaatgaattactctttaacaaacgact 22800  
atagatatacatgaaaaattggaaggacttattatgcatatgataatcaat 22850  
ttaaagacaaacacttaaaatttatattgttgccactctcaaaaagtgttaa 22900  
tagaacagctaattggtttaaaaagcagagtacagaagttcccaaaccttat 22950  
ggcaccttaatatcgagaaaaactttttaagcatgcctaggccacaaaa 23000  
aatacctgtattttgattattaaattgtaaggtctacacaaacctaatagt 23050  
aatagggtccaatagtaatgctgtccaatagatgttgatgtttttttcctt 23100  
gcaaaacttaaaagatccctacagtgctctgtaaatagcactgcctggtta 23150  
gagttgaatttcagataaaataattttttcatgttaattatttttctttt 23200  
ctttactttttttttgtttttttgtttttttgttttttttttgagaca 23250  
gggtctcattctgttggccaggctgctgtgcaatggcatgatcatggctc 23300  
actgcagccttgacctccctgggctcaggtgatcctcccacctcagcctc 23350  
ccaagtagctagctgggactacaggtgcttaccatcatgcccggtcaatt 23400  
tttgtgtttttgttagagatgtggttttgccatgttgccaggctggtct 23450  
tgaactcctgggctcaagtgatccgcccgctcgccctcccaaagtgtcta 23500  
ggaatgacaggcatgagccactgcacctggccctgggagagattttctt 23550  
aatggttacatagacatactaaacattattttattgtctatatgaagt 23600

Fig. 16 (continued)

24/34

tcaagtttaactaggtgcccctgcacttttttagttgctaatacctgtagctg 23650  
 taccatgcattcactgggtgctccccagcttgcccttgacagagtttgga 23700  
 aaccatagtcctataaactctagggcaatttttttaattgtaaaatttgattc 23750  
 attttaaattaataaataaataacaggaatttttttaaaaattgttttaaa 23800  
 tataattaaaattatcaaaatattttttaactgaacttgtagactagagat 23850  
 atttagattatgaagagtgggtttatgctaactaatgacagctctggcta 23900  
 tgcattgtggagcactgagctataaattgtggcttcccccaattctcctgat 23950  
 gtcacttgaacaaaacctaagtgtcagaccagagcttctggtatcttcca 24000  
 tgggatttccattcaacagctggagcaaatgaagtcagattgatttttttt 24050  
 aatttgtcccaattttgttgcctcaaaaacataattataatcattttattag 24100  
 aactagaatttcttcagtttaacaacagaaatagttattcattatgaaaa 24150  
 gcgaatctggaggccttcatttgggtgccaatctaaccattaaattgtga 24200  
 cgtttttcttttagGAAGCTCTGTAGATGTGCTATACACTTTTGCAAACT 24250  
 R S S V D V L Y T F A N  
 GCTCAGGACTGGACTTGATCTTTGGCCTAAATGCGTTATTAAAGAACAGCA 24300  
 C S G L D L I F G L N A L L R T A  
 GATTTCAGTGGAAACAGTTCTAATGCTCAGTTGCTCCTGGACTACTGCTC 24350  
 D L Q W N S S N A Q L L L D Y C S  
 TTCCAAGGGGTATAACATTTCTTGGGAACTAGGCAATGgtgagtacccca 24400  
 S K G Y N I S W E L G N  
 gggaacaatttcattataaaggagattccccactagcattattttctttct 24450  
 tttctttttctttttcttttttttttttttttttgagacagagtctcgactgc 24500  
 tggccaggctggagtgcagtggcgccacctcggtcacttgaagctctgc 24550  
 ctcccaaaacgccattctcctgcctcagcctcccgagtagctgggactac 24600  
 aggcaccccgccacccgcgcggcctaatttttttttttttttttttttttt 24650  
 tttttttgcattttttagtagagacgggtttccacgtgttagccaggatg 24700  
 gtcttgatctcctgacctcgatctgacctcctcgccctcccaagtgc 24750  
 tgggattacaggcgtgagccaccaggcccggtagcattattttcttatga 24800  
 cacttt 24850  
 agtgcagtggcgccatctcggtcactgcaagctccacctccaggttca 24900  
 cgccattctcctgcctcagcctcccgagtagctgggactacacgcacccg 24950  
 ccaccacgcccggctaatttttttttttttttttttttttttttttttt 25000  
 ccgtgttagccaggatgggtctctatatcctgaccccatgatctgcccgc 25050  
 tcggcctcccaagtgggtgggattacaggcgtgagccactgcgcggccg 25100  
 aacactctttttattattagcaaatatacttctgctgggcacattcttg 25150  
 caagtgtcaacaatgcaacttttgaagtgcattgtggcagaaactcctg 25200  
 ctgtatttattccagaaactattattgttaattccagttttattgttacatt 25250  
 tgaagtgagaaccagttggagccagcaacgttcccagctccaaagtccct 25300  
 ttgagattttcagaatcacttaacctattatgcttggcaacctggactc 25350  
 agcaaaactgggaagtgcagcagtttgtttttatccatcccttctttctca 25400  
 gtttctcaaatgtgtcagtttaattctcagtaaccccatgcaaccttcatt 25450  
 acctgcccgaagcgtctagaacttgccagtatagaatcctacgtgggtca 25500  
 agctcctgactgtctccttcttcaactcttttttttgcaagaacttgtaaa 25550  
 ttttaactataagttattcatgattcgccacatttattcaaaacatagagt 25600  
 gctttttccacatatcagccaatggaaataaggattaaatgggaaatgaa 25650  
 atgtagtaataaggataaagcacaagtcttcttctgctcaaaacttttttt 25700  
 ttttttttttcagacaagatcttgcctctgttaccaggtctggagtgcagt 25750  
 ggctgtttcatagctcaatgtaacctccaactcctgggctcatgcaatct 25800  
 ctcacacctcagccccctgattagctaggactacactatgcctagccaat 25850  
 tttttttcttttgcctgtgtgttgccaggtgtctcgtatctcctggc 25900  
 ctcaagtaatcctcctgcctcgcccttctaaagtgtctgggattataggca 25950  
 tgagccactgtgcccgtctcaaacctttttttccaaagtaaatgaagt 26000  
 attagatatggaatatagtctagtcccagatatccatatccattggttt 26050  
 attaccctcatttataacttcaaatgtttaatagaccctcatatctcag 26100  
 ttatacagtttaaaattttgtttttgtttttctggagtattttattataa 26150  
 ctatgagttttactttactttatttttttttttttttttttttttttttt 26200  
 ctctgtcactcaggctggagtgcgggttgcgtgatcatggctcactatggc 26250  
 ctgcaccttctgggtcaagtgcctcctcctcagcctcccaagctgag 26300  
 actacaggcatgcaccaccacatctagctaatttttttttttttttttttt 26350  
 aacaaggctttactatgttaccagagtggtctcaaaactcctggcctcag 26400  
 gggatcctcctgtctcagcctaccaaaatgctgggattacaggcatgagc 26450

Fig. 16 (continued)

25/34

catagcgccagacctggttttacttttcttgacttttgaattacaagtttt 26500  
tgtaatttggaaaatgtttgtgtcttttaataactgctgtatgtttgct 26550  
tttaataacaacatttctcgatatataatttggagaattgctgtctttcag 26600  
AACCTAACAGTTCCTTAAGAAGGCTGATATTTTCATCAATGGGTCGCAG 26650  
E P N S F L K K A D I F I N G S Q  
TTAGGAGAAGATTTTATTCATTCATAAACTTCTAAGAAAGTCCACCTT 26700  
L G E D F I Q L H K L L R K S T F  
CAAAAATGCCAAACTCTATGGTCCTGATGTTGGTCAGCCTCGAAGAAAGA 26750  
K N A K L Y G P D V G Q P R R K  
CGGCTAAGATGCTGAAGAGgttaggaactagaggatgcagaatcactttac 26800  
T A K M L K S  
ttttcttcttttcttcttttggagacagagtctcactctgtcagccagactg 26850  
gagtgcagtggtacaatcatggctcactgcaacttcgacctcccaggctc 26900  
aagcaatcctcccatctcagtcaccacaaatagctgggactacaggtgcac 26950  
atcaccacacctggctacttttaaaaaaattttttttagagatggggctc 27000  
cctgtgtgtgcccaggctgtctcttgaaattcctgtgctcaagccatcct 27050  
tccacctcagcctcccagagtgcaggattacaggcatgagccaccacac 27100  
ccagccaccacttttcttaaaaaaaagattctctctggttagacaa 27150  
tcttcaatagtccacatggttattaaacaatctgctgcctgaatacatgat 27200  
ttacaaaaaaaaggaaatttggacgggttcagaatatcaagggatctgag 27250  
gcaaatgtcacctatgataaaatttgctatcaaaattaggaagtgtgtgt 27300  
ttacctgatcctaagcagtaaccagcccatttctagggataaaaactct 27350  
catgcgtatatattgtgcatatatatgtattatatgactgagtgataataa 27400  
attttttttctagCTTCCTGAAGGCTGGTGGAGAAGTGATTGATTCAAGTT 27450  
F L K A G G E V I D S V  
ACATGGCATCAgtaagtatgtctcctattcttaataactaggaaagtaagg 27500  
T W H H  
ctagctttattttattacctagtagtattcaaaaagttagttcatttaactgcc 27550  
aattgactgcagttcaataaagaaacaaatagtgctcaagtagcactgt 27600  
actccaatttttaattttaataaaaaaaattttaagttatttttaataatg 27650  
tagtggttttctataaagatcactttatacagaagaacagtgccaattaac 27700  
ccatggaacatataagtagctaaaaaccaattgcttgccaaagaaccagta 27750  
acccaggagtagatgtccttgccactgtgttttttcaagacagagtaact 27800  
gatttctagttacttgcatagaatggactcctcctcataactcccttcca 27850  
tcttggtctttccctagtagaacttctaccttttttagtaacagggtgag 27900  
tgggagaggtaagaaggagaataaaggctcagcaattaacctaaaagcagaa 27950  
agtaaaatttgttatttttttctgaatatcttctgtgaatttagCTAC 28000  
Y  
TATTTGAATGGACGGACTGCTACACAGGGAAGATTTTCTAAACCTGATGT 28050  
Y L N G R T A T R E D F I N P D V  
ATTGGACATTTTATTTCATCTGTGCAAAAAGTTTCCAGgtaatagtct 28100  
L D I F I S S V Q K V F Q  
ttttaaaactttttaatgtaaaaccagaatccttattttatagttctagcta 28150  
gttctaaattctataggtatgtatatttacatgtttttctaatttttagag 28200  
aacaagcactatgacttatccactgttagttttcccttagcattgggtc 28250  
ttaccccatgtacgtgattagaaatttgaaatatttccaatagcctttag 28300  
tagaattaactcacatagatgataagaatgggttggttacttcatgttc 28350  
cttccacagcctactatttcaataaaagaaagtttccaagacctaataatg 28400  
actatgaacatattttataactatataggaggggtgggtctaggaatata 28450  
aagttttgaatgctgttaattcttcaacaccacagttgaaaccacaggtca 28500  
gcttttttgcaattaccatggatacttttctgttctatagGTGGTTGAGA 28550  
V V E  
GCACCAGGCTGGCAAGAAGGCTGGTTAGGAGAAACAAGCTCTGCATAT 28600  
S T R P G K K V W L G E T S S A Y  
GGAGGCGGAGCGCCCTTGCTATCCGACACCTTTGCAGCTGGCTTTATgtg 28650  
G G G A P L L S D T F A A G F M  
agtgaagcagcgctggccttaggggtcagagtgcagctcttctccatcct 28700  
tctattctgctgaaatagctccccagccaaaagcagatcaaagaccgtt 28750  
tcagtggtgagccccaaaattcatgccagattttgcaagaaaatgattt 28800  
actaaagcttgaggagactctttaacaagtggtccaaattaatcactata 28850  
aggatgaattgtttcagaaattttggcctttaattatggcccataaataat 28900

26/34

Fig. 16 (continued)

gtcaagtagtccttactctaaagaagtacactgtaaaagaatgcatatag 28950  
 ccggatatggtagttccctgtaatcccaatactttgggaggccaaggtgg 29000  
 gaggattgcttgagcccaggagtttgaggctgcagtgagttatgatgggtg 29050  
 ccactgcactctagactgggcaacagagtgagactgtctttttttttccc 29100  
 ctctgtcaccagactggagggcagtgccacgatctcacctcactgcaac 29150  
 ctctgcctcccggttgaagcgattctcctgcctcagcgtcctgagtagc 29200  
 tgggactacaggagtatcccgcaactgggctaatttttgatttttagta 29250  
 gagacggggttttgacatgttgcccaggctgggtctgaaacccatgagctc 29300  
 aagtgtatctgcctacctcagccttccaaaatgctgggattacggacatga 29350  
 gctaccacgcccggccacacccctgtctcttaaaaaaaatgcaag 29400  
 ttagagcatattacagctttgtctctcaggaggatacttagtgtatgtag 29450  
 ctataattcatagattcccaagaagtttagagcctaaagtatgagggtccc 29500  
 accagaggggtctatcattaaatttaaagatttggttaaatcatctcattgt 29550  
 ccaacaccacaacttgattgctttaaataactgggttagttacatttag 29600  
 taactctattagtgcttttaattctatactgctatactcactcattgagat 29650  
 tttttttctttctcttccatcttcattctttttctctcctcctcattc 29700  
 ttataagcctagaatacatcacaaatcctttatgcccattggaagcaagag 29750  
 gaataaagaatggagatgtttgttttgccattaaactaaagatctggggtg 29800  
 tcgggggagaaggggatagagaaggagaagtgggaagaggtgtccataat 29850  
 agcttaggtgcaattctgcttattttacattttacccccgctgactgcca 29900  
 cttttcttcagccctcacacattgtttgtgcagggacctcataggacca 29950  
 ggaattgtctatagaggtgggaatttgtctcaccctgaaagggatactc 30000  
 tagcatggtataatagctctctaggtttgttatcatatggaaagatgtaaa 30050  
 gggagggattctgctgctgctgctgctgctgctgcatgcagttgccatttcat 3 100  
 ttaaatgactttatttataattgatgacacttttctggcttctctgttaatt 30150  
 cctccctcaaagatcaataaaaccagaaccaggcatgggtggcatgcacttg 30200  
 tggctcctgtaaccaccaacaggttcaccttgctgctgcttagatagag 30250  
 ccaattatcaagacaggggaattgcaaaggagaaagagtaatttatgcag 30300  
 agccagctgtgcaggagaccagagttttattattactcaaatcagttctcc 30350  
 ccgaacattcgaggatcagagcttttaaggataatttggccggtaggggc 30400  
 ttaggaaagtggaagtgctggttggtcagggtggagatggaaacacaggg 30450  
 agtggaaagtggaggttttcttgctgctctctggttctggatgggag 30500  
 aactggttggggccagattaccgggtctgggtggtctcaaatgatccacca 30550  
 gttcagggtctgcaagatatctcaagcactgatcttaggttttacaacag 30600  
 tgatgttatccccaggaacaatttggggaggttcagactcttgagccag 30650  
 aggtctgattatccctaaccgtaattctctaattgttgtagctaatttgtt 30700  
 agtctgtcaaaaggttagctttgtccccaggcaagaaggggtcttttcaga 30750  
 aaagggctattatcatttttgtttcagagtcacaacatgaactgaatttc 30800  
 ttcccaaagttagttcagcctacaccaggaatgaagaaggacagcttaa 30850  
 aggttagaagcaagatggagtcattgaggtctgatctctttcactgtcat 30900  
 aatttctcagttataatttttgcaaaggcggtttcagtcacagctactt 30950  
 gggaggctgagacaggaggattaatggagccaggagtttgaggttgag 31000  
 agagctatgatcacgccaactgcactccagcctgggtgacagagtgaacc 31050  
 ctgtctctaaataaaataaataagtaataaaataacataaaataaaatc 31100  
 aagatggtgtgcaattagaattgagcgattttgtttccaaacctcaagaa 31150  
 agcttggtcttgctctgtccagGTGGCTGGATAAATTGGGCCTGTCAGC 31200  
 W L D K L G L S A  
 CCGAATGGGAATAGAAAGTGGTGATGAGGCAAGTATTCTTTGGAGCAGGAA 31250  
 R M G I E V V M R Q V F F G A G  
 ACTACCATTTAGTGGATGAAAACCTTCGATCCTTTACCTgtaagtgaacct 31300  
 N Y H L V D E N F D P L P  
 tattttcttaattctagtgaggtagattaaagtcaactcaggacctctgg 31350  
 tgtaacctcctatgaacagtcagtcctctcagtaactagccaaatcatg 31400  
 agatgatgaattagaaggagccttagatagcatccaatctaactttttt 31450  
 tgtgtgtttgaagagaagaaatcaagagctaggaataaactttttaagggt 31500  
 aagccatttgcagtatagtgtggattttgtttaaaaggggataatttgaa 31550  
 attttatgactcattatacaagacaaaataagttggattttcaaatgttt 31600  
 tacaaagtaaatcaagttataattgcctacagtcacgcaagcttcaaaa 31650  
 cattttttatgttatgaaattgtaattttttaaccttaaaatgagccag 31700  
 taccatgtgtttgtcttaaaatctcatgctaagaatttactatgttgta 31750  
 ataacttcaagatattttagaataaagctctatttctaatccttctctcc 31800

Fig. 16 (continued)

27/34

aactgtatctggtgctaaatcaggaaatgtttcttcccaaaaagcctcgt 31850  
ggaagatctgtatgtctctaaatataatgtcagggataatacagatgtagccc 31900  
tgcaagcatgaccttgatttttatagtctaaatgtcatttgcagatat 31950  
ctattttctaaagaataatctctaaagaattatttgatgtgttaggaaa 32000  
gctaagaaattttgcaagagcgtacgtgaaaatataagctaggcttttg 32050  
tgggttggatagacttcccaacaaaattgctttttatctatagtgtatc 32100  
caagcttgtggaacatattagtcacatcttttttagaaaattcttagaaaa 32150  
gtgatcttgcaaaaatggaatttatctttcccaagtataattctgtcatg 32200  
tatagagtttaactaagcatagtaatttcaccagacaaacattcaaaatc 32250  
tactctgacctttttatctcatccaaattttccaggggccagacataa 32300  
acctttgccttacgaactctttgtatatgcactaaatagtcttctccttc 32350  
aaggttctcagtcagctagaaaaatgtgcaagagtaaatggtacccttct 32400  
caactgttagatcccaagagaattagacttaaaactcactctacatgtctgtg 32450  
actttatttttatgtcatgacagtcctgtgaggtggcaaggcaggtatct 32500  
tggatccattttttatagataaggaagttcaaattgagaagaggttgcatga 32550  
tttacaggaagccatactgtagtctctatgttactcttaaaaaatcccatc 32600  
aaatcctgtctctgagcctgcataactttctaccctaccagtcattgacc 32650  
catgcttatgtctcctttgaaaaacattgattccactcttgtctccagtgga 32700  
aaaagtggaaattttaagcagagaaacaaaagccatttgtcttgtaagtct 32750  
actttccctctactttcaagaaggaaagttggggatgtgttgtaagtgtg 32800  
atattttatattttatatttttaaaaattgatacaaggctcttactgtga 32850  
ttgtgcaggctgtgtctcaaaactcctgggctcaagtgtatcatccacctca 32900  
gcctcccaagtggtgggattacagcatgaaccattgtgcccaccaccgatc 32950  
cgagtttttttaagaaaaacttttactatagaaaattttaatcatataca 33000  
aaatacagaggaaggtatatgaaccacatttaggagactagaatatgccca 33050  
ccccaaaatagccactttggcataaggattatttcgagctaaaggcaac 33100  
tgggaagaaacacatagaagaaaagtctctgtccttctccatttgccta 33150  
aaagcaggacatgaattcttaaaagtccccctccttccctttctaccagga 33200  
aaaaacaagagttaatcactgaagataacttcagacccttatcagtgtaga 33250  
gatggcactagaagaatctatattacatactcatttatttctctccac 33300  
aacttgccaccaccagagactaaaaatccttttcttctgtcatgtctcttg 33350  
tccaaaaatttgcctctataagctggagttctaagccacctcttgagaat 33400  
tacttgctccctgggtattttctgttaacatacatgtattaatatacatgt 33450  
taacaagcttctgtttgtttttctcctgttttctgtctgttacagaggt 33500  
ccatcccaactaagaaactaaagagtaggaggaataataatttctcctctg 33550  
catactttgatcttgtttaatccgtaacccttcccaacttttcacctccta 33600  
cctattagattactttgaagcaaaatttcagatatattactttatctataa 33650  
atatctcagtatgtgctaggtgtggtggctcacacctgtaatcccaacac 33700  
tttgggaagctgaggcaggaggtacttgagccagaggttcaagacca 33750  
gctacggcaacaaaaatcaaaaacttatctgggcatggtggcacatgcc 33800  
tgtggtcccagctacatgagaggtgagggcaggaggtcgttttagccca 33850  
ggaggttgaggctgcagtaagctgcattcacaccactgcactccagcctg 33900  
ggtgacagagtaagaccatgtctcaaaaaatacatattttagttatgtat 33950  
cctttttgtaaaaacacaatactttttatcatactttaataataacaata 34000  
attccttagtatcaccaaataattttgtcagtgctcacattttccttatt 34050  
gtctaaaaatattgttgatagttattcaaatcagaatccaaacaaggtcca 34100  
tatattacatttgggttgacaagtctcttaagtttgttcacctttaagttc 34150  
ttcctccctctctttcatctcttgtaatttattaatgtgaaaaaacaggt 34200  
aatttgtctatagtattttctacattatagagtttgcctacatttatcc 34250  
ctatgatcatatttagcatgttctctgtccctgtgtttctctgtaaaact 34300  
ggtagttatacctagaagcttgagtttattcagggtttttaattgtatttt 34350  
ttttgcaagaattctttattatctgcttctggaagcacagaatgtctgggt 34400  
tgtgtctgggttttgatcttgacagctactgatgaccattgcctaataccat 34450  
tactttattgggtggggggaataaggttttaaaataaattttttttaa 34500  
gatttttttaactgttattttgagacagtgctcatttcgtttccaggc 34550  
tggagtgagtgccacaaatcacggctcactgcagccttgacctcctggga 34600  
tcaggtgatcttctcacctcagcctcctgggtacctggaactacaggtgc 34650  
acaccaccacacctggctaattttttgtattttgtgtacagaaggggttt 34700  
catcatgtttccagactgggtctgaactcctgggttaagtgatctacc 34750  
cacttcagcttcccaaaatcctgggattacactttggccaccgtgctgg 34800  
cctaaatgaaattatttgtctctaaacagacagaagttttactttaaaaa 34850

Fig. 16 (continued)

28/34

tttgtctttgtgtgtacatgtgtttgtgtatgtgtgtgtgtctaaaagtt 34900  
 tggcttttgagcttttgcctttgaattcttggatgaacaataaccaagaatac 34950  
 ttaaactctgtatcattcttgacagatatccctacaggctatggcctttt 35000  
 gaattgtgtcctccagtgataaaaagcagcaagcagcagatactgctctcag 35050  
 attcatgtgtgttcacatgtgaggtgaaaaaaaaaaaaagatgaatccta 35100  
 tttaaatgccccaggataaacagtgatactctttgtaggataactattttg 35150  
 cttgccactgggtttcattaaataaggacataagtaaaagatctatttttgt 35200  
 ctctttctccccaaccaccacaactagGATTATTGGCTATCTCTTCTGTT 35250  
 D Y W L S L L F  
 CAAGAAATTGGTGGGCACCAAGGTGTTAATGGCAAGCGTGCAAGGTTCAA 35300  
 K K L V G T K V L M A S V Q G S  
 AGAGAAGGAAGCTTCGAGTATACCTTCATTGCACAAACACTGACAAGtaa 35350  
 K . R . R . K . L . R . V . Y . L . H . C . T . N . . . T . D . N  
 gtatgaaacacacccctttaccaatcatcaagttttagtgggtaagcctgt 35400  
 aactttactcaaacacccctgttgcattgtgtctatacattgcataagtata 35450  
 ggcagttgcaatttagtaaaagttttatacaacgatttttattttattttat 35500  
 ttttagaagaaaaatgctactttttgtgtgtgtgttttttgagacggggc 35550  
 ctgctcgctcaccaggtggagtgagtgagtgcaatctcagctcactgc 35600  
 aacctccgcctcccggttcaagtgattcttgaagaggagaacaataata 35650  
 acaacaatatttttcaaaagttgtgacccgagtttctggagttgagaa 35700  
 gacatcgagattttgtagcctcatactcttgccttaggtagcaaaaaat 35750  
 gttcttaaatctcaggaatattctctagataggtttcaatctatcattcc 35800  
 tgataagatgatgtgaaataactaattctagccaaaaaagaccagctacc 35850  
 atttccgattgttggggactgggaactctggatagttaggacccagtag 35900  
 gaagtgcgaggggaatggtttgaatggataaattcataaaaaatgtcag 35950  
 tagatttaattttcttatacatttcagtcctttttataaggctaggaaaaag 36000  
 cccctgtttttatggtttataaatttgaattcacatgaaccacaaaattt 36050  
 gccttttaccttctctatgtctgaaaatggatagtctggtggtcctcttaa 36100  
 caaccagctggcagagctgtgaggatctcagtggtgctctagcccagaca 36150  
 ttggtagcatgaacggcaacatttttaattgtgttttcaaaataggagca 36200  
 cactagcgggtctaaaacgatcataaaagaaggataactaagaggggccact 36250  
 gtcatatttgatccttaacttaggatgcattatggattgtcattatgga 36300  
 tactaatacttaggatcacatttgaattgagtttttaattgcttaaat 36350  
 agatacatatttctattaagttaacctcttgccttttagTCCAAGGTATA 36400  
 P R Y  
 AAGAAGGAGATTAACTCTGTATGCCATAAACCTCCATAATGTCACCAAG 36450  
 K . E . G . D . L . T . L . Y . A . I . N . L . . . H . N . . V . T . K  
 TACTTGCGGTACCCCTATCCTTTTCTAACAAGCAAGTGGATAAATACCT 36500  
 Y L R L P Y P F S N K Q V D K Y L  
 TCTAAGACCTTTGGGACCTCATGGATTACTTTCCAAGtaagtaattttcc 36550  
 L . R . E . L . G . . P H G L L S K  
 ttgttcatccaaactttcaataaattttattgggtgtttatcagaatagag 36600  
 agtttggacagggagcaaaagcaaaagtcactatatcaagttctaataa 36650  
 ttcttaatatcaggaaattttatgtatgaatacttactaataatgagtata 36700  
 actcatcctaagagtctaaagcaaaaggatgtgaacacaaactagcagtt 36750  
 atcttagagaataagtttgcatttcaaaaataactgacataatcaagatcc 36800  
 actcaacgcattttaaattatttactctaaaaagacataaattcttggtaac 36850  
 acattcactaaagcaaaatatacctttatataattgctatcaaagggtatg 36900  
 tgggttggatataaaatatcataccatgtgagatcagtggtgattcctttac 36950  
 agcataaatttttattggttagagtaagaaaaagaatagctagagtata 37000  
 ttcttaagtagattctcatataccttgggttcaaaaaccaattattgact 37050  
 acatcttataaaagcctgtattcaatggagtgccaaaaaatgactatgag 37100  
 tcttaaaagagttaggcatataaatttttaaggttctgttcaatgtatg 37150  
 ttggaaggagtttcttctcatgactattctcatatttgagcataaaaaag 37200  
 agtttacaggcttggcgcagtggtcatgcctgtaatcccaataactttgg 37250  
 gaagctgaagcaggcagatcacttcagcccaggagtttgagaccagcctg 37300  
 ggcaatatggcaaaactctctctacaaaatatacaaaaattagccaggcg 37350  
 tgggtgtgcatgctgtagtcccgactacttgggaagctgaggtgggagg 37400  
 attgcttgagcccaggggggtcatggctgcagtgagctgtgatggtgcct 37450  
 ctgtcaccagcctgggtgacagagtgcagccctgtctcaaaaaataaaa 37500  
 taaataaaaaatgaagagtttcaaaaattctcacatctcctcccatctt 37550

Fig. 16 (continued)

29/34

gcaaatgccacataagtgatgtgttccaggactattagcctcggaaacctg 37600  
aggcagtacagtaagcacgctttctccaaagtcctgtcccccacagacaa 37650  
acattatttacactgggtactgctcttttattttttccctctatgcttt 37700  
attttactataactataatcatataacatgtaataaggaaaaaggcagggt 37750  
cgggggagagatccagaagtcttcccaagagcctttccaacatagcctct 37800  
gtagacatttttctttctctttttttttttttttttttttctgagaca 37850  
gagtctcactctgttgcctcaggctagagtgcagtggcgtgatctaggctc 37900  
actgcaacctccgacctcctgggttcaagcaattctccacctcagcctcc 37950  
ctagtagctgggattagaggcatgcacaccagcctggctaatttttgt 38000  
attttttagtagagatgagggttcaccatgtgggccaggctggtcttgaac 38050  
tccctgacctcaagtgtaccacctgccttagcctcccaaagtgttaggatt 38100  
acacgagttagcaccgtgccttgccttaccattctgatcacacatt 38150  
tcatgttttataattggaaaaactggtgaaattatagacaatgttttgttc 38200  
ccctaaattctctttgatgagtataattacttacactctctgtcttta 38250  
aaattttgcaaaatagtatccctagataagtttatgagtgcacagtctgta 38300  
cgcttactcatattaaatgacctcggagagttaacaacagtcacctttaa 38350  
aaattattactatcatatcatattttttagggcgggggtctcattctgt 38400  
ctcccaggctggagagtagtgggtgcagctcactgcagccaccgc 38450  
tacctgggctcaagtgtccttccctcctcagcctctctgtagctgagac 38500  
cacaggcttatgtaccacacctggctaattttttaactttttgtagaga 38550  
cgatgtctcattatgttggccaggctggtctcaaaactcctaagctcaagt 38600  
gatcttctcagcctcccaaagtgtggtgattacaggcatgaaaaactgc 38650  
accagccctaaaaattatttagggctcctgcatagtaagactttaataaat 38700  
atttaaatgaacatctgggttttttaaaaaaaaaatagagacaaggtctc 38750  
actatattgcccagctggtctcgaactcctggactcagcgaatcctgct 38800  
gccttagcgcgccaaagtgtgggtattacaggcatgaccacacctcatctg 38850  
ggctgagtgaacatatttttaacataaaggccgtattttatatttatctc 38900  
atacattttgcccagcatcccatattccgcgaatctgttgccttgctaatt 38950  
tccttccagcttcatttcattetgaaatttgacaaacatcttctatttctt 39000  
tgtcgtcatgtttatttgacttcagaatataaaaataaaacactatacccaa 39050  
ttaaacccaccctcattgcccagcctgatgtgaaaaataatcagcataca 39100  
ttaagcttaccttgatataatgtgtagcatcttttagataaatatacagc 39150  
tgattaaagcaatatagcctgatgggtataatatcttgcccatgtacctcat 39200  
cttatctccagcaggattaaattcacagtgatcagatttacctttaacttt 39250  
tgttagcaaaatatcctctccaaaagcatatctaaaacttttgtgtgtact 39300  
cttgcaagtttcttaatttcattgcagaacaggtcttaccactgttagct 39350  
ggagatattttcaagacctatttttgtttgtgggttctctgatgtggtca 39400  
tggcattttcccttccatctctaaaaattgaggtgatacaggctttt 39450  
aaacaaaaccaaactcatatagactgagtacaactgcaatgcaggcatgct 39500  
aacctctgtacaatcatgggcgtgctattgatgtcttaagttacaga 39550  
acacgggctgagcgtctcattagggtcaaaatgtaaacagtttttctgc 39600  
tactgatgcttaatgaggacagggtgtgagagatttctttaaggaaaac 39650  
aaatatataaatgtcatatggaataatcttaacattagagaattaaag 39700  
taataaaactaatatactcacaccatggaatcttgtgcagacattaaaat 39750  
tatgtagtggatggatgttttaattgggtgtgagaaaaagttaggatgtgctg 39800  
gggtggggggaagaatcaagttttaagaaaaatagataacccatactta 39850  
agtaaaaaaaaaaaaaaaggatgtacagtcagtggtgttaatgatgg 39900  
ggatacattccgagaaaatgtgtcgatagtgatttcatccttgtgtgaac 39950  
atcatagagtgaacttacacaaacctagatgggtctagcctactatgtatc 40000  
taggttatatgactagcctgttgcctcctaggctacaaacctgtaaagcat 40050  
gttactgtagcgaatatacaataacttaacacaaatggcaagctatcattg 40100  
tgttaagtgtgtgtatctaaacatatctaaaacatagaaaaactaatgt 40150  
gttgtgtacaaatgttaacatgactatgacattgctaggcaataggaatt 40200  
ataattttatccttttatggaaccacacttatatagcgtccatgggtgg 40250  
accaaaacatccttatgtggcatatgactgtatacatgtacacaaaaat 40300  
agatgaaagaaatgaatataacataaaatatttaaatgggtataatgact 40350  
taggttacttttatttatcttagtaataataatgatgatagataatactt 40400  
ttatagtgtttactatataaaaagacactgttataagtggtctacatactt 40450  
tacctgtattacctaagtataaaatataaactctgacagtaactaatct 40500  
tatacgttctcttttcttttttttttttttttttttttagacagaatctt 40550  
gctctaccaggctggagtgcagggtgcaatctcggctcactgcacacctcc 40600



30/34

Fig. 16 (continued)

gcctcccagggttcaaacgattctcatgtctcagcctcctgagtagctggg 40650  
 actacaggcacacaccaccatgcccggctaatttttgtatttttgggtag 40700  
 agatggaggttttgccatgttggccaggctgatcttgaactcctggcctca 40750  
 agtgcctgcctgcctcagcctcccaagtgctgggattacaggtgtgaa 40800  
 ccactgtgctcggcctaattcttacaagttttcaatatttaaagagtgtca 40850  
 actttgttgacaatatataaacatatttgagaaaaagagatatagcatct 40900  
 tatttagaattatgaaaaatcaatagacctacagccgactaaagctttt 40950  
 cttcataagctcttgccctatttgattcgctcctgtgaatatgcattaat 41000  
 ttgatttaaataataagtagtataaagaaataacacttttcttaatttt 41050  
 taagaacgttoaacagtttttaatttgaaattccaatagtgaatacatag 41100  
 aaaaataaaaaattttctgtagtttagccaaattgtttttgtttcaccaca 41150  
 gcattctacccaaaattttcttaataacagtaagaaaaatgaatgcatacctc 41200  
 ctgcaggggagaggggagtaggtaggtttatgggcatagttacaagtgtgaga 41250  
 aatttcattggctaccattttacgctaaattcataaaaaactgcattcaatt 41300  
 ctatatatctattttctttacataaaaaagggtttcaattatttgccatta 41350  
 aataaaatagccaccattccagaagttgtgtcatgtttatcctttttata 41400  
 ccaccatcatattgcctattatagattgtgtgtgttccattttctgtat 41450  
 atgggcccagacagtaagttttctggtttggagtcctatgtgtctctat 41500  
 cataactactcatctctgcccattgttagcttaaagattatctaggccaat 41550  
 gcctaagtgatatagtggtgaaatacaagttatataataggtgccac 41600  
 aaaaaaaaattttatttgggtctaaaaaagatttcatgactttttagcagc 41650  
 atgggtggggcatgcaccacttggttaactcgggtgtatctttctccttg 41700  
 cagATCTGTCCAACCTCAATGGTCTAACTCTAAAGATGGTGGATGATCAAA 41750  
 S V Q L N G L T L K M V D D Q  
 CCTTGCCACCTTTAATGGAAAAACCTCTCCGGCCAGGAAGTTCAGTGGGC 41800  
 T L P P L M E K P L R P G S S L G  
 TTGCCAGCTTTCTCATATAGTTTTTTTGTGATAAGAAATGCCAAAGTTGC 41850  
 L P A F S Y S F F V I R N A K V A  
 TGCCTGCATCTGAAAAATAAAATATACTAGTCCTGACACTGaatttttcaa 41900  
 A C I \*  
 gtataactaagagtaaaagcaactcaagttataggaaggaagcagatacct 41950  
 tgcaagcaactagtggtgtcttgagagacactgggacactgtcagtgct 42000  
 agatttagcacagtattttgatctcgctaggtagaacactgctaataata 42050  
 atagctaataataacctgtttccaaatactgcttagcattttgcatgtttt 42100  
 acttttatctaaagttttgtttgttttattattttttttttttttttt 42150  
 ttgagacagaatctctctctgtcaccaggttgagtgccatgggtgcgat 42200  
 ctggctcactgcacattttaagcaattctcctgcctcagcttctctgagta 42250  
 gctgggattatagggctgtgccaccacgcccagctactttctatatatttt 42300  
 ttagagagatggagtttcgccaatttgcccaagctgggtctcgaactcctgt 42350  
 cctcgaactcctgtcctcaagtgatccaccgcctcagcctctcaaagtgt 42400  
 ctgggattacaggtgtgagccaccacacccagcagtggtttatttttgag 42450  
 acaggggtatcattctgttgcccaggttgagtgagtggtgtgcaatcatag 42500  
 atcactgcagccttttaactcctgggtcaagtcacctcctgcttagcc 42550  
 tcccaagtagctaggaccacagacacatgccatcacacttggtattttt 42600  
 aaaaaattttttgtagagatgggtgtcgtatgttaccctaaactgggtcc 42650  
 tgaactcctggactcaattgatcctccaccttggccttccaggtgtgtg 42700  
 gatttctttgggagtagacagcatggttacagcaggagatcatttgatgttac 42750  
 ctctgtgcagtggtgctagtcagcgaaagactataatacctgtggggaca 42800  
 gcgattagccaccacaaccagtccttatttaaagttattaaaaatggctg 42850  
 ggcgcagtggtcacacctgtaatcctagcactttgggagggcaggcag 42900  
 atggatcacctgacgtgaggaatttgagaccagcctggccaacatgggtga 42950  
 aaccccatctctactaaaaaatacaaaaaattagctgggtgtgtgtcctgta 43000  
 gtcccagctacttgggaggtcggggcaggagaattacttgaaccaggag 43050  
 gcagaggttgagtgagcggagattgtgccactgcactccagcctgggtg 43100  
 acagagagagattccatctcaaaaaaacaagttattaaaaatgtatatga 43150  
 atgctcctaataatggtcaggaagcaaggaagcgaaggatattatgagt 43200  
 ttaagaaggtgttagctgtatatttatctttcaaaatgtattagaaga 43250  
 ttttagaattctttccttcatgtgccatctctacaggcaccatcagaaa 43300  
 aagcatactgcccgtgtaccgtgaaactgggtgtaaaagagaaactatctat 43350  
 ttgcaccttaaaagacagctagattttgtgtattttcttcttctcggtttt 43400

Fig. 16 (continued)

31/34

```
ctttgtcagcaataatatgtgagaggacagattgtagatatgatagtat 43450
aaaaaatgggttaatgacaattcagagggcaggagattctgtaaaacttaaa 43500
attactataaaatgaaattgatttgtcaagaggataaaatttagaaaacac 43550
ccaataaccttataactgtctgttaatgcttgcttttctctacctttctt 43600
ccttggtttcagttgggaagcttttggctgcaagtaacagaaactcctaatt 43650
tcaaattggcttaagcaataaggaaatgtatattccacataactagacgt 43700
tcaaacaggccagggtccagcacttcagtacgtcaccagggatctgggtt 43750
cttcccagctctctgctctgcatcttttagcgctggcttcattctcagac 43800
tctggtagcatgatggctgtagctgtttcatgggccccttcaaacctcat 43850
agcaaccagaggaagaaaatgagccatttttgagtctccttcatagact 43900
tgaataactctttttcagagcttctcacagcaaacctctcctcatgtctc 43950
ctcatgtcttattgttcagaaaatgggtaatgtggccatttcaccagtcac 44000
tgccaacaacaacgaggttcctataattgtctctgagtaaccctttggaa 44050
tggagaggggtgttggtcagctctacaaactgaacactgcagttctgcgctt 44100
tttaccagtgaaaaaatgtaattattttccctcttaaggattaatatctc 44150
ttcaaattgtatgcctgttatggatatagtatctttaaaattttttatttt 44200
aatagcttttaggggtacacactttttgcttacaggggtgaattgtgtagt 44250
ggtgaagactcggcttttaattgtacttgtcacctgagtgatgtacattgt 44300
acccaataggtaatttttcatccattaccctccttccgccctcttccctt 44350
ctgagctctccaacatcccttataccactgtgtatgttcttgtgtacctac 44400
agctaagcttccacttataagtggagaacatgcagttatttggttttccatt 44450
cctgagttacttcccttaggataacagccccagttccgtccaagttgct 44500
gcaaaatataattattcttctttatggctgagtaatagtccatggtacata 44550
tataccacattttctttatccacttatcagttgatggacacttaggttaa 44600
ttccattcaatttcattcaatttaagtataatttgtaaggagctaaagctg 44650
aaaattaaatttttagatctttcaatactcttaaatttttatatgtaagtgg 44700
tttttatattttcacatttgaaataaagtaatttttataaccttgatatt 44750
gtatgactattcttttagtaatgtaaagcctacagactcctacatttgga 44800
accactagtggtgtgtttcacccttggtatactatcaggatcctcga 44898
```


32/34

Figure 17

					50
human	MLLRSKPALP	PPLMLLLLGP	LGPLSPGALP	RPAQAQDVVD	LDFFTQEPFH
mouse	~~~~~ML	RLLLWLWGP	LGALAQGAPA	GTAPTDDVVD	LEFYTKRPLR
rat	~~~~~	~LLLLWLWGR	LRALTQGTTPA	GTAPTKDVVD	LEFYTKRLFQ
					100
human	LVSPSFLSVT	IDANLATDPR	FLILLGSPKL	RTLARGLSPA	YLRFGGTKTD
mouse	SVSPSFLSIT	IDASLATDPR	FLTFLGSPRL	RALARGLSPA	YLRFGGTKTD
rat	SVSPSFLSIT	IDASLATDPR	FLTFLSSPRL	RALSRLSPA	YLRFGGTKTD
					150
human	FLIFDPKES	TFEERSYWQS	QVNQDICKYG	SIPPDVEEKL	RLEWPIQEQL
mouse	FLIFDPKEP	TSEERSYWKS	QVNHDICRSE	PVSAAVLRKL	QVEWPFQELL
rat	FLIFDPNNEP	TSEERSYWQS	QDNNDICGSD	RVSADVL~~~	~~~~~
					200
human	LLREHYQKKE	KNSTYSRSSV	DVLYTFANCS	GLDLIFGLNA	LLRTADLQWN
mouse	LLREQYQKEF	KNSTYSRSSV	DMLYSFAKCS	GLDLIFGLNA	LLRTPDLRWN
rat	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
					250
human	SSNAQLLLDY	CSSKGYNISW	ELGNEPNSFL	KKADIFINGS	QLGEDYIQLH
mouse	SSNAQLLLDY	CSSKGYNISW	ELGNEPNSFW	KKAHILIDGL	QLGEDFVELH
rat	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
					300
human	KLLRKSTFKN	AKLYGPDVGQ	PRRKTAKMLK	SFLKAGGEVI	DSVTWHHYLL
mouse	KLLQRSFAQN	AKLYGPDIGQ	PRGKTVKLLR	SFLKAGGEVI	DSLTHWHHYLL
rat	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
					350
human	NGRTATREDF	LNPDVLDIFI	SSVQKVFQVV	ESTRPGKKVW	LGETSSAYGG
mouse	NGRIATKEDF	LSSDALDIFI	LSVQKILKVT	KEITPGKKVW	LGETSSAYGG
rat	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
					400
human	GAPLLSDTFA	AGFMWLDKLG	LSARMGIEVV	MRQVFFGAGN	YHLVDENFDP
mouse	GAPLLSNTFA	AGFMWLDKLG	LSAQMGIEVV	MRQVFFGAGN	YHLVDENFEP
rat	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
					450
human	LFDYWLSLLF	KKLVGTVKVM	ASVQGSKRRK	LRVYLHCTNT	DNPRYKEGDL
mouse	LFDYWLSLLF	KKLVGSPRVL	SRVKGPDRSK	LRVYLHCTNV	YHPRYQEGDL
rat	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
					500
human	TLYAINLHNV	TKYLRLPYPF	SNKQVDKYLL	RPLGPHGLLS	KSVQLNGLTL
mouse	TLYVNLHNV	TKHLKVPPPL	FRKPVDTYLL	KPSGPDGLLS	KSVQLNGQIL
rat	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~L
					543
human	KMVDDQTLPP	LMEKPLRPGS	SLGLPAFSYS	FFVIRNAKVA	ACI~
mouse	KMVDEQTLPA	LTEKPLPAGS	ALSILPAFSYG	FFVIRNAKIA	ACI~
rat	KMVDEQTXPA	LTEKPLPAGS	SLSVPAFSYG	FFVIRNAKIA	ACI~

33/34

Figure 18



34/34

Figure 19

```
|MLLRSKPALPPPLMLLLGLPLGPLSPGALPRPAQAQDVVDLDFFTQEPHLVSPSFLSVT| 60
PHD |          EEEEE          HHH          EEEE   EEE|

|IDANLATDPRFLILGSPKLRTLARGLSPAYLRFGGTKTDFLI FDPKKESTFEERSYWQS| 120
PHD |EEE          EEEEE  HHHHHH  HHHHE  EEEEE          HHHHHH|

|QVNQDICKYGSIPPDVEEKLRLWPYQEQLLLREHYQKKFKNSTYSRSSVDVLYTFANCS| 180
PHD |HHHHHHHH  HHHHHH  HHHHHHHHHHHHHHH  EEEEEEEEEEE |

|GLDLIFGLNALLRTADLQWNSSNAQLLLDYCSSKGYNISWELGNPNPNSFLKKADIFINGS| 240
PHD | HHHHHHHHHHHHHHHHHHHHHHHHHHHHHHH  EEEEE  HHHHHH EEEE |

|QLGEDYIQLHKLLRKSTFKNAKLYGPDVGQPRRKTA KMLKSFLKAGGEVIDSVTWHHYL| 300
PHD | HHHHHHHHHHHHHHHHHHHHHHHHHHHHHHH  HHHHHHHHHHHHH  EEEEEEEEEEE |

|NGRTATREDFLNPDVLDIFISSVQKVQVVESTRP GKQVWLGETSSAYGGGAPLLSDTFA| 360
PHD |          HHHHHHHHHHHHEEEEEEE  EEEEE  HHHHHH|

|AGFMWLDKLGLSARMGIEVVMRQVFFGAGNYHLVDENFDPLPDYWLSLLFKKLVGTVKLM| 420
PHD |HHHHHHHH  HHHH HHHHHHHHHHH  EEEEE  HHHHHHHHHHH  EEEEE|

|ASVQGSKRRKLRVYLHCTNTDNPRYKEGDLTLYAINLHNVT KYRLPYPF SNKQVDKYLL| 480
PHD |EEE  E EEEEEEE  EEEEE  EEEEE  HHHHHHHH|

|RPLGPHGLLSKSVQLNGLTLKMVDDQTLPPIMEKPLRPGSSLG LPAFSYSFFVIRNAKVA| 540
PHD |HH  EEEEEEE  EEEEE          EEEEEEE EE |

|ACI|
PHD | |
```

543

## 1

## SEQUENCE LISTING

- (1) GENERAL INFORMATION:
- (i) APPLICANT: Iris Pecker, Israel Vlodavsky and Elena Feinstein
- (ii) TITLE OF INVENTION: POLYNUCLEOTIDE ENCODING A POLYPEPTIDE HAVING HEPARANASE ACTIVITY AND EXPRESSION OF SAME IN GENETICALLY MODIFIED CELLS
- (iii) NUMBER OF SEQUENCES: 47
- (iv) CORRESPONDENCE ADDRESS:
- (A) ADDRESSEE: Mark M. Friedman c/o Anthony Castorina
- (B) STREET: 2001 Jefferson Davis Highway, Suite 207
- (C) CITY: Arlington
- (D) STATE: Virginia
- (E) COUNTRY: United States of America
- (F) ZIP: 22202
- (v) COMPUTER READABLE FORM:
- (A) MEDIUM TYPE: 1.44 megabyte, 3.5" microdisk
- (B) COMPUTER: Twinhead\* Slimnote-890TX
- (C) OPERATING SYSTEM: MS DOS version 6.2, Windows version 3.11
- (D) SOFTWARE: Word for Windows version 2.0 converted to an ASCII file
- (vi) CURRENT APPLICATION DATA:
- (A) APPLICATION NUMBER:
- (B) FILING DATE:
- (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
- (A) APPLICATION NUMBER: 08/922,170
- (B) FILING DATE: 2 SEP 1997
- (A) APPLICATION NUMBER: 09/109,386
- (B) FILING DATE: 10 JUL 1998
- (A) APPLICATION NUMBER: PCT/US98/17954
- (B) FILING DATE: 31 AUG 1998
- (A) APPLICATION NUMBER: 09/258,892
- (B) FILING DATE: 1 MAR 1999
- (viii) ATTORNEY/AGENT INFORMATION:
- (A) NAME: Friedman, Mark M.
- (B) REGISTRATION NUMBER: 33,883
- (C) REFERENCE/DOCKET NUMBER: 910/62
- (ix) TELECOMMUNICATION INFORMATION:
- (A) TELEPHONE: 972-3-5625553
- (B) TELEFAX: 972-3-5625554
- (C) TELEX:
- (2) INFORMATION FOR SEQ ID NO:1:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 27
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:  
CCATCCTAAT ACGACTCACT ATAGGGC 27
- (2) INFORMATION FOR SEQ ID NO:2:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 24
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single

2

- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:  
GTAGTGATGC CATGTAAC TG AATC 24
- (2) INFORMATION FOR SEQ ID NO:3:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 23
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:  
ACTCACTATA GGGCTCGAGC GGC 23
- (2) INFORMATION FOR SEQ ID NO:4:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 22
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:  
GCATCTTAGC CGTCTTTCTT CG 22
- (2) INFORMATION FOR SEQ ID NO:5:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 15
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:  
TTTTTTTTTT TTTT 15
- (2) INFORMATION FOR SEQ ID NO:6:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 23
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:  
TTCGATCCCA AGAAGGAATC AAC 23
- (2) INFORMATION FOR SEQ ID NO:7:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 24
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:  
GTAGTGATGC CATGTAAC TG AATC 24
- (2) INFORMATION FOR SEQ ID NO:8:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 9
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:  
Tyr Gly Pro Asp Val Gly Gln Pro Arg

## (2) INFORMATION FOR SEQ ID NO:9:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1721  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

CTAGAGCTTT CGACTCTCG CTGCGCGGCA GCTGGCGGGG GGAGCAGCCA GGTGAGCCCA 60  
 AGATGCTGCT GCGCTCGAAG CCTGCGCTGC CGCGCGCGCT GATGCTGCTG CTCCTGGGGC 120  
 CGCTGGGTCC CCTCTCCCCT GGCGCCCTGC CCCGACCTGC GCAAGCACAG GACGTCGTGG 180  
 ACCTGGACTT cTTCACCCAG GAGCCGCTGC ACCTGGTGAG CCCCTCGTTC CTGTCCGTCA 240  
 CCATTGACGC CAACCTGGCC ACGGACCCGC GGTTCCTCAT CCTCTGGGT TCTCCAAAGC 300  
 TTCGTACCTT GGCCAGAGGC TTGTCTCCTG CGTACCTGAG GTTTGGTGGC ACCAAGACAG 360  
 ACTTCCTAAT TTTCGATCCC AAGAAGGAAT CAACCTTTGA AGAGAGAAGT TACTGGCAAT 420  
 CTCAAGTCAA CCAGGATATT TGCAAATATG GATCCATCCC TCCTGATGTG GAGGAGAAGT 480  
 TACGGTTGGA ATGGCCCTAC CAGGAGCAAT TGCTACTCCG AGAACACTAC CAGAAAAAGT 540  
 TCAAGAACAG CACCTACTCA AGAAGCTCTG TAGATGTGCT ATACACTTTT GCAAACTGCT 600  
 CAGGACTGGA CTTGATCTTT GGCCTAAATG CGTTATTAAG AACAGCAGAT TTGCAGTGGG 660  
 ACAGTTCTAA TGCTCAGTTG CTCCTGGACT ACTGCTCTTC CAAGGGGTAT AACATTTCTT 720  
 GGGAACTAGG CAATGAACCT AACAGTTTCC TTAAGAAGGC TGATATTTTC ATCAATGGGT 780  
 CGCAGTTAGG AGAAGATTAT ATTCAATTGC ATAACTTCT AAGAAAGTCC ACCTTCAAAA 840  
 ATGCAAAACT CTATGGTCTT GATGTTGGTC AGCCTCGAAG AAAGACGGCT AAGATGCTGA 900  
 AGAGCTTCCT GAAGGCTGGT GGAGAAGTGA TTGATTCACT TACATGGCAT CACTACTATT 960  
 TGAATGGACG GACTGCTACC AGGGAAGATT TTCTAAACCC TGATGTATTG GACATTTTAA 1020  
 TTTCACTCTG GCAAAAAGTT TTCCAGGTGG TTGAGAGCAC CAGGCCTGGC AAGAAAGTCT 1080  
 GGTTAGGAGA AACAACTCTT GCATATGGAG GCGGAGCGCC CTGTCTATCC GACACCTTTG 1140  
 CAGCTGGCTT TATGTGGCTG GATAAATTGG GCCTGTCAGC CCGAATGGGA ATAGAAGTGG 1200  
 TGATGAGGCA AGTATTCTTT GGAGCAGGAA ACTACCATTT AGTGGATGAA AACTTCGATC 1260  
 CTTTACCTGA TTATGGCTA TCTCTTCTGT TCAAGAAATT GGTGGGCACC AAGGTGTTAA 1320  
 TGGCAAGCGT GCAAGGTTCA AAGAGAAGGA AGCTTCGAGT ATACCTTCAT TGCACAAACA 1380  
 CTGACAATCC AAGGTATAAA GAAGGAGATT TAACTCTGTA TGCCATAAAC CTCCATAACG 1440  
 TCACCAAGTA CTTGCGGTTA CCCTATCCTT TTTCTAACAA GCAAGTGGAT AAATACCTTC 1500  
 TAAGACCTTT GGGACCTCAT GGATTACTTT CCAAATCTGT CCAACTCAAT GGTCTAATC 1560  
 TAAAGATGGT GGATGATCAA ACCTTGCCAC CTTTAATGGA AAAACCTCTC CGGCCAGGAA 1620  
 GTTCACTGGG CTTGCCAGCT TTCTCATATA GTTTTTTGT GATAAGAAAT GCCAAAGTTG 1680  
 CTGCTTGCAT CTGAAAAATA AATATACTAG TCCTGACACT G 1721

## (2) INFORMATION FOR SEQ ID NO:10:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 543  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Met Leu Leu Arg Ser Lys Pro Ala Leu Pro Pro Pro Leu Met Leu Leu  
 5 10 15  
 Leu Leu Gly Pro Leu Gly Pro Leu Ser Pro Gly Ala Leu Pro Arg Pro  
 20 25 30  
 Ala Gln Ala Gln Asp Val Val Asp Leu Asp Phe Phe Thr Gln Glu Pro  
 35 40 45  
 Leu His Leu Val Ser Pro Ser Phe Leu Ser Val Thr Ile Asp Ala Asn  
 50 55 60



Leu Ala Thr Asp Pro Arg Phe Leu Ile Leu Leu Gly Ser Pro Lys Leu  
65 70 75 80

Arg Thr Leu Ala Arg Gly Leu Ser Pro Ala Tyr Leu Arg Phe Gly Gly  
85 90 95

Thr Lys Thr Asp Phe Leu Ile Phe Asp Pro Lys Lys Glu Ser Thr Phe  
100 105 110

Glu Glu Arg Ser Tyr Trp Gln Ser Gln Val Asn Gln Asp Ile Cys Lys  
115 120 125

Tyr Gly Ser Ile Pro Pro Asp Val Glu Glu Lys Leu Arg Leu Glu Trp  
130 135 140

Pro Tyr Gln Glu Gln Leu Leu Leu Arg Glu His Tyr Gln Lys Lys Phe  
145 150 155 160

Lys Asn Ser Thr Tyr Ser Arg Ser Ser Val Asp Val Leu Tyr Thr Phe  
165 170 175

Ala Asn Cys Ser Gly Leu Asp Leu Ile Phe Gly Leu Asn Ala Leu Leu  
180 185 190

Arg Thr Ala Asp Leu Gln Trp Asn Ser Ser Asn Ala Gln Leu Leu Leu  
195 200 205

Asp Tyr Cys Ser Ser Lys Gly Tyr Asn Ile Ser Trp Glu Leu Gly Asn  
210 215 220

Glu Pro Asn Ser Phe Leu Lys Lys Ala Asp Ile Phe Ile Asn Gly Ser  
225 230 235 240

Gln Leu Gly Glu Asp Tyr Ile Gln Leu His Lys Leu Leu Arg Lys Ser  
245 250 255

Thr Phe Lys Asn Ala Lys Leu Tyr Gly Pro Asp Val Gly Gln Pro Arg  
260 265 270

Arg Lys Thr Ala Lys Met Leu Lys Ser Phe Leu Lys Ala Gly Gly Glu  
275 280 285

Val Ile Asp Ser Val Thr Trp His His Tyr Tyr Leu Asn Gly Arg Thr  
290 295 300

Ala Thr Arg Glu Asp Phe Leu Asn Pro Asp Val Leu Asp Ile Phe Ile  
305 310 315 320

Ser Ser Val Gln Lys Val Phe Gln Val Val Glu Ser Thr Arg Pro Gly  
325 330 335

Lys Lys Val Trp Leu Gly Glu Thr Ser Ser Ala Tyr Gly Gly Ala  
340 345 350

Pro Leu Leu Ser Asp Thr Phe Ala Ala Gly Phe Met Trp Leu Asp Lys  
355 360 365

Leu Gly Leu Ser Ala Arg Met Gly Ile Glu Val Val Met Arg Gln Val  
 370 375 380  
 Phe Phe Gly Ala Gly Asn Tyr His Leu Val Asp Glu Asn Phe Asp Pro  
 385 390 395 400  
 Leu Pro Asp Tyr Trp Leu Ser Leu Leu Phe Lys Lys Leu Val Gly Thr  
 405 410 415  
 Lys Val Leu Met Ala Ser Val Gln Gly Ser Lys Arg Arg Lys Leu Arg  
 420 425 430  
 Val Tyr Leu His Cys Thr Asn Thr Asp Asn Pro Arg Tyr Lys Glu Gly  
 435 440 445  
 Asp Leu Thr Leu Tyr Ala Ile Asn Leu His Asn Val Thr Lys Tyr Leu  
 450 455 460  
 Arg Leu Pro Tyr Pro Phe Ser Asn Lys Gln Val Asp Lys Tyr Leu Leu  
 465 470 475 480  
 Arg Pro Leu Gly Pro His Gly Leu Leu Ser Lys Ser Val Gln Leu Asn  
 485 490 495  
 Gly Leu Thr Leu Lys Met Val Asp Asp Gln Thr Leu Pro Pro Leu Met  
 500 505 510  
 Glu Lys Pro Leu Arg Pro Gly Ser Ser Leu Gly Leu Pro Ala Phe Ser  
 515 520 525  
 Tyr Ser Phe Phe Val Ile Arg Asn Ala Lys Val Ala Ala Cys Ile  
 530 535 540 543

## (2) INFORMATION FOR SEQ ID NO:11:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1721  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

CT AGA GCT TTC GAC 14  
 TCT CCG CTG CGC GGC AGC TGG CGG GGG GAG CAG CCA GGT GAG CCC AAG 62  
 ATG CTG CTG CGC TCG AAG CCT GCG CTG CCG CCG CCG CTG ATG CTG CTG 110  
 Met Leu Leu Arg Ser Lys Pro Ala Leu Pro Pro Pro Leu Met Leu Leu  
 5 10 15  
 CTC CTG GGG CCG CTG GGT CCC CTC TCC CCT GGC GCC CTG CCC CGA CCT 158  
 Leu Leu Gly Pro Leu Gly Pro Leu Ser Pro Gly Ala Leu Pro Arg Pro  
 20 25 30  
 GCG CAA GCA CAG GAC GTC GTG GAC CTG GAC TTC TTC ACC CAG GAG CCG 206  
 Ala Gln Ala Gln Asp Val Val Asp Leu Asp Phe Phe Thr Gln Glu Pro  
 35 40 45

CTG CAC CTG GTG AGC CCC TCG TTC CTG TCC GTC ACC ATT GAC GCC AAC 254  
 Leu His Leu Val Ser Pro Ser Phe Leu Ser Val Thr Ile Asp Ala Asn  
 50 55 60

CTG GCC ACG GAC CCG CGG TTC CTC ATC CTC CTG GGT TCT CCA AAG CTT 302  
 Leu Ala Thr Asp Pro Arg Phe Leu Ile Leu Leu Gly Ser Pro Lys Leu  
 65 70 75 80

CGT ACC TTG GCC AGA GGC TTG TCT CCT GCG TAC CTG AGG TTT GGT GGC 350  
 Arg Thr Leu Ala Arg Gly Leu Ser Pro Ala Tyr Leu Arg Phe Gly Gly  
 85 90 95

ACC AAG ACA GAC TTC CTA ATT TTC GAT CCC AAG AAG GAA TCA ACC TTT 398  
 Thr Lys Thr Asp Phe Leu Ile Phe Asp Pro Lys Lys Glu Ser Thr Phe  
 100 105 110

GAA GAG AGA AGT TAC TGG CAA TCT CAA GTC AAC CAG GAT ATT TGC AAA 446  
 Glu Glu Arg Ser Tyr Trp Gln Ser Gln Val Asn Gln Asp Ile Cys Lys  
 115 120 125

TAT GGA TCC ATC CCT CCT GAT GTG GAG GAG AAG TTA CGG TTG GAA TGG 494  
 Tyr Gly Ser Ile Pro Pro Asp Val Glu Glu Lys Leu Arg Leu Glu Trp  
 130 135 140

CCC TAC CAG GAG CAA TTG CTA CTC CGA GAA CAC TAC CAG AAA AAG TTC 542  
 Pro Tyr Gln Glu Gln Leu Leu Leu Arg Glu His Tyr Gln Lys Lys Phe  
 145 150 155 160

AAG AAC AGC ACC TAC TCA AGA AGC TCT GTA GAT GTG CTA TAC ACT TTT 590  
 Lys Asn Ser Thr Tyr Ser Arg Ser Ser Val Asp Val Leu Tyr Thr Phe  
 165 170 175

GCA AAC TGC TCA GGA CTG GAC TTG ATC TTT GGC CTA AAT GCG TTA TTA 638  
 Ala Asn Cys Ser Gly Leu Asp Leu Ile Phe Gly Leu Asn Ala Leu Leu  
 180 185 190

AGA ACA GCA GAT TTG CAG TGG AAC AGT TCT AAT GCT CAG TTG CTC CTG 686  
 Arg Thr Ala Asp Leu Gln Trp Asn Ser Ser Asn Ala Gln Leu Leu Leu  
 195 200 205

GAC TAC TGC TCT TCC AAG GGG TAT AAC ATT TCT TGG GAA CTA GGC AAT 734  
 Asp Tyr Cys Ser Ser Lys Gly Tyr Asn Ile Ser Trp Glu Leu Gly Asn  
 210 215 220

GAA CCT AAC AGT TTC CTT AAG AAG GCT GAT ATT TTC ATC AAT GGG TCG 782  
 Glu Pro Asn Ser Phe Leu Lys Lys Ala Asp Ile Phe Ile Asn Gly Ser  
 225 230 235 240

CAG TTA GGA GAA GAT TAT ATT CAA TTG CAT AAA CTT CTA AGA AAG TCC 830  
 Gln Leu Gly Glu Asp Tyr Ile Gln Leu His Lys Leu Leu Arg Lys Ser  
 245 250 255

ACC TTC AAA AAT GCA AAA CTC TAT GGT CCT GAT GTT GGT CAG CCT CGA 878  
 Thr Phe Lys Asn Ala Lys Leu Tyr Gly Pro Asp Val Gly Gln Pro Arg  
 260 265 270

7

AGA AAG ACG GCT AAG ATG CTG AAG AGC TTC CTG AAG GCT GGT GGA GAA 926  
 Arg Lys Thr Ala Lys Met Leu Lys Ser Phe Leu Lys Ala Gly Gly Glu  
 275 280 285

GTG ATT GAT TCA GTT ACA TGG CAT CAC TAC TAT TTG AAT GGA CGG ACT 974  
 Val Ile Asp Ser Val Thr Trp His His Tyr Tyr Leu Asn Gly Arg Thr  
 290 295 300

GCT ACC AGG GAA GAT TTT CTA AAC CCT GAT GTA TTG GAC ATT TTT ATT 1022  
 Ala Thr Arg Glu Asp Phe Leu Asn Pro Asp Val Leu Asp Ile Phe Ile  
 305 310 315 320

TCA TCT GTG CAA AAA GTT TTC CAG GTG GTT GAG AGC ACC AGG CCT GGC 1070  
 Ser Ser Val Gln Lys Val Phe Gln Val Val Glu Ser Thr Arg Pro Gly  
 325 330 335

AAG AAG GTC TGG TTA GGA GAA ACA AGC TCT GCA TAT GGA GGC GGA GCG 1118  
 Lys Lys Val Trp Leu Gly Glu Thr Ser Ser Ala Tyr Gly Gly Gly Ala  
 340 345 350

CCC TTG CTA TCC GAC ACC TTT GCA GCT GGC TTT ATG TGG CTG GAT AAA 1166  
 Pro Leu Leu Ser Asp Thr Phe Ala Ala Gly Phe Met Trp Leu Asp Lys  
 355 360 365

TTG GGC CTG TCA GCC CGA ATG GGA ATA GAA GTG GTG ATG AGG CAA GTA 1214  
 Leu Gly Leu Ser Ala Arg Met Gly Ile Glu Val Val Met Arg Gln Val  
 370 375 380

TTC TTT GGA GCA GGA AAC TAC CAT TTA GTG GAT GAA AAC TTC GAT CCT 1262  
 Phe Phe Gly Ala Gly Asn Tyr His Leu Val Asp Glu Asn Phe Asp Pro  
 385 390 395 400

TTA CCT GAT TAT TGG CTA TCT CTT CTG TTC AAG AAA TTG GTG GGC ACC 1310  
 Leu Pro Asp Tyr Trp Leu Ser Leu Leu Phe Lys Lys Leu Val Gly Thr  
 405 410 415

AAG GTG TTA ATG GCA AGC GTG CAA GGT TCA AAG AGA AGG AAG CTT CGA 1358  
 Lys Val Leu Met Ala Ser Val Gln Gly Ser Lys Arg Arg Lys Leu Arg  
 420 425 430

GTA TAC CTT CAT TGC ACA AAC ACT GAC AAT CCA AGG TAT AAA GAA GGA 1406  
 Val Tyr Leu His Cys Thr Asn Thr Asp Asn Pro Arg Tyr Lys Glu Gly  
 435 440 445

GAT TTA ACT CTG TAT GCC ATA AAC CTC CAT AAC GTC ACC AAG TAC TTG 1454  
 Asp Leu Thr Leu Tyr Ala Ile Asn Leu His Asn Val Thr Lys Tyr Leu  
 450 455 460

CGG TTA CCC TAT CCT TTT TCT AAC AAG CAA GTG GAT AAA TAC CTT CTA 1502  
 Arg Leu Pro Tyr Pro Phe Ser Asn Lys Gln Val Asp Lys Tyr Leu Leu  
 465 470 475 480

AGA CCT TTG GGA CCT CAT GGA TTA CTT TCC AAA TCT GTC CAA CTC AAT 1550  
 Arg Pro Leu Gly Pro His Gly Leu Leu Ser Lys Ser Val Gln Leu Asn  
 485 490 495

GGT CTA ACT CTA AAG ATG GTG GAT GAT CAA ACC TTG CCA CCT TTA ATG 1598

8

Gly Leu Thr Leu Lys Met Val Asp Asp Gln Thr Leu Pro Pro Leu Met  
 500 505 510

GAA AAA CCT CTC CGG CCA GGA AGT TCA CTG GGC TTG CCA GCT TTC TCA 1646  
 Glu Lys Pro Leu Arg Pro Gly Ser Ser Leu Gly Leu Pro Ala Phe Ser  
 515 520 525

TAT AGT TTT TTT GTG ATA AGA AAT GCC AAA GTT GCT GCT TGC ATC TGA 1694  
 Tyr Ser Phe Phe Val Ile Arg Asn Ala Lys Val Ala Ala Cys Ile  
 530 535 540 543

AAA TAA AAT ATA CTA GTC CTG ACA CTG 1721

## (2) INFORMATION FOR SEQ ID NO:12:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 824  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:12

CTGGCAAGAA GGTCTGGTTG GGAGAGACGA GCTCAGCTTA CGGTGGCGGT GCACCCTTGC 60  
 TGTCACACAC CTTTCAGCT GGCITTATGT GGCTGGATAA ATTGGGCTG TCAGCCAGAG 120  
 TGGGCATAGA AGTCGTGATG AGGCAGGTGT TCTTCGGAGC AGGCAACTAC CACTTAGTGG 180  
 ATGAAAACCTT TGAGCCCTTA CCTGATTACT GGCTCTCTCT TCTGTTCAAG AAAGTGGTAG 240  
 GTCCAGGGT GTTACTGTCA AGAGTGAAAG GCCCAGACAG GAGCAAACTC CGAGTGTATC 300  
 TCCACTGCAC TAACGTCTAT CACCCACGAT ATCAGGAAGG AGATCTAACT CTGTATGTCC 360  
 TGAACCTCCA TAATGTCACC AAGCACTTGA AGGTACCGCC TCCGTTGTTT AGGAAACCAG 420  
 TGGATACGTA CCTTCTGAAG CCTTCGGGGC CGGATGGATT ACTTTCCAAA TCTGTCCAAC 480  
 TGAACGGTCA AATTCTGAAG ATGGTGGATG AGCAGACCCT GCCAGCTTTG ACAGAAAAAC 540  
 CTCTCCCGC AGGAAGTGCA CTAAGCCTGC CTGCCTTTTC CTATGGTTTT TTTGTCATAA 600  
 GAAATGCCAA AATCGCTGCT TGTATATGAA AATAAAGGC ATACGGTACC CCTGAGACAA 660  
 AAGCCGAGGG GGGTGTATT CATAAAACAA AACCTAGTT TAGGAGGCCA CCTCCTTGCC 720  
 GAGTTCAGAG GCTTCGGGAG GGTGGGGTAC ACTTCAGTAT TACATTCAGT GTGGTGTCT 780  
 CTCTAAGAAG AATACTGCAG GTGGTGACAG TTAATAGCAC TGTG 824

## (2) INFORMATION FOR SEQ ID NO:13:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1899  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:13

GGGAAAGCGA GCAAGGAAGT AGGAGAGAGC CGGGCAGGCG GGGCGGGGTT GGATTGGGAG 60  
 CAGTGGGAGG GATGCAGAAG AGGAGTGGGA GGGATGGAGG GCGCAGTGGG AGGGGTGAGG 120  
 AGGCGTAACG GGGCGGAGGA AAGGAGAAAA GGGCGCTGGG GCTCGGCGGG AGGAAGTGCT 180  
 AGAGCTCTCG ACTCTCCGCT GCGCGGCAGC TGGCGGGGGG AGCAGCCAGG TGAGCCCAAG 240  
 ATGCTGCTGC GCTCGAAGCC TGCCTGCCC GCGCGCTGA TGTGCTGCT CCTGGGGCCG 300  
 CTGGGTCCCC TCTCCCTGG CGCCCTGCC CGACCTGCGC AAGCACAGGA CGTCGTGGAC 360  
 CTGGACTTCT TCACCAGGA GCCGCTGCAC CTGGTGAGCC CCTCGTTCCT GTCCGTCACC 420  
 ATTGACGCCA ACCTGGCCAC GGACCCGCGG TTCCTCATCC TCCTGGGTTT TCCAAAGCTT 480  
 CGTACCTTGG CCAGAGGCTT GTCTCCTGCG TACCTGAGGT TTGGTGGCAC CAAGACAGAC 540  
 TTCTAATTT TCGATCCCAA GAAGGAATCA ACCTTTGAAG AGAGAAGTTA CTGGCAATCT 600  
 CRAAGTCAAC AGGATATTG CAAATATGGA TCCATCCCTC CTGATGTGGA GGAGAAGTTA 660  
 CGGTTGGAAT GGCCCTACCA GGAGCAATTG CTAATCCGAG AACACTACCA GAAAAAGTTC 720  
 AAGAACAGCA CTAATCAAG AAGCTCTGTA GATGTGCTAT ACACCTTTGC AAAGTGTCTA 780

9

GGACTGGACT TGATCTTTGG CCTAAATGCG TTATTAAGAA CAGCAGATTT GCAGTGGAAC 840  
 AGTTCTAATG CTCAGTTGCT CCTGGACTAC TGCTCTTCCA AGGGGTATAA CATTTCCTGG 900  
 GAACTAGGCA ATGAACCTAA CAGTTTCCTT AAGAAGGCTG ATATTTTCAT CAATGGGTCG 960  
 CAGTTAGGAG AAGATTATAT TCAATTGCAT AAACCTCTAA GAAAGTCCAC CTTCAAAAAT 1020  
 GCAAACTCT ATGGTCTCTGA TGTGGTCAG CCTCGAAGAA AGACGGCTAA GATGCTGAAG 1080  
 AGCTTCCTGA AGGCTGGTGG AGAAGTGATT GATTCAGTTA CATGGCATCA CTAATAITTTG 1140  
 AATGGACGGA CTGCTACCAG GGAAGATTTT CTAAACCCCTG ATGTATTGGA CATTTTTATT 1200  
 TCATCTGTGC AAAAAGTTTT CCAGGTGGTT GAGAGCACCA GGCCTGGCAA GAAGGTCTGG 1260  
 TTAGGAGAAA CAAGCTCTGC ATATGGAGGC GGAGCGCCCT TGCTATCCGA CACCTTTGCA 1320  
 GCTGGCTTTA TGTGGCTGGA TAAATPGGC CTGTGAGCCC GAATGGGAAT AGAAGTGGTG 1380  
 ATGAGGCAAG TATTCCTTTG AGCAGGAAAC TACCATTAGT TGGATGAAAA CTTCGATCCT 1440  
 TTACCTGATT ATGGCTATC TCTTCTGTC AAGAAATTGG TGGGCACCAA GGTGTTAATG 1500  
 GCAAGCGTGC AAGGTTCAAA GAGAAGGAAG CTTGAGTAT ACCTTCATTG CACAAACACT 1560  
 GACAAATCAA GGTATAAGA AGGAGATTTA ACTCTGTATG CCATAAACCT CCATAACGTC 1620  
 ACCAAGTACT TGCGGTTACC CTATCCTTTT TCTAACAGC AAGTGGATAA ATACCTTCTA 1680  
 AGACCTTTGG GACCTCATGG ATTACTTTCC AAATCTGTCC AACTCAATGG TCTAACTCTA 1740  
 AAGATGGTGG ATGATCAAAAC CTTGCCACCT TTAATGGAAA AACCTCTCCG GCCAGGAAGT 1800  
 TCACTGGGCT TGCCAGCTTT CTCATATAGT TTTTGTGA TAAGAAATGC CAAAGTTGCT 1860  
 GCTTGATCT GAAAATAAAA TATACTAGTC CTGACACTG 1899

## (2) INFORMATION FOR SEQ ID NO:14:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 592  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: singl  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:14

Met Glu Gly Ala Val Gly Gly Val Arg Arg Arg Asn Gly Ala Glu  
                   5                  10                  15  
 Glu Arg Arg Lys Gly Arg Trp Gly Ser Ala Gly Gly Ser Ala Arg  
                   20                  25                  30  
 Ala Leu Asp Ser Pro Leu Arg Gly Ser Trp Arg Gly Glu Gln Pro  
                   35                  40                  45  
 Gly Glu Pro Lys Met Leu Leu Arg Ser Lys Pro Ala Leu Pro Pro  
                   50                  55                  60  
 Pro Leu Met Leu Leu Leu Gly Pro Leu Gly Pro Leu Ser Pro  
                   65                  70                  75  
 Gly Ala Leu Pro Arg Pro Ala Gln Ala Gln Asp Val Val Asp Leu  
                   80                  85                  90  
 Asp Phe Phe Thr Gln Glu Pro Leu His Leu Val Ser Pro Ser Phe  
                   95                  100                 105  
 Leu Ser Val Thr Ile Asp Ala Asn Leu Ala Thr Asp Pro Arg Phe  
                  110                 115                 120  
 Leu Ile Leu Leu Gly Ser Pro Lys Leu Arg Thr Leu Ala Arg Gly  
                  125                 130                 135  
 Leu Ser Pro Ala Tyr Leu Arg Phe Gly Gly Thr Lys Thr Asp Phe  
                  140                 145                 150  
 Leu Ile Phe Asp Pro Lys Lys Glu Ser Thr Phe Glu Glu Arg Ser  
                  155                 160                 165  
 Tyr Trp Gln Ser Gln Val Asn Gln Asp Ile Cys Lys Tyr Gly Ser  
                  170                 175                 180  
 Ile Pro Pro Asp Val Glu Glu Lys Leu Arg Leu Glu Trp Pro Tyr  
                  185                 190                 195  
 Gln Glu Gln Leu Leu Leu Arg Glu His Tyr Gln Lys Lys Phe Lys  
                  200                 205                 210  
 Asn Ser Thr Tyr Ser Arg Ser Ser Val Asp Val Leu Tyr Thr Phe

10

215	220	225
Ala Asn Cys Ser Gly Leu Asp Leu Ile Phe Gly Leu Asn Ala Leu		
230	235	240
Leu Arg Thr Ala Asp Leu Gln Trp Asn Ser Ser Asn Ala Gln Leu		
245	250	255
Leu Leu Asp Tyr Cys Ser Ser Lys Gly Tyr Asn Ile Ser Trp Glu		
260	265	270
Leu Gly Asn Glu Pro Asn Ser Phe Leu Lys Lys Ala Asp Ile Phe		
275	280	285
Ile Asn Gly Ser Gln Leu Gly Glu Asp Tyr Ile Gln Leu His Lys		
290	295	300
Leu Leu Arg Lys Ser Thr Phe Lys Asn Ala Lys Leu Tyr Gly Pro		
305	310	315
Asp Val Gly Gln Pro Arg Arg Lys Thr Ala Lys Met Leu Lys Ser		
320	325	330
Phe Leu Lys Ala Gly Gly Glu Val Ile Asp Ser Val Thr Trp His		
335	340	345
His Tyr Tyr Leu Asn Gly Arg Thr Ala Thr Arg Glu Asp Phe Leu		
350	355	360
Asn Pro Asp Val Leu Asp Ile Phe Ile Ser Ser Val Gln Lys Val		
365	370	375
Phe Gln Val Val Glu Ser Thr Arg Pro Gly Lys Lys Val Trp Leu		
380	385	390
Gly Glu Thr Ser Ser Ala Tyr Gly Gly Gly Ala Pro Leu Leu Ser		
395	400	405
Asp Thr Phe Ala Ala Gly Phe Met Trp Leu Asp Lys Leu Gly Leu		
410	415	420
Ser Ala Arg Met Gly Ile Glu Val Val Met Arg Gln Val Phe Phe		
425	430	435
Gly Ala Gly Asn Tyr His Leu Val Asp Glu Asn Phe Asp Pro Leu		
440	445	450
Pro Asp Tyr Trp Leu Ser Leu Leu Phe Lys Lys Leu Val Gly Thr		
455	460	465
Lys Val Leu Met Ala Ser Val Gln Gly Ser Lys Arg Arg Lys Leu		
470	475	480
Arg Val Tyr Leu His Cys Thr Asn Thr Asp Asn Pro Arg Tyr Lys		
485	490	495
Glu Gly Asp Leu Thr Leu Tyr Ala Ile Asn Leu His Asn Val Thr		
500	505	510
Lys Tyr Leu Arg Leu Pro Tyr Pro Phe Ser Asn Lys Gln Val Asp		
515	520	525
Lys Tyr Leu Leu Arg Pro Leu Gly Pro His Gly Leu Leu Ser Lys		
530	535	540
Ser Val Gln Leu Asn Gly Leu Thr Leu Lys Met Val Asp Asp Gln		
545	550	555
Thr Leu Pro Pro Leu Met Glu Lys Pro Leu Arg Pro Gly Ser Ser		
560	565	570
Leu Gly Leu Pro Ala Phe Ser Tyr Ser Phe Phe Val Ile Arg Asn		
575	580	585
Ala Lys Val Ala Ala Cys Ile		
590	592	

## (2) INFORMATION FOR SEQ ID NO:15:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1899  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double

11

(D) TOPOLOGY: linear  
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15

```

                                GGG      3
AAA GCG AGC AAG GAA GTA GGA GAG AGC CGG GCA GGC GGG GCG GGG  48
TTG GAT TGG GAG CAG TGG GAG GGA TGC AGA AGA GGA GTG GGA GGG  93
ATG GAG GGC GCA GTG GGA GGG GTG AGG AGG CGT AAC GGG GCG GAG  138
Met Glu Gly Ala Val Gly Gly Val Arg Arg Arg Asn Gly Ala Glu
      5              10              15

GAA AGG AGA AAA GGG CGC TGG GGC TCG GCG GGA GGA AGT GCT AGA  183
Glu Arg Arg Lys Gly Arg Trp Gly Ser Ala Gly Gly Ser Ala Arg
      20              25              30

GCT CTC GAC TCT CCG CTG CGC GGC AGC TGG CGG GGG GAG CAG CCA  228
Ala Leu Asp Ser Pro Leu Arg Gly Ser Trp Arg Gly Glu Gln Pro
      35              40              45

GGT GAG CCC AAG ATG CTG CTG CGC TCG AAG CCT GCG CTG CCG CCG  273
Gly Glu Pro Lys Met Leu Leu Arg Ser Lys Pro Ala Leu Pro Pro
      50              55              60

CCG CTG ATG CTG CTG CTC CTG GGG CCG CTG GGT CCC CTC TCC CCT  318
Pro Leu Met Leu Leu Leu Gly Pro Leu Gly Pro Leu Ser Pro
      65              70              75

GGC GCC CTG CCC CGA CCT GCG CAA GCA CAG GAC GTC GTG GAC CTG  363
Gly Ala Leu Pro Arg Pro Ala Gln Ala Gln Asp Val Val Asp Leu
      80              85              90

GAC TTC TTC ACC CAG GAG CCG CTG CAC CTG GTG AGC CCC TCG TTC  408
Asp Phe Phe Thr Gln Glu Pro Leu His Leu Val Ser Pro Ser Phe
      95              100             105

CTG TCC GTC ACC ATT GAC GCC AAC CTG GCC ACG GAC CCG CGG TTC  453
Leu Ser Val Thr Ile Asp Ala Asn Leu Ala Thr Asp Pro Arg Phe
     110              115             120

CTC ATC CTC CTG GGT TCT CCA AAG CTT CGT ACC TTG GCC AGA GGC  498
Leu Ile Leu Leu Gly Ser Pro Lys Leu Arg Thr Leu Ala Arg Gly
     125              130             135

TTG TCT CCT GCG TAC CTG AGG TTT GGT GGC ACC AAG ACA GAC TTC  543
Leu Ser Pro Ala Tyr Leu Arg Phe Gly Gly Thr Lys Thr Asp Phe
     140              145             150

CTA ATT TTC GAT CCC AAG AAG GAA TCA ACC TTT GAA GAG AGA AGT  588
Leu Ile Phe Asp Pro Lys Lys Glu Ser Thr Phe Glu Glu Arg Ser
     155              160             165

TAC TGG CAA TCT CAA GTC AAC CAG GAT ATT TGC AAA TAT GGA TCC  633
Tyr Trp Gln Ser Gln Val Asn Gln Asp Ile Cys Lys Tyr Gly Ser
     170              175             180

ATC CCT CCT GAT GTG GAG GAG AAG TTA CGG TTG GAA TGG CCC TAC  678
Ile Pro Pro Asp Val Glu Glu Lys Leu Arg Leu Glu Trp Pro Tyr
     185              190             195

```



CAG GAG CAA TTG CTA CTC CGA GAA CAC TAC CAG AAA AAG TTC AAG	723
Gln Glu Gln Leu Leu Leu Arg Glu His Tyr Gln Lys Lys Phe Lys	
200 205 210	
AAC AGC ACC TAC TCA AGA AGC TCT GTA GAT GTG CTA TAC ACT TTT	768
Asn Ser Thr Tyr Ser Arg Ser Ser Val Asp Val Leu Tyr Thr Phe	
215 220 225	
GCA AAC TGC TCA GGA CTG GAC TTG ATC TTT GGC CTA AAT GCG TTA	813
Ala Asn Cys Ser Gly Leu Asp Leu Ile Phe Gly Leu Asn Ala Leu	
230 235 240	
TTA AGA ACA GCA GAT TTG CAG TGG AAC AGT TCT AAT GCT CAG TTG	858
Leu Arg Thr Ala Asp Leu Gln Trp Asn Ser Ser Asn Ala Gln Leu	
245 250 255	
CTC CTG GAC TAC TGC TCT TCC AAG GGG TAT AAC ATT TCT TGG GAA	903
Leu Leu Asp Tyr Cys Ser Ser Lys Gly Tyr Asn Ile Ser Trp Glu	
260 265 270	
CTA GGC AAT GAA CCT AAC AGT TTC CTT AAG AAG GCT GAT ATT TTC	948
Leu Gly Asn Glu Pro Asn Ser Phe Leu Lys Lys Ala Asp Ile Phe	
275 280 285	
ATC AAT GGG TCG CAG TTA GGA GAA GAT TAT ATT CAA TTG CAT AAA	993
Ile Asn Gly Ser Gln Leu Gly Glu Asp Tyr Ile Gln Leu His Lys	
290 295 300	
CTT CTA AGA AAG TCC ACC TTC AAA AAT GCA AAA CTC TAT GGT CCT	1038
Leu Leu Arg Lys Ser Thr Phe Lys Asn Ala Lys Leu Tyr Gly Pro	
305 310 315	
GAT GTT GGT CAG CCT CGA AGA AAG ACG GCT AAG ATG CTG AAG AGC	1083
Asp Val Gly Gln Pro Arg Arg Lys Thr Ala Lys Met Leu Lys Ser	
320 325 330	
TTC CTG AAG GCT GGT GGA GAA GTG ATT GAT TCA GTT ACA TGG CAT	1128
Phe Leu Lys Ala Gly Gly Glu Val Ile Asp Ser Val Thr Trp His	
335 340 345	
CAC TAC TAT TTG AAT GGA CGG ACT GCT ACC AGG GAA GAT TTT CTA	1173
His Tyr Tyr Leu Asn Gly Arg Thr Ala Thr Arg Glu Asp Phe Leu	
350 355 360	
AAC CCT GAT GTA TTG GAC ATT TTT ATT TCA TCT GTG CAA AAA GTT	1218
Asn Pro Asp Val Leu Asp Ile Phe Ile Ser Ser Val Gln Lys Val	
365 370 375	
TTC CAG GTG GTT GAG AGC ACC AGG CCT GGC AAG AAG GTC TGG TTA	1263
Phe Gln Val Val Glu Ser Thr Arg Pro Gly Lys Lys Val Trp Leu	
380 385 390	
GGA GAA ACA AGC TCT GCA TAT GGA GGC GGA GCG CCC TTG CTA TCC	1308
Gly Glu Thr Ser Ser Ala Tyr Gly Gly Gly Ala Pro Leu Leu Ser	
395 400 405	

## 13

GAC ACC TTT GCA GCT GGC TTT ATG TGG CTG GAT AAA TTG GGC CTG	1353
Asp Thr Phe Ala Ala Gly Phe Met Trp Leu Asp Lys Leu Gly Leu	
410 415 420	
TCA GCC CGA ATG GGA ATA GAA GTG GTG ATG AGG CAA GTA TTC TTT	1398
Ser Ala Arg Met Gly Ile Glu Val Val Met Arg Gln Val Phe Phe	
425 430 435	
GGA GCA GGA AAC TAC CAT TTA GTG GAT GAA AAC TTC GAT CCT TTA	1443
Gly Ala Gly Asn Tyr His Leu Val Asp Glu Asn Phe Asp Pro Leu	
440 445 450	
CCT GAT TAT TGG CTA TCT CTT CTG TTC AAG AAA TTG GTG GGC ACC	1488
Pro Asp Tyr Trp Leu Ser Leu Leu Phe Lys Lys Leu Val Gly Thr	
455 460 465	
AAG GTG TTA ATG GCA AGC GTG CAA GGT TCA AAG AGA AGG AAG CTT	1533
Lys Val Leu Met Ala Ser Val Gln Gly Ser Lys Arg Arg Lys Leu	
470 475 480	
CGA GTA TAC CTT CAT TGC ACA AAC ACT GAC AAT CCA AGG TAT AAA	1578
Arg Val Tyr Leu His Cys Thr Asn Thr Asp Asn Pro Arg Tyr Lys	
485 490 495	
GAA GGA GAT TTA ACT CTG TAT GCC ATA AAC CTC CAT AAC GTC ACC	1623
Glu Gly Asp Leu Thr Leu Tyr Ala Ile Asn Leu His Asn Val Thr	
500 505 510	
AAG TAC TTG CGG TTA CCC TAT CCT TTT TCT AAC AAG CAA GTG GAT	1668
Lys Tyr Leu Arg Leu Pro Tyr Pro Phe Ser Asn Lys Gln Val Asp	
515 520 525	
AAA TAC CTT CTA AGA CCT TTG GGA CCT CAT GGA TTA CTT TCC AAA	1713
Lys Tyr Leu Leu Arg Pro Leu Gly Pro His Gly Leu Leu Ser Lys	
530 535 540	
TCT GTC CAA CTC AAT GGT CTA ACT CTA AAG ATG GTG GAT GAT CAA	1758
Ser Val Gln Leu Asn Gly Leu Thr Leu Lys Met Val Asp Asp Gln	
545 550 555	
ACC TTG CCA CCT TTA ATG GAA AAA CCT CTC CGG CCA GGA AGT TCA	1803
Thr Leu Pro Pro Leu Met Glu Lys Pro Leu Arg Pro Gly Ser Ser	
560 565 570	
CTG GGC TTG CCA GCT TTC TCA TAT AGT TTT TTT GTG ATA AGA AAT	1848
Leu Gly Leu Pro Ala Phe Ser Tyr Ser Phe Phe Val Ile Arg Asn	
575 580 585	
GCC AAA GTT GCT GCT TGC ATC TGA AAA TAA AAT ATA CTA GTC CTG	1893
Ala Lys Val Ala Ala Cys Ile	
590 592	
ACA CTG	1899

## (2) INFORMATION FOR SEQ ID NO:16:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 594  
 (B) TYPE: nucleic acid

14

- (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear  
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:16

```

ATTACTATAG GGCACGCGTG GTCGACGGCC CGGGCTGGTA TTGTCTTAAT GAGAAGTTGA 60
TAAAGAATTT TGGGTGGTTG ATCTCTTTCC AGCTGCAGTT TAGCGTATGC TGAGGCCAGA 120
TTTTTTCAGG CAAAAGTAAA ATACCTGAGA AACTGCCTGG CCAGAGGACA ATCAGATTTT 180
GGCTGGCTCA AGTGACAAGC AAGTGTTTAT AAGCTAGATG GGAGAGGAAG GGATGAATAC 240
TCCATTGGAG GCTTTACTCG AGGGTCAGAG GGATACCCGG CGCCATCAGA ATGGGATCTG 300
GGAGTCGAA ACGCTGGGTT CCCACGAGAG CGCGCAGAAC ACGTGCCTCA GGAAGCCTGG 360
TCCGGGATGC CCAGCGCTGC TCCCCGGGCG CTCCTCCCCG GGCGCTCCTC CCCAGGCCTC 420
CCGGGCGCTT GGATCCCGGC CATCTCCGCA CCCTTCAAGT GGGTGTGGGT GATTTCGTAA 480
GTGAACGTGA CCGCCACCGG GGGGAAAGCG AGCAAGGAAG TAGGAGAGAG CCGGGCAGGC 540
GGGGCGGGT TGGATTGGGA GCAGTGGGAG GGATGCAGAA GAGGAGTGGG AGGG 594

```

- (2) INFORMATION FOR SEQ ID NO:17:  
 (i) SEQUENCE CHARACTERISTICS:  
   (A) LENGTH: 21  
   (B) TYPE: nucleic acid  
   (C) STRANDEDNESS: single  
   (D) TOPOLOGY: linear  
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:17  
 CCCCAGGAGC AGCAGCATCA G 21

- (2) INFORMATION FOR SEQ ID NO:18:  
 (i) SEQUENCE CHARACTERISTICS:  
   (A) LENGTH: 21  
   (B) TYPE: nucleic acid  
   (C) STRANDEDNESS: single  
   (D) TOPOLOGY: linear  
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:18  
 AGGCTTCGAG CGCAGCAGCA T 21

- (2) INFORMATION FOR SEQ ID NO:19:  
 (i) SEQUENCE CHARACTERISTICS:  
   (A) LENGTH: 22  
   (B) TYPE: nucleic acid  
   (C) STRANDEDNESS: single  
   (D) TOPOLOGY: linear  
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:19  
 GTAATACGAC TCACTATAGG GC 22

- (2) INFORMATION FOR SEQ ID NO:20:  
 (i) SEQUENCE CHARACTERISTICS:  
   (A) LENGTH: 19  
   (B) TYPE: nucleic acid  
   (C) STRANDEDNESS: single  
   (D) TOPOLOGY: linear  
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:20  
 ACTATAGGCG ACGCGTGGT 19

- (2) INFORMATION FOR SEQ ID NO:21:  
 (i) SEQUENCE CHARACTERISTICS:  
   (A) LENGTH: 21  
   (B) TYPE: nucleic acid  
   (C) STRANDEDNESS: single  
   (D) TOPOLOGY: linear

15

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:21  
CTTGGGCTCA CCTGGCTGCT C 21
- (2) INFORMATION FOR SEQ ID NO:22:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 23
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:22  
AGCTCTGTAG ATGTGCTATA CAC 23
- (2) INFORMATION FOR SEQ ID NO:23:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 22
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:23  
GCATCTTAGC CGTCTTTCTT CG 22
- (2) INFORMATION FOR SEQ ID NO:24:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 23
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:24  
GAGCAGCCAG GTGAGCCCAA GAT 23
- (2) INFORMATION FOR SEQ ID NO:25:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 23
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:25  
TTCGATCCCA AGAAGGAATC AAC 23
- (2) INFORMATION FOR SEQ ID NO:26:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 23
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:26  
AGCTCTGTAG ATGTGCTATA CAC 23
- (2) INFORMATION FOR SEQ ID NO:27:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 24
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:27  
TCAGATGCAA GCAGCAACTT TGGC 24

16

- (2) INFORMATION FOR SEQ ID NO:28:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 22
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:28  
GCATCTTAGC CGTCTTCTT CG 22
- (2) INFORMATION FOR SEQ ID NO:29:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 24
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:29  
GTAGTGATGC CATGTAAC TG AATC 24
- (2) INFORMATION FOR SEQ ID NO:30:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 22
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:30  
AGGCACCCCTA GAGATGTTCC AG 22
- (2) INFORMATION FOR SEQ ID NO:31:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 24
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:31  
GAAGATTCTT GTTCCATGA CGTG 24
- (2) INFORMATION FOR SEQ ID NO:32:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 25
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:32  
CCACACTGAA TGTAACTG AAGTG 25
- (2) INFORMATION FOR SEQ ID NO:33:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 22
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:33  
CGAAGCTCTG GAACTCGGCA AG 22
- (2) INFORMATION FOR SEQ ID NO:34:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 22

17

- (B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34  
GCCAGCTGCA AAGGTGTTGG AC 22
- (2) INFORMATION FOR SEQ ID NO:35:  
(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 23  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35  
AACACCTGCC TCATCAGCAGC TTC 23
- (2) INFORMATION FOR SEQ ID NO:36:  
(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 22  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36  
GCCAGGCTGG CGTCGATGGT GA 22
- (2) INFORMATION FOR SEQ ID NO:37:  
(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 22  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37  
GTCGATGGTG ATGGACAGGA AC 22
- (2) INFORMATION FOR SEQ ID NO:38:  
(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 22  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38  
GTAATACGAC TCACTATAGG GC 22
- (2) INFORMATION FOR SEQ ID NO:39:  
(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 19  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39  
ACTATAGGGC ACGCGTGGT 19
- (2) INFORMATION FOR SEQ ID NO:40:  
(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 27  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

18

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40  
CCATCCTAAT ACGACTCACT ATAGGCG 27

(2) INFORMATION FOR SEQ ID NO:41:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 23  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41  
ACTCACTATA GGGCTCGAGC GGC 23

(2) INFORMATION FOR SEQ ID NO:42:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 44848  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: double  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42

GGATCTTGGC TCACTGCAAT CTCTGCCTCC CATGCAATTC TTATGCATCA 50  
GCCTCCTGAG TAGCTTGGAT TATAGGTCTG CGCCACCACT CCTGGCTACA 100  
CCATGTTGCC CAGGCTGGTC TTGAACCTTT GGGCTCTAGT GATCCACCCG 150  
CCTTGGCCTC CCAAAGTGCT GGGATTACAG GTGTGAGCCA TCACACCCGG 200  
CCCCCGTTT CCATATTAGT AACTCACATG TAGACCACAA GGATGCACTA 250  
TTTAGAAAAC TTGCAATGGT CCACTTTTCA AATCACCCAA ACATGTTAAA 300  
GAAATTTGTA TGACTGGGCA TGGCACAGTG GCTCATGCCT GCAATCCTAG 350  
CATTTTGTGA GGCTGAGACG GGCAGATCAC GAGGTACAGG GATTGAGACC 400  
ATCCTGACAG ACATGGTGAA ATCCCATCTC TACTAAAAAT ACAAAACAAT 450  
TAGCCGGGGG TGATGGCAGG CCCCTGTAGT CCCAGCTACT CGGGAGGCTG 500  
AGGCAGGAGA ATGGCGTGAA TCCAGGAGGC AGAGCTTGCA GTGAGCCGAG 550  
ATGGTGGCAC TGCACTCCAG CCTGGGCGAC AGAGCGAGAC TCCGTCTCAA 600  
AAAAAAAAAA AAAGAAAGAA ATTGGTATGA CTGTTGACTC ACAACAGGAG 650  
TCAGGGGCAT GGGGTGGGGT GTAAGATTAA TGTCAAGACA AATGTGGAAA 700  
AGAAACTTCT GTTTTTCCAA CTCCACGTCT GCTACCATAT TATTACACTC 750  
TTCTGGTAGT GTGGTGTGTA TGTGTGAATT TTTTTCATA TGTATACAGT 800  
AATTGTAGGA TATGAACCTG ATTCTAGTTG CAAAACCTAC TATGAGCTTA 850  
GCTTTTAAGT TGCTTAAGAA TAGGTAGATC TATGCAAATA ATGATAATTA 900  
TTATTATTAT TTTAAGAGAG GGTCTCACTT TGTACCCAG GCTGGAGTGC 950  
AGTGGTGTGA TTAAGGGTCA CTGCAACCTC CACCTCCCAG GCTCAAATAA 1000  
ACCTCCCACC TCAGCCTCCC CAGTAGCTGG AACCACAGGC ACGGGCCACC 1050  
ACGCTGGCT AATTTTTTGT ATTTTTTGTA GAGATGGGGT TTCATCATGT 1100  
TGCCAGGCT GTTCTTGAAT TCCTCGGCTC AAGCAATCCT CCCACCTTGG 1150  
CCTCCCAAAA TGCTGGCATC ACAGGCATGA TGGCATCACT GGCATCAGAT 1200  
ACCATGCCTG GCCTGATTTA TGCAAATTAG ATATGCATT TCAAATAATC 1250  
TATTTTTATT TGTTGCCTTA TTGGTGGTAC AATCTCAAGT GGAATAATCT 1300  
AAGGGTTTTG GTGTTATTTG CTACTCAAC CAATATTTAT TAGACTCTTA 1350  
CTAAGCACCA ACATGATCAC ATGCCTGAGC TATGGCTAGC ATAGCGTGTG 1400  
AGACAAACTT AATCTCTGTT TTGGTGGAGC ATATAATCTA GTAGATGAAG 1450  
CCAATGTTGA GCAACATCAC AATACTAACA AATTGAGGAT GCTACGAGAG 1500  
TGTCTAACAA ATTGAGGATG CTACGAGAGT GTCTAACAAA TTGAGGATGC 1550  
TATGAGAGTG TGTATGGAG AGCTGCCTGG AGATTGAGAG AAAGCTTCCT 1600  
TGAGGGAAGT TACATTTGAG CTGAAACACA CTGCCATCTG CTCGAGGTTT 1650  
TGTAACGTCA TTCACATCCC GATTCTGACA CTTCACATCC CGATTCTGAC 1700  
ACTTCACCCA GTTACTGTCT CAGAGCTTGG GTCCGCATGT GTAAAACAAG 1750  
GACAGTATGC ACTTGGCAGG GTTGTGAGAA GGGAAAGAGAA CACAAGTAAA 1800  
GCACCTGTAT CAGGCATACA GTAGGCACTA AGCGTGCGAT GCTTGCTATG 1850  
ATTATACATC AGTGTAAACA TCAAGGAAAA GCTGAAGAAA AGTCTGACCA 1900  
ACAGCGAAG ATAAATGCGC AGAGGAGAAA TTTGGCAAAG GCTCCAAATT 1950  
CAGGGGCACT CCGTACTCTA CACTTTGTAT GGGGGCTTCA GGTCTTGAGT 2000  
TCCAGACATT GGAGCAACTA ACCCTTTAAG ATTGCTAAAT ATTGTCTTAA 2050  
TGAGGAAGTG ATAAAGAATT TTGGGTGGTT GATCTCTTTC CAGCTGCAGT 2100  
TTAGCGTAGT CTGAGGCCAG ATTTTTCCTA GCAAAAAGTAA AATACCTGAG 2150  
AAACTGCTTG GCCAGAGGAC AATCAGATT TGGCTGGCTC AAGTGACAAG 2200  
CAAGTGTTTA TAAGCTAGAT GGGAGAGGAA GGGATGAATA CTCCATTGGA 2250  
GGTTTTACTC GAGGGTCAGA GGGATACCCG GCGCCATCAG AATGGGATCT 2300  
GGGAGTCGGA AACGCTGGGT TCCCACGAGA GCGCGCAGAA CAGTGCCTC 2350  
AGGAAGCCTG GTCCGGGATG CCCAGCGCTG CTCCCCGGGC GCTCCTCCCC 2400

GGGCGCTCCT	CCCCAGGCCT	CCCGGGCGCT	TGGATCCCG	CCATCTCCGC	2450
ACCCTTCAAG	TGGGTGTGGG	TGATTTCTGT	AGTGAACGTG	ACCGCCACCG	2500
AGGGGAAAGC	GAGCAAGGAA	GTAGGAGAGA	GCCGGGCAGG	CGGGGCGGGG	2550
TTGGATTGGG	AGCAGTGGGA	GGGATGCAGA	AGAGGAGTGG	GAGGGATGGA	2600
GGGCGCAGTG	GGAGGGGTGA	GGAGGCGTAA	CGGGGCGGAG	GAAAGGAGAA	2650
AAGGGCGCTG	GGGCTCGGCG	GGAGGAAGTG	CTAGAGCTCT	CGACTCTCCG	2700
CTGCGCGGCA	GCTGGCGGGG	GGAGCAGCCA	GGTGAGCCCA	AGATGCTGCT	2750
CGCTCGAAG	CCTGCGCTGC	CGCGCGCGCT	GATGCTGCTG	CTCTGGGGGC	2800
CGCTGGGTCC	CCTCTCCCTT	GGCGCCCTGC	CCCGACCTGC	GCAAGCACAG	2850
GACGTCTGTG	ACCTGGACTT	CTTCACCCAG	GAGCGGCTGC	ACCTGGTGAG	2900
CCCTCGTTTC	CTGTCCGTCA	CCATTGACGC	CAACCTGGCC	ACGGACCCGC	2950
GGTTCCTCAT	CCTCTGGGGG	TAAGCGCCAG	CCTCCTGGTC	CTGTCCCTTT	3000
TCCTGTCTCT	CTGACACCTA	TGTCTGCCCC	GCCAGCGGCT	CTCCTTCTTT	3050
TGCGCGGAAA	CAACTTCACA	CCGGAACCTC	CCCGCTGTGC	TCTCCCAACC	3100
CCACTTCCCG	CCTCTCATTC	TCCCTCTCCC	TCCCTTACTC	TCAGACCCCA	3150
AACCGCTTTT	TGGGGGGTAT	CATTTAAAAA	ATAGATTAGG	GGGTACCAAG	3200
TGCAGTTCTG	TTCCATGGGT	ATATTGCATT	GTGGTGGCAT	CTGGGCTCTT	3250
AGTGTAAGTG	TCACCCGAAT	GTTGTACATT	GTATCTAATA	GGTAATTCTT	3300
CATCCCTCAT	CCCTCTCCCA	CCCTCCACCC	TTTTGGAGTC	TCCAGTGTCT	3350
ACTATTCCAC	TAAGTCCATG	TGTACACATT	GTTTAGCGCC	CACCTCTAAAT	3400
GAGCCTTTTT	GTTTCATTCA	TTCTGTAAAT	GTTGAATAGG	CACCACCTAA	3450
GGTCAGGTAT	AAGTGGAAAT	TTGAAAAAGA	AACCTGCCAC	TTGCCCCAGT	3500
ACTTCCCTAG	CCAAGAGGAG	GGAACACAGG	CAGGTGCACC	TGAAGGCCCTG	3550
TGAGTGCTTG	ATTGTCTGTG	CAGTGTAGGA	CAAGTAAGAT	TGTGCATAGC	3600
CTTCTGTATT	TAAGACTGTG	TTAGGAAGAT	TTCTCTTTCT	TTTCTTTTCT	3650
TTTTCTTTTT	TCTTTTCTTT	TTTTTTTTTA	GGCAGATGAA	AAGGGCGTCA	3700
CAGAACAGGA	ATAAAATCT	AAATATTCAA	TAAATGAGAC	CTAGGAGACT	3750
ACTGCAGTGA	CTTACAAAGT	CCTAATAAAA	AGATGTCTCT	CCAAAATGGG	3800
GCTGCAAAAT	GTGGTGCTGC	CTTATCAGCT	CTAAGTTTTT	TCCTTACCTG	3850
AGAAAGAAAG	AACCTGATGC	AGGTTCAGGG	CTCCTGCCCC	ATGAATGCAG	3900
GCTGACTCCA	AGATGGGGAG	CTACAGGGAC	AATCCCAAGT	CTTCTAGGCC	3950
TCTTATTTAG	GCCCTGGGAG	CCTCCAGAGA	TGGCCACATC	TTGACCAGCC	4000
CAGATAGAGG	GAAAGATCAC	CATTATCTCA	CCTCTGTGTC	AAATACCTAG	4050
ATGCTGTCTT	CCCTGAGCCC	ACACTATAGT	TGCCAGCGCT	AATTTAATGG	4100
GTAGTGTACT	GGTTAAGAGA	TGGACAGACC	ATCCTGGCTT	GACTCTCAGC	4150
TCTGGCAAGG	ATGAGTGAAT	TGGTTTTTCC	ATATCTCTTG	GCCACACCAA	4200
CCTTGATTTC	TTCACTGTGA	GAATGGAATT	TCTCAAGCTT	GCCTCAAGGA	4250
TTATTGCCCG	AGGATTGTAT	GATATGGTAA	GAGCTTCTCA	GTGTTTGACC	4300
CATAGTAAGT	GTTTGACGTT	TCAAACGAAT	TGTTTCTTTC	TAGGACATGG	4350
TGAGCAATTT	GTAGCCATTC	ACCGGTTTTT	TGTTTCTTTG	GATCATAGTT	4400
AACCTCTCCT	TTTCTTCTTG	GCACTACAAT	TTTCTGGTGG	GGAAGAATCC	4450
TTACTTTCTG	CCCTTCCCTT	TAAGGATAGG	AAGCTGATAC	TAGGCAGCAA	4500
CTAGTTGGGG	GATAGGAAGA	TTGTTCCAGA	GAAATGCTGA	ACCATAGGGC	4550
TCCAGATCAC	AGGACCCGAG	TCTTAGCTTG	CTGGGGTGTG	GGGTGGGGGG	4600
GGGCGGTTAC	TGAACATGGG	TATGAAGTAG	ATGTCCATT	ACTGAAATGT	4650
GAGGACCTGA	GGCTCTTCT	ATTGCTGTAG	CCAGCATATT	CCCCAACCTC	4700
TCCCAAGAA	AGGACAGATG	GGGGTTCCCC	CCTGGAGTAA	CAGGTCCAAA	4750
AGAAAAAACA	TACAGTGGGA	CTTCCAGGAT	CTGGGCTCTG	TCACCCAGCA	4800
GTCAAGCTCC	CCGCAATTGA	CTAACACCCC	CCTAACACGT	AGAAATTCCT	4850
ATCTGCAATT	TAGTGAGGAT	GATACCTTTA	TTCTTCTTAA	ATACATCTCT	4900
TCATTTCCCA	GAGCACCTTT	TTTTCCCTTC	CTCTGCACCT	TTTTGTTAAA	4950
GACTGGAGTA	TAATGAAATA	CCAAGAGAGC	ATAACATGTG	ATACATAAAA	5000
CTTTTTTTCT	GGTTTACAAA	ACAGTTTCAAT	CTTGTCCATA	CGTGCTTCTC	5050
TCCAAGGCTG	GCTGCTGTCT	GTTCCAGCCC	GCTTGGCTTG	GAGAGGCCAT	5100
CTGCCATACC	TGCTCCCCAG	AOCATCGAC	AAGCACACCC	AGAGTGTAT	5150
CTGCTAAGAC	CTAAAAGAGG	GAGGAACCCC	CTCTCTCAT	CTAAGACCTA	5200
GCTTCTAAAT	TAGAGTGTGA	GGGTCCATCT	CCCCAGGAGG	GGCACAGGGC	5250
CCAAACAGCC	CAGCCATCTC	AGAAGACAAC	ACTAAGCTTT	GTAGGGGTCC	5300
ACAGTAGAGG	AGAGTAAGAC	GCCTGTTGTT	TAATTTATTA	CAGTTCCTCA	5350
AAAGTGAAGA	TGTGTGGGCG	GGATGGCAAG	AGCTGAGCAG	ACGAAAGCTG	5400
AAGGAATAAG	GAAAGAGAGG	AGGACACAAA	CAGCTGACAC	TTCTCTAGTT	5450
CTTGTCAATT	GCCTGGCCCT	GTTCTAAGCA	CCTTCTAGGT	ATTAATCCAT	5500
TTAGTCTTGG	CTACAACACT	GTGAGTAACT	AGTTTTGTCA	CCCCCATTTT	5550
AAAAATGAAG	AAAGTGAGGC	TCAGGGAGGT	TAAGTAACTT	GGCCACAGTT	5600
TGAACTAGAG	CTCTGATCAC	ATGAGATAAT	AGTGCCCAT	AAAAGGGAAA	5650
GCAGATTATA	TTTTTTAAAG	GAAAGAGAGT	AGGATATGGT	AGAAAAAGAT	5700
TGTTTGGAAA	GGAATTGAGA	GATTGATATA	ATGAAAAGAA	GCATTACAT	5750
GAGAGTAACA	GTATCAGGGC	CCAAACCTTC	ATCTAAGGTA	CTTCAAGAG	5800
GCCTAAGCAA	ACTTAGTCAC	TGGCGTGGTT	CTAGTCTCCA	TGATGGCAAA	5850
TACATTGTGT	ACAGCCCAAC	TCCACACAAA	ACTTAAATAC	CAATGATAGA	5900
GCAATCTAAA	ATTTGAAAAG	AAAAATCTTT	CAATTTGTCT	TCTTCCAGTA	5950
GGGACTTAAT	CAAGAAACCA	ATCAAAATAC	TTCTTAAGCC	TAACTGTGTG	6000
CAGAACTCCA	AAGAGAGCCC	AGCCCTAAAT	CAACACTGTC	CAATGGAAAT	6050
ATAATATAAT	GTGGGCTTCA	TATGCAAGGT	CATATGTAAT	TTTAAATTTT	6100
CTAGTAGCCA	TATTAATAAG	GTAATAAGAA	ACAAGTGAAG	TTAATTTTAA	6150



TAATTTTATT TAGTTCATA GATCCAAAAT GTTTTCTCAG CATGTAATCA 6200  
ATATAAAAAT ATTAATGAGG TATTTATTAT TCCTTTTCTC AAACCAAGTC 6250  
TATTTCTATA TCTGGCGTGT ATTATTITACA GCACCTTCTCA GACTATATTT 6300  
CTTTCTTTCT TTTTITTTTC CGAGACAATT TTGCTCTTGT CACCCAAGCT 6350  
AGAGTACAAT GGGCTTACCT CGGCTCACTG CAACCTCCGC CTCCCGGGTT 6400  
CAAGTTATTC TCGTGCCTCA GTCTCCCAAG TAGCTGGGAC TAGAGGCATG 6450  
CACCACCACG CCTGGCTAAT TGTGTATTTT TAGTAGAGAC AGGGTTTCAC 6500  
CATGTTGGCC AGGCTAATCT CAAACTCCTG AGCTCAGGTG ATATGCCAC 6550  
CTCGGCTCTC CAAAGTGTG GGATTACAGG CGTGAGCCAC TGCACCCGGC 6600  
CTCAGATTAA CTATATTTCA AGCGTTCACT AGCCACATGT AGCTAGTGCT 6650  
ATGGTAGTGG ACAGTACAGA TCTGCATTTT AATTAAGACA CGTATACAAG 6700  
CATAGTTCAC TAATGCACGG TAAAAAAG TATAGTGCTG AGTCGGTGGT 6750  
AGAAATCCTA AATAGTGCAG AGCAAAAGTG GTACGAACAG CAATCTCAGT 6800  
GATAATGCAA CCATGCTTGC TTTTCATTGC AATTGTCTTA TTTTCTTCA 6850  
GCAAGTTTCA TCCATTTTTC CCAATTCAAT AAATATTAC TGATAAAAAC 6900  
TTTCAATATT AGATTCTTGC ATCTTCATAG ACAGAGTTGC TTTTCACATT 6950  
TAGAAAAATTA CTATCAATG TTAACACACG GTTTTGATAA CCAGTGTGG 7000  
AAAGAGGTGC AGACTCCCA TGTGCTTAT GTAGGCAGAA ATATTCCAGT 7050  
CCAAAGGCAA ACAAGGGCT GGGGACAATC ACACACCTCA TGTCTCTAA 7100  
CTCCTGGGAA GTGCTGTCCC TCTGATTGAG CTCTTATTAT TGCCTTCCC 7150  
ACTAACCTTG TCCACTGTGC CCTGGAGCCC TTGTCAGGTG TACCTGTCT 7200  
GTCTCTCTCA CAGAAATATCT CCTTACCTC CTGTCCAAAG CTACAACTTG 7250  
GCTATTCTCT GATGACACTG TCTTCCCTGT AGCCCTTTTG AGTAATGGCT 7300  
GCATATTCTC CCATAGTCCA GTTCTTTTCC TGTCTCCAG TCTGGCTCT 7350  
GGATGACAGC CCACTAGTTT GAACTCCATA CTGCTATAGT TCAAGTCCCT 7400  
TTTGACTTGT TACCTTGGGC AAATACCTC CTTTTGTCA GGTTCCTTGT 7450  
TTGTAATAAG ACGATAATAA TGCCATTGCT TCCAGTGGGT TATTTGAAA 7500  
TTGAGTGAAA GAAGGCGGGT AGCTTCCCTA CACGCTCAGT GTAGACTAGC 7550  
CTGATGTGCA TTACGGGTGA TGCCATGACT CAGTGTGTTT TCCTCATCTC 7600  
CACATCTGGC TCTCATCCAG TGCTCTGCT TACGGCACTC TGTCCCCCTC 7650  
TTACTTACTC CCCCTTATTA ACTGAAGACT GGCAGTATC TCACAGTTT 7700  
CTCTCCACTT CCTAGTCTCA CCATCATCTC AGATGACTTC AAGTCACCTA 7750  
GATAAACTGT CTCAGTTTCT TCACTCACAT TTTTATATA CAGATAATGT 7800  
TACACTCAAG TTGTAACAGA ACCAGCTTAT CCAGCTCATG AAATGTATGC 7850  
ATTTCATCTC AACTCTGTAT TCAGTGACAT CCTGTGGGTA TCTGGAAATC 7900  
AGCCATGGTG AGAATATTTA CCATGGAAAT TGGCAAAATC TAAAAAGCAG 7950  
AGCAGCTTTT TTTCTGAGAG CCAGACCATA GCTCTTCTAC TCCATAGCAC 8000  
CCATCATAAC AATTTTAAA TACCTCCACT GAACAGCTTC TTCTCTCTC 8050  
TACTTCTTCC ATATCTGATT TGAGCTTCTT AATTATCAT GTGAACCACT 8100  
CTTGTAAATA TAACCCCAA TCCCTGTTC AITGTTCTTC CTGCTAAAAT 8150  
ACTAAACCTG GTTTAGTCCA ACCATATTTT CTCTCTTGG AATCTACAGG 8200  
GTGGCCCAA AACTTGGAAA TGGAAAAATA TTACTTATTA ATTTAAATGT 8250  
ATATTAATAA GCCATTTTAA TGCTTCATTT CCAGTCTCAG TGGCCACCT 8300  
GTATAGCTGG GCTATTGAGC TCTTGCGGGA GGAGGGAGTG GACAGTCTCC 8350  
CAGCCACACA GACTAGTGT GCACCAACA TTTTATAGCT TCCAGACTTC 8400  
CCTGGCCCTT AGTGTATCCC TTAACCTCC ATTTCTCTGC CTTCACATT 8450  
CTCTACTTTT TAAAAATCTC TGACTCCACC TTCACCTTAT CATTCCTAGC 8500  
ACATGACCAT ACTTCTGCTT CCCAAAGAAA ATGAGCAATT ACTTCTTTT 8550  
CCTTTTCTCT CTGTCATCAA ATCTGCAGAC ATGTCATGCC TAAGTCCAGC 8600  
TTTCTCTCTT TCTCTGATCT CAGTCTGCTT CTCCATTTT TGCCCTGAAT 8650  
CCGCTCCCTT CCCCAACCCC CAAGGACTTC GCTCTATCAG TCACCTCTTC 8700  
CCTCTCTCTG ATCTTCAACT CCTCCCATTT TACTGGCTTC TTCTCAAGC 8750  
CTTTCCCAA GCTTTTCCCA TCTCAATTAC CTCCTCGCAC ATGCTCTGTC 8800  
AGAAACCAAC CCGTTTCTTC CCTCCCTCG GCAGCCTGTT CTCTCTGTT 8850  
TGCCCTCATG ATGGCACCAT CATGTGTCA CTAAATCAA TCTCTCCGAC 8900  
ATCATCAATG GCCTTCTTT GTTGGGAAAC CTAATAAACA CTTTATCTTA 8950  
TTTGGTCTTT GTTATGGGTT GAATGAGGTT ACCCCGAAAT CCATATTAGA 9000  
AGTCTTAACC CCCAGTACCT CAGAATGTGA CTTTATTGG GAATAGGGTC 9050  
ATTGCAGACG TTATTAGTTA GGATGAGGTC ATACTGGAAT GTGATGGGCT 9100  
GCTTATCTAA TATGACTGAT GTCTTATAA CAAGGAGAAA TTTGGAGACA 9150  
GACACGCACA TAGGGAGAAT ACCATGTGAT GACAGGAGTT ATGGAGTTGG 9200  
AGTCAAAAGC CTATGGGAAC TTAGGAGAAA GACCTGGAAC AAATCCTTTC 9250  
CTGCGCTAG AGAGGGAGTA TGGCCCTGCC ACTACCTTGA ATTCAACGTT 9300  
TCGGCTTTTC AAAACTGTAA GACAATACAT TTCTGTTGTT CAAACCAATT 9350  
AGTTTGCAGT ACTCTGCGAC TGCAGCCCTA ACAAACTAAT ACAGTCTCTT 9400  
GGAGGCAATT GGAAGGTTG ACAATGGAAG CACTTCTTCA CCCCTTTAGG 9450  
TCTGTGCGCT TTCTTGTGG GGGGTGTTTT CTAACAAATC CTCTCCATCT 9500  
CTCTCTCTCT AGTTTGTCTT AAACATTGGT GTTCTTCAGA CTTCTGACCT 9550  
AGGCCTTCTT TTCACCTTAC ATATTCCCT GGGTGGTCTC ACCCACTTCC 9600  
AGAAATTAAT TAAATTAAGT CTATGACAGT ACTGTGCTGG AAACGTGTTA 9650  
ACAACTGGCT CTCTGGGAAG AGGGGAGACT GGTGATGGT TTTTGTGAT 9700  
TTCTGTGGTG TAAATACTCC CTCCATGGCC AATCCAAAC TGCCAACAGT 9750  
TTAAACAATG GCTCACAAA TTTCTCCAAA TTTAACATTT GGCTTTCACA 9800  
GGCCAACAAC GTGGTACAGC CAACTCCAGC ACACCTCTGC TTTTGTGTCA 9850  
GAGAGAAGTA ACTTATTTTT GTACAAAAGG TAAAAAATAA ACACCTGCAG 9900

GCCCCCTTTT TTCTCTTAAC AAAGTCTCT AGAAATAGAA TAGCTGAAGC 9950  
 TTCTTTTATG CATTTCATCTG TTATTTCCAT GTCACGTGGG TGGTGGGATT 10000  
 ATTTTTCCCTT TATTTTCTT GTATATGGTT GAAATACTGT ACCTTTTGATC 10050  
 AGTTTTAGTT TTATGGCATG TTTTGCAACC ATATTAAATC TAGTTTTGTG 10100  
 CAGAGGGCGT CAATATTATT TTCTCAAAAC AAGAAAAATAT TTCATTGCAA 10150  
 AGGAGACAAA CAAAAAGGTC CTAAATACCA AAACCTTGAA ATGTGATTTC 10200  
 TTGTACTTGG CAGTGTCCAA GTGGTAAACC CAAACAGTAT TGGGTTTTCA 10250  
 TTTTGTTCAG GAAAGTCTTT GTCTGGCAGC GACTTACCCT TACATCAGGC 10300  
 GGGCCTTGCT CATTTCATTCA CTAAAGTATT TATTAAACAC CAGCGGTGTG 10350  
 CCAAGTACTT ATCTAGGTAT CGGGTAGATT CTGATAAGTC AGTCAGGTCC 10400  
 CTGCTCTCAG GGAGCTTGCA GCAGAGATGG GGGCTGCAAT AGAGAGTAAG 10450  
 CCAAGGAAAT GAAAAAGGAA GTTGATTTCAG GAGAGTGATG AATGCTATGA 10500  
 AGAAAAATGAA GGCAGCGCAG TGTGATGGAG AGTGACCCAA GGTGGTACAG 10550  
 TTTGTACCTC TAAGGACCAG ACTGTGACCC AGGTCACTCA CAGATGCCCG 10600  
 TCATGTGATG CCACAGCAAC TTTTCCAGGT GCTCGTTTCC TCCCCTTCC 10650  
 CAGTCTCTTG CCCAGCCGCG ACTGCTTACA AATACAGCTA GAGGAATCTA 10700  
 AATGAGGTTT CTCTATCATC AAACCCAATC AAAATGCCAA GGAACAGAAT 10750  
 CAGTGCCTGG CTGAAGGCAG TGGAAACAGG CCAGCCTGGA GTGGTTCTCT 10800  
 CTGAGGAAGT TCCTCATCTT GGTTTTAGGG CCATACCTTG TGACCTGTGA 10850  
 GCTAGGGGTT GCCAGTCCCT GACATTTCTA CTGAGGACTC GCCTGTCTAT 10900  
 ATTCCCGGCC TGTATGTGTC TCCTGAGTTC CAGACACACA GGGCGAAGCG 10950  
 CCTGATGGAT GGAAGTATGT TTTTGGTGT TCCATTGGTA TCTCAAATTC 11000  
 TACAAAACCT AGTGCCCTCT CTCTCCCTG TTCTCCCTCA TCTTCAGTCT 11050  
 ATCAGCTGTT CCTCATCCAG CAAATGATAT TACCATCTTC CAAGGAGCTT 11100  
 CCCAGGAGTA ATCCTTGACT CCTCCTCAAC ATCCAATTAA TAATCAAATC 11150  
 TAGGGCAGGT ACAATAGCTC ACGCCTATAA TCCAGCACTT TGGGAGGCT 11200  
 GAGGAGGAGT GATCATTGGA GGCCAGGAGT TCAAGACCAG CTTGGCCAAC 11250  
 AAGGTGAAC CTGTCTCATT TAAAAAAGT TATTTTAAAA ACTCAAATCT 11300  
 ATTAATTCTA CCTCTAAGTG TGTCTTGAAT TTATCCATCT CTCTCCATCT 11350  
 CTGAGCTGTT ACCTTACCTC AGTCCATCAC GTTTTGTCTA CGTTAACATG 11400  
 ACCAGAGTCT TGTCTTAGT CTGGTGAGGT CACTCCAGCT GCTTCAGATC 11450  
 CTTCCATGGC TCACCGTTGC CCTCATATAA AGTTGGCACT CTTGGACATG 11500  
 TGGCTTACGG GGCCTCCCT GATGTGGCCC TATTTGCTTC TCCATTCTGT 11550  
 TCTCTCCAG CCTCTCTGCC CCCATCTCTA GGCACCAACC ACACCTTCT 11600  
 GCTCGTCAAT GGTGCCAGCT TCTCTCTAT CTCTGGTCTT TGGACAGACT 11650  
 TTTCCCTTCA CCTGGAATGC TTTCTTCAAT CCTACCCAC TCTCTTAAAT 11700  
 CTAGATAAGG TTTATTCTTT TTGAATGTCT AGCAGTGAAA CCATTTCCCC 11750  
 TGAAAAACCT TCTCTAACCA ACCCCCTACC CTCAGCCCAA GGTCTAGATT 11800  
 AGGAGTCCCT CTGAATGTTT CCATAGCATT TTTAAAGAAAT TGCCCTATTTA 11850  
 TCTGTTCCTA TCTATACCTA AACTACAAAT TGTATGAGAA CAGCCACTAT 11900  
 CTCTGCCTGG TTACCAATTC ATCTCCAGCA ACTAGCATAA TGCCTGGCAG 11950  
 AGTCAGCCTG CAACAAATAT TTGTTGAATA AATTAACAGA TGGCTTTATC 12000  
 TCCTTAAGTA AATCTTGCTT TTTTCACCTA TTAACACAGA CGCACAGGCC 12050  
 AGGTGTGGTG GCCCATGCCCT GTAATCCAG CACTTTGGCA GGCTGAGGTG 12100  
 GGCGGATCAC CTGAGGTCAG GAGTTCAAGA CCAGCCTGGC CAACATGGTG 12150  
 AAACCCCATC TCTAATAAAA ATACAAAAAT TAGCTGGGCA TGGTGGTGGG 12200  
 TGGGTATAGT CCCAGCTACT AGGGAGGCTG AGGCAAGAGA ATCGCTTGAA 12250  
 CCCAGGAGGC AGAGGTGGCA GTGAGCCGAG ATCATGCCAC TGTACTCCAG 12300  
 CCTGGATGAC AGAGACCCCTG TCTCAAAACA CACACACACA CACACACACA 12350  
 CACACACACA CACACACACA CACACACACC AAGTTGTATA ATTTAAAAATA 12400  
 TAACGTGCTT GTTATGGAAC ACTTGTAAAA TACAGGAAAG TAATGAAAAA 12450  
 GTCTACCATC TAGCTCACCA CATAATGACC ATGTCTATCA TCCTGGCATA 12500  
 ATTCTCTCCT GTATATAAAT ATATATTCTT TTATTGTTAA AATTACACTA 12550  
 TGAGTACTAT TTATTATTAT TACTGTGGCA AAATGCGCAA AACATAAAAT 12600  
 CTTGCCATTT TAAGGTATGC AGTTTGGTGC ATTCACCACA CTCACATTGT 12650  
 TGTGCAATA TCACCACTAT CTATCTCAGA ACTTCTTGT CTTCCTCAAC 12700  
 TGAAACTCTG TACCCATTAA ACAATAGTGC ATCTCTGTT TTCCCTCCCT 12750  
 TACAATTAT TTTTATTGG GTTTGTACCA AACTGAAAAAT AGCTGCTTCT 12800  
 TCCTTACTTA GTTCAGATTA GCATTTCCAT TTATTAGCC GTGGTTTTGA 12850  
 GGATGCCATG ACAGATGCCA TCCTTCCTAG AGCTCTTTGG GGCCTGTCAGG 12900  
 TATTTAGCTC AGGGTGAATT CGGGTTGATA ACATTTTAAA ATCTCACTTT 12950  
 ATTCTGAGGT TCCTAGTGTG AGAGCCCACT GTATTTTATG GGAATCCCAA 13000  
 GTTACAAACA AAAATATGGT GAGGAGGAAT CACTGAAGTT TTAACACAAG 13050  
 AGACTTACAT TTTGTTCAAT TTCTATCTTT TAGTTTATTT CCTAAGCATA 13100  
 AAGAAATACT TTGAAATTTT TACATAGCAT TATACATATT TAATTAAAGCA 13150  
 TGAGCACATC TTAACACTTT AAATTTTGA TCAGATCTTT AATTCCTAGG 13200  
 ATATTAAAGG GTACTGGCAA TTTGGCCAGG TGTGGTGGTT CACGCCTATA 13250  
 ATCCCAACAC TTTGGGAGGG TGAAGTGGGC GAATTGCTAG AGCCAGGAG 13300  
 GTGGAGGCTG CAATGGCCTG AGATCAGGCC ATCGTACTCC AGCCTGGATG 13350  
 ATGAGAATGA AATCTGTCT CAAAAAATAA AAAAAAATAA AAAAGAAGAA 13400  
 GAAGAAGTAT TGGCAATCAG TGCTCCAGGA ATAATTCTCT GACTTGAAT 13450  
 AAACCTACAT GTAGACAAAC TAATTAGGCC ATTCCAAGAG TTGCTAGCAT 13500  
 TGGTTTAAAT TGTTTTCAGA GCATTCAGG AAGCAGTGTG GCCAGCATTG 13550  
 CATGTTTGTG ACTTCAGAAA TGTATGACAG GTGTTTCTCT TACCAGGTC 13600  
 TTCTGTTTTC TTAGTTTTCG TCATGTAAT ATTTATGAAC ATCCTCATCT 13650

TTTTGAGGGA	AGGGATTATA	GATCATTCTA	ATTCCATTTT	CTAGCATTTG	13700
GTACCATTCT	AAGCACATGA	TAGGCACCCA	TTTGGAGCAT	TTTTGGCTTG	13750
ACAGAAATATG	CATTTAGAAAT	TGTTCAAATT	AGAGGTGTCA	GTGATGGGAA	13800
TTAGAATACT	ATATAATTCT	AAGTCATTTG	ACTTAAATAC	AAAAGAATGA	13850
TTTTCTCTGG	TGGGGAATGG	TGAAGGGAGG	CAGGAGTTAA	GAAGAGGAGA	13900
AGAGATCCTA	AGTCATTTAT	AAACTTCTCT	GGAAAGACAG	GTGTGTGAAG	13950
ACTTTTTTAA	AAGTCATTCA	CCAAATTGTG	TGTGTGTGTG	TGTGTGTGTT	14000
TTAAATAGAC	TTTATTTTTT	AGAGCAGTTT	TAGGTTTACA	GCAAAATTGA	14050
ATGCAAGGAC	AGAGATTTCC	CATAAACCCC	CTGCCACAC	ACATGCATAG	14100
CCTCCCTCAT	TATCAACATC	CCCACCAGAG	AGGTGTTTGT	TCTAGTTGAT	14150
GAACCTACAC	TGACACATCA	TTATCACCCA	AAGTCCATAG	TTACGGGCAG	14200
GGTTCACGTG	CGGTGTACAT	TCTATGGGTT	TGAGCAAATG	TATAATGACA	14250
TGTATCCACC	ATTATAGTAA	CATACAGAGT	ATTTTCAGTG	CCCTGCAAAT	14300
CCCCTGTTCT	CCACCTATTC	ATCCCTCCCT	CTCTGCATTI	CCACCCCCAG	14350
CCCCTGGTAA	CCGCTGATCT	TTTACTGTCT	CCATAGTTTC	GGACGATCTA	14400
TTTTTCAGAC	AGACACAGAG	CTGTCTTTCC	CTTAGTTTCT	ATTCTATCAT	14450
TTCTTTCTCC	CCATCCATCA	TAAAAGGCTA	TGAGTTTTTT	TTAAGTGTGG	14500
AACACCATCC	TACTTGTCAA	GTTAAAACAT	AAGCTCCTGG	CTGGGTACAG	14550
TGGCTCAATG	CTGTAATCTC	AGCATTTTGG	GAGGCTGTGG	CAGAAGCATC	14600
ACTTGAAGCC	AGAAGTTTGA	GACCAGCCTG	GGCAACATAG	CAAGACCCCA	14650
TCCCTCCACA	CACAAAACACA	CACACACACA	CACACACACA	CACACACACA	14700
CACACACACA	CACAAAACACA	AGCTCTTGCC	AGAATTAGAG	CTACAAATTG	14750
CCCTCAGGTT	CCTAGAAGAT	CAGTCCCTCA	ATTAGATTCA	GATTGAGATG	14800
CTTCTCTTTT	TAAACAATGA	TTCCCTTTCT	ATCATGCCCA	ATAAGAAAAC	14850
AAATAAAAAAT	TAAACAATAC	TGCCTGTAAT	CTCAGCTACC	CAGGAGGCAG	14900
AAGCAGAACT	GCTTCAACCC	GGCAAGCAGA	AGTTGCAGTG	AAGTGAGATC	14950
GCGCCACTGC	ACTCCAGCCT	GGGAAACAGA	GCAAGATTCT	GTCTCAAAAA	15000
CAAAACAATG	TGATTTCTCT	CTCTAAGTCC	TGCACAGGGA	AATGTTAAGA	15050
AATAGGTCCA	CCAGGAAAGA	AGGAAGTAAG	AATGTTTGAC	TAGATTGTCT	15100
TGGAAAAAAT	AGTTTATACT	TCTTGCTTGT	CTTCTTAACA	GTTCTCCAAA	15150
GCTTCGTACC	TTGGCCAGAG	GCTTGTCTCC	TGCGTACCTG	AGGTTTGGTG	15200
GCACCAAGAC	AGACTTCCTA	ATTTTCGATC	CCAAGAAGGA	ATCAACCTTT	15250
GAAGAGAGAA	GTTACTGGCA	ATCTCAAGTC	AACCAGGGTG	AAAATTTTTA	15300
AAGATTCAC	CTATATTTTA	ATTAACGTCA	GTCCGTCATG	AGAATGCTTT	15350
GAGAAAAACT	TTATTCTCA	CACCTAACAA	TAAATGAGAT	TAACCTTCCTC	15400
TCCCTCATC	TGACCTGTGG	AGGAATCTGA	ACAAGAGGAG	GAGGCAAGTG	15450
GCAGGTTTCC	TTATCATGAT	GTTTGTCTATG	TTCAGTGTGA	GGCCTCACAA	15500
AAAAAATAA	AAAAAATAA	GGCGTCCTGG	ATATAACTGA	GAGCTCATTTG	15550
TACAGTAAAT	ATTAATAAAA	CAGTGATTGT	AGCTGAAGGA	TAGAAGTCTG	15600
TGGAGGGAGC	AAGTGGGTAG	AATCGCGTCA	AACTAAAGAG	CATTCTAGC	15650
CAAAGACACA	ATGATAGATT	GAAGGATATT	TATTCTAAAT	ATAGAATATG	15700
GGTGAACGAG	ATCTGTGGAC	TTCTGGGCTC	CAACGTTAGA	TTCTGATTTT	15750
AGCAAGCTTG	TCAGGGGATT	CTGATATTGA	AAGGCTGTGG	CCTTCACCTG	15800
AGAAACCTGC	CCTAGGGGGC	CATGAAAAAT	TGTCCTGTCT	TTCAGAAGTG	15850
CTATCAGACA	TCAAAATGGA	GTTAAATCGT	ATCTTAACAA	TTACTAGGAT	15900
GGGCGCAGTG	ACTCACACCT	GTAATCCCAA	CACCTTGGGA	GGCTGAGGCA	15950
GGAGGATCAC	TTGAGCCCAG	GAGTTCGGGA	CCAGCCTGGG	CAACATAGAG	16000
AGAGGTTGTC	TCTATTTTTT	AATAATTTAA	AGAGAAAAAA	ATACTGAAAA	16050
TATTGTATAC	ACCACCTGAAT	TATAATAATG	TGTATATAAT	GTATATATTC	16100
ATTATGAGGA	ATATTTGATT	ATTTCATATA	TTATATCTTT	TCCTTCTGTT	16150
TATTTTATCC	AGTTATGAAG	TATTTAGAAC	AATTCATCAG	TAAATGGGGC	16200
TAAATTGACA	GAATAGTAAT	CAGAGAAAAAT	AGAAAAAGAC	AGATGGGTTA	16250
TCTTTGAATA	CCAGGTTGGA	GTTGTTTATG	GGTTTGTGTT	TTGTTTGGG	16300
GGCGTTTTTT	TAGACAGAGT	CCCACTCTGT	TGCCAGGCT	GGAGTGCACT	16350
GGCACAAGCA	TGGCCCACTG	CATCCTTGAC	CTCTTGGGCT	CAAGCAATCT	16400
TCCACCTTA	GCCTCCTGAG	TAGCTGGGAC	CACAGGTGCA	TGTCACCACA	16450
CCCAGCTAAT	TTTTTTATTT	TTTGTAGAGA	CAGTCTTTCT	ATGTTATCCA	16500
GGCTGATCTC	AAACTCCTGC	ACTCAAGTGA	TCCCCCTGCC	TTGGCGTCCC	16550
AAAGTATTGG	GATTATAGGC	ATAGCCACCA	CACCCAACTT	AGTTTCTATT	16600
TAGACTTGGC	CCTTTCCAC	CAGTCATTTG	TGTCCAAAAG	ATCTCATAAA	16650
TGTAGACAGG	AAACTGTCCCT	TTGCTCATCA	GTTTTCTTCA	TCCTGTGTCT	16700
AGGGGGATGG	TCGGTGGGGG	AAACTGGGGT	TATGCAAGTI	CCTCTGAAAC	16750
ATCCTCTGTG	AGCCCAAGGA	TGGATGAGGC	ACCAGCCGCC	AGCGAGTCAG	16800
TGTGCAGCTT	TCCAGAAAGG	AAGTCATCAG	CCAGTCAGCC	GGCCCTGGCA	16850
GCCAGCAGCC	GGCAACCCCTG	CTGTCTTGTG	ATAAAGAAAT	GGTCTGCCTG	16900
ACAGGATGGT	GTGGATTTTT	CTTTTTTCTT	TTTTTTTTTT	TGAGACAGG	16950
GTCTGGCTCT	GTGCGCCAGG	CTGGAGTGCA	ATGGCGGGAT	CTTGGCTCAC	17000
TGCAGCCTCT	GCCTCCAGG	CTCAAGGCAT	CCTCCACCTT	CGGTCTCCCG	17050
AGTAGCTGGG	ACCACAGGCA	CACACCACCA	CGCCCAACTA	AGTTTTCGTA	17100
TTTTTAGTAG	AGGCAGGGTT	TTACTATGTT	GTCCAGGCTA	GTCTCAAAT	17150
CCTGAGCTCA	AGCTATCCAT	CTGCCCTTGG	CTCCCAAAGA	GCTGGAATTA	17200
CAAGCGTGAG	CCACTGTGCC	TGACCAGGGT	GGATTTTTTC	AAGTGCACAT	17250
GTTGTGTGCC	CAGAAGCTCT	GATGGTACCA	AATTCCAAGC	GAAAAAAGT	17300
CAATGGTTCC	CACCCATCCT	ACCTCCCATG	ATGGCAAGAG	GAAATCACCA	17350
CACGTCAGAT	ACAGTCCATG	TAAAACAAT	TGCTATGGAT	TTTGAAAGTG	17400

AACCTTAAGA	GAAGTGCAGT	ATGTTTTCTT	CATTAGAGTT	CTCTGGTAAT	17450
TTCCAGCTTT	TTTTTTTTTT	TTTTTTAGAC	AGTGTCTCGC	TTTGTGCGCC	17500
AGTGTCAACC	AGGCTGGAGT	GCAGTGACGT	GATCTCGGCT	CAGTCAACC	17550
TCGCGCTCGT	GGGTTGAAGT	GATTCTCCTG	CCTCAGCCTC	CTGAGTAGCT	17600
GTATTTTAGT	AGAGACGAGG	TTTCAACATT	TGGCCAGGCT	GGTCTCGAAC	17650
TCCTGACCTC	AAGTAGATTG	CCCATCTCAG	CCTCCCAAAG	TGCTGGGATT	17700
ACAGGTGTGA	GCCACTGCAC	CCGGCCAGTA	ATTCAAGCT	TCTGAGGAGC	17750
CCTTTGAATT	GTTAAATAAC	TTGTAGCTAT	GTCCAACATA	TCCATGTTCA	17800
GTGTATGTTT	GATATTTCTT	AGGAAACCTG	CCCTTGGTTG	TTTTCTTTGT	17850
GGTAATTCAT	GAGCCGGCAA	ATTTGACATG	TGTTACAGAA	TATACCTTTT	17900
CTCTGCTCTC	CTACCTCATA	ACCAGAACTT	AATTATCCTG	CTTTAGTCAC	17950
ATAAATAGCT	AACATAATAA	ATATATGAGA	TTTCAGTCTG	CTCACTGTGA	18000
AAATAGACCT	TCTAAATGAT	CTCTTCCACT	TGCAGATATT	TGCAAAATATG	18050
GATCCATCCC	TCCTGATGTG	GAGGAGAAGT	TACGGTTGGA	ATGGCCCTAC	18100
CAGGAGCAAT	TGCTACTCCG	AGAACACTAC	CAGAAAAAGT	TCAAGAACAG	18150
CACCTACTCT	AGTAAGAAAT	GAAAGGCACC	CTAGAGATGT	TCCAGCCCCA	18200
AAGATATTTG	AATAGGTTGG	ACTCGGGCAC	CAATCTAGCA	AGTCTACGG	18250
AAGTTGTATA	AAGCTGAAAA	TACTGAAGCA	TTTCCCAAT	GGGAAATCCT	18300
AAACTCAAAA	CTTGCTTTTT	GGTTTTTTG	TTTGTGTTGT	TTTTCTTCAT	18350
CTGACATTGC	TTAGTAGTCA	CAGAATGAAA	GATAAATCAA	TCATTTCATGA	18400
TCTAACAAATG	ACCTTCAGTG	CTCTAAAAAA	CTACGGAGTC	AAGGAAAAACA	18450
TGAATATATT	CCTCATGTAA	AATTAATAA	CAGACATATA	AAGGGCAAAA	18500
CATGAACATC	ATTTCATACCT	TGAGGTCCGT	CCCCCTCCCA	GAATAAACCC	18550
CCAGTATGCC	TTGGTTTAGA	GCATTAAAGCA	GGAGGGCCCT	GAGTCACTCC	18600
AGACAGTCTT	GACCACCAAG	CAGCATCTCT	TTTTTGTGTT	CTCTGTGGCT	18650
TTTGCAAAACA	CAGGGCTAGC	TCAGCTACCC	ATTAGTATGT	TTTCAGTCAC	18700
TAAACAGTCC	TTCCAGTCTT	CAAAATTAGGA	TGACATTGTC	ACATGGGGCT	18750
TTAAAGCAAG	TGAACAAGG	AACCCCTTTT	TTTTTTTTTT	TTGAGATGGA	18800
ATCTCACTCT	TGTCGCCGAG	CCTGGAGTGC	AATGGCGCAA	TCTTGGCTCA	18850
CTGCAACCTC	CACCTCCGAG	GTTCAAGAGA	TCTCCTGCC	TTAGCCTCCT	18900
ATTCAATTATG	AGGAATATTT	GATTATTCAG	TTCTGTAGG	GTAAAGATAT	18950
TACCCCGATG	CATATTATTG	ATTATTGAGT	AGCTGAGATT	ACAGGTGCCT	19000
GCCACCACGA	CCGGCTAATT	TTTTGTATTT	TTTAGTAGAG	ACAGGGTTTC	19050
ACCATGTTGG	CCAGGCTCCA	GGCTCGTCTC	GAATCCTGTA	CCTCAGGTGA	19100
TCCACCCACC	TCAGCCTCCC	AAAGTTCTGG	GATTACAGGC	GTGAGCCACC	19150
ACTCCTGGCC	ACAATCCTTT	TTTAACTATG	AAATATATTT	TTATCTGAAG	19200
TTTGAATGTT	ATACCCAACT	GAGGGATGAT	GTTCCCATAT	CTCAGTTAAA	19250
GAAATAACCT	GCTCAGATAC	TTCAAGCTCT	TCTTTGACT	TTTGAAAATA	19300
AATGATCTTG	AAGTTACTAT	ACTTTGTTTG	GGTTAGTTAA	CATTATTTAA	19350
AGTATATTAT	TTTAAATTAAT	TATCTTTGTA	AGATTTTACT	GTATACTACC	19400
TGGAGTTCAA	TGTATCAGAT	GGATTTCAAA	TTTATGTACA	TTTTTTATGT	19450
ATATGTTACA	GAAAAAATG	TGATCCATAA	GAAATCAGAA	AATAGCGCAT	19500
ATGCTAATAG	CTAATGTTGT	CCTCTAAAAA	ACTTATTTTT	GCATTTTTAA	19550
GAGGGGGATA	TACTCTGACA	CTTTAATAAG	TGTAATTAAT	TATTGACTGG	19600
AATTTGGCAT	GAGGCAGGGC	CATTTCAGAT	CCCATTAAG	GAATGACACA	19650
TACCAGAGAA	CCACAGAAAT	AAGGCCACAT	TTGTAATAAA	TCATTATAGC	19700
TCTGTAGGTA	GAAGACCCAG	TTGTATTAGG	TAATTAATGG	ATTGTCTCTT	19750
AAAACACATG	TCCCGGAAGA	TATAGGTGAG	TCTTGGGGGG	CCGCATTAAA	19800
CATTATACCA	ATGTATCTTA	CATTTCCTAG	AAAGTTTAC	TACTTTACAG	19850
GATCTTTCTG	TTACCAAAAT	GGAAGGTTTC	CAACTCCAGG	ACTTGGCTTT	19900
CATAGTTCTT	ACACCAGGGG	AAATGCCTTC	CTTTGCTAAC	TATGCAACCA	19950
GGTTAGTTAG	TGTAAGTCCA	GCCACCCTGT	TGGCAATGCT	AAAAGGTACA	20000
ACAAACACAG	AATTTTATTT	GCATTTGTAA	ACATTTGATT	TCTGGCTCGA	20050
AATTTTCAGT	TTTCATGGGC	ACGTCAATGA	AACAGAAATC	TTCTGTGTTT	20100
AGTTTGGGCA	CCTACTCATT	GTAGTGACAA	ATATTTGAGA	AGCCAATAGG	20150
GGATTCCACA	AATGTTCTG	AACCTGTGGC	TGAGACTGGT	AATGGCTGAG	20200
TGACATGGGG	ACATACCACA	AAAGAAGAGG	TAGCAAAAGG	CTGCTGAGAT	20250
AAGGACATGT	TCATTGCTTA	GCTAGTGGCC	TGCACCTTA	AAACACATGT	20300
CCCAGGCTGG	GTGCTGTGGC	TCACGCCTGT	AATCCAGCA	CTTTGGGAGG	20350
CTGAGGCGGG	TGGATTACCT	GAGGTCAGGA	GTTGAGACC	AACCTGGCCA	20400
ACATAGTGAA	ACCTCATTTT	TACTAAAAAT	ACAAAAATTA	GCCAGGCATG	20450
CTGGCGGGGG	CCTGTAGTCC	CAGCTACTCA	GGAGGCAGGC	AGGAGAATTA	20500
CTTGAATCTG	GGAGGCAGAG	GTTGTGGTGA	GCCGAGATTG	CGCCACCGCA	20550
CGCTAGCCTG	GGCGACAAAG	TGAGACTCTG	TCTCAAAAAA	ACAAAAACAA	20600
AAAAACAAACA	AACAAAAAAC	AACAACAACA	AAAAAACGGG	TATCCAGAAA	20650
GATACAGGTA	AGTTTTCTAA	CACAGGTCCT	CTTGTATGGT	GCGTTCCACT	20700
TAACTAGAAG	ATGACAAAAA	CATTTGTCTAT	GAGAAATATAG	ACTCACATTT	20750
TAAACCTGTT	TGAGCAGGAA	AAGGAAGCAA	TGTTACAGAT	GTAATTCCTG	20800
GTGTGACTGC	AGAAAGGATG	ACTCCCTTAT	TAAAGTAGTC	ATCCTGAGTG	20850
AGCTAACTCT	TTGTACTTCC	TCTTCTCCTC	CTGTTCCCTT	CATCACCCCA	20900
TTCTTCCGTT	GCCTACACCC	AGGCCACAT	TGGATGCTGA	CATAGACTTA	20950
CATGGTACAG	TCCAAGGGAA	AGATCTGCCA	TTTTTTTCAA	TGTGTCATCT	21000
TGTTTATCTT	CATTCCAAGG	ATCTCTCCAC	TCTTTATACA	GTAAGAGATG	21050
AGAGTCTGGA	AAGGATTGGG	AATAAGATAA	TGAATTGTAA	GTTTTAAATT	21100
GTTCTTCGTA	TTTTGGGGAA	GGAGTAGGCT	AGGTGGTCCT	TCTGTTTTTT	21150

TTTTGTTTTT	TTTTTTAAAG	TAGATGTGGC	CAGACGTGGT	GGCTCACGCC	21200
TGTAATCCCA	GCACCTTGAG	AGGCTGAGGC	AGGTGGATCA	CTTGATGTCA	21250
GGAGTTCAAG	ACCAGCCTGG	CCAACACAGT	GAAACCCCGT	CTTTACTAAA	21300
AATACAAAAA	CTAGCCGGGC	TTGGTGGCGT	CCACCTGTAG	TCCCAGCTAC	21350
TGCAGAGGTG	GAGGCAGGAG	AATCACTTGA	ACCCGGGAGG	TGGAGGTTGC	21400
AGTGAGCCAA	GATCATGCCA	TTGTACTCCA	GCCTGGGCGA	CAGAACAATA	21450
CTCTGTCTCA	AAAAAAAAGA	GAAAAAGAAA	GAAAAAAGA	ATGGATTGTA	21500
ACTCAGTCGT	CAATAGCCTC	TATTCAGGA	GATGTTACAG	TTGATTATGT	21550
TATAGGGGGT	GTATAATAGA	ATTTCGAGCT	ATGTAATTC	CAAGTGCATT	21600
TGGAAGAATG	AAGAAATGGA	GGAAGGGTAA	AGTATGAGTG	CAAGCATTCC	21650
AGGTTTTTTG	AAAAATGCTAT	AATCTTTGTT	CAGGGCTAGT	ACAAAGTGCT	21700
ATTTAGCTGT	AAGGGTTTTT	TGTGATTAC	AGACAGTTTT	CACATGTGTC	21750
ATTTCAACCT	TGGTTTTATG	GCGAAGGCAT	GTGATGGTGC	TTGTCCCGAG	21800
ACTTTAGATC	CATATCTGAG	GTTCTGTGCG	GGCAAAGATA	TTACCCCTGA	21850
TCATATTATA	GTCTATAAGT	GGGAGAGTTG	TGCCTGGAGC	TCAAGTCTTA	21900
TGATTTCCTGA	TCCAGGGCAC	TTCTACAAC	ATGATTTTGC	AATATAAAAG	21950
CCTATAATGT	GTGACTAAAG	CAGGTCACTC	ACCCCTTGTA	ACAGACTCTA	22000
GTAATGGTAC	TGCCACCAAA	CGGCTGCGTG	ATATTGGGGA	AAGACTTACC	22050
TTATTTGAAT	CTCAGTTTCC	TCCTAGAAAA	ATGAGGGTGG	AGGTTAAGCA	22100
TAGGCTGATG	ATCCTAAAGC	CTCCATACTG	CCCTAAACTG	TGGCTCTAAG	22150
ATCCAGTAGA	ATGCTGGGTC	ACAGGACTCT	AGGGAGCTTT	TCAAACCCAA	22200
ATGCTGTGCA	TTCTTTGATG	GTAGGCAGCA	GTTTATGGAA	GTGGGCGACA	22250
CAGCAAAATAT	CAAAATACCT	AAAGCAGCTT	GCAAGAGTTG	TTTCTGCCTA	22300
GTGGTCTTTA	TAGTTAATAT	TAAATAGTTA	ATTTTTTTTT	TTTTTGAGAC	22350
AGAGTCTTGC	TCTGTTACCC	AGGCTGCAGT	GCAGTGGCAC	AATCTCGGCT	22400
CAGTGCACCC	TCCACCTCCC	GGGTTTGAGC	AATTCGTGCT	CAGCCTCCCA	22450
AGTAGCTGGG	ACTACAGGTG	CATGCCACTG	CACCCAGCTA	ATTTTTGTAT	22500
TTTTAGTAGA	GACGGGGTTT	CACCATATTG	GGCAGGCTGG	TCTCGAACTC	22550
TTGACCTCAG	GTGATCCACC	TGCTTCAGCC	TCCCAAAGTG	CTGGGATTAC	22600
AGGCATGAGC	CAGTGCACCC	AGCTTAAATA	GCTAATATTT	AATATTATTC	22650
TATAGTTATT	CAAGTAATTC	AGGCCAAAGA	CTTAGAAACA	AAACAAAAG	22700
CCACTTTTAA	GAGAAAGGG	TGTAAGTTTG	CCAGATAGAT	AGAGATCTTT	22750
CTTTTTTAAC	TACAAGAGTT	CAGGAATGAA	TTACTCTTTA	ACAAACGACT	22800
ATAGATATAC	ATGAAATTTG	GAAGGACTTA	TTATGCATAT	GATAATCAAT	22850
TTAAAGACAA	CACCTTAAAT	TATATTGTTG	CCACTCTCAA	AAAGTGGTAA	22900
TAGAACAGCT	AATGGTTTAA	AAAGCAGAGT	ACAGAAGTTC	CCAAACTTAT	22950
GGCACCTTAA	TATCGCAGAA	AACTTTTTAA	AGCATGCCTA	GGCCACAAAA	23000
AATACCTGTA	TTTTGATTAT	TAAATTGTAA	GGTCTACACA	ACCTAATAGT	23050
AATAGGTCCA	ATAGTAATGC	TGTCCAATAG	ATGTTGATGT	TTTTTTCCTT	23100
GCAAACCTTA	AAGATCCTAC	AGTGCCTCTG	TAAATAGCAC	TGCTTGTTA	23150
GAGTTGAATT	TCAGATAAAT	AATTTTTTTC	ATGTTAATTA	TTTTTCTTTT	23200
CTTTACTTTT	TTTTTTGTTT	TTTTGTTTTT	TTGTTTTTTT	TTTTTGAGACA	23250
GGGTCTCATT	CTGTTGCCCA	GGCTGCTGTG	CAATGGCATG	ATCATGGCTC	23300
ACTGCAGCCT	TGACCTCCCT	GGGCTCAGGT	GATCCTCCCA	CCTCAGCCTC	23350
CCAAGTAGCT	AGCTGGGACT	ACAGGTGCTT	ACCATCATGC	CCGGCTAATT	23400
TTTGTTTITT	TTGTAGAGAT	GTGGTTTTGC	CATGTTGCC	AGGCTGGTCT	23450
TGAACCTCTG	GGCTCAAGTG	ATCCGCCCGC	CTCGGCTCC	CAAAGTGCTA	23500
GGATGACAGG	CATGAGCCAC	TGCACCTGGC	CCCTGGGCGA	AGTATTCTTT	23550
AATGGTTACA	TAGGACATAC	ACTAAACATT	ATTTATTTGC	TATATGAAGT	23600
TCAGGTTTAA	CTAGGTGCC	TGCACTTTTA	GTGCTAAAT	CCTGTAGCTG	23650
TACCATGCA	TTCACTGGTG	CTCCCCAGCT	TGCTTGCAC	AGAGTTTGGG	23700
AACCATAGTC	CTATAACTCT	AGGCCAATTT	TTTTAATGTA	AATTTGATTC	23750
ATTTTAAATT	AATAAATAAT	AACAGGAATT	TTTTTAAAAA	TTGTTTAAAA	23800
TATAATTAAA	ATTATCAAAA	TATTTTTTAA	CTGAACCTGT	GACTAGAGAT	23850
ATTTAGATT	TGAAGAGTGG	GGTTTATGCT	AACTAATGAC	AGTCTGGCTA	23900
TGCATGTGGA	GCACTGAGCT	ATAAATTGTG	GCTTCCCCAA	TTCTCCTGAT	23950
GTCACTTGAA	CAAAACCTAA	GTGTCAGACC	AGAGCTTCTG	GTATCTTCCA	24000
TGGGATTTC	TTCAACAGCT	GGAGCAAATG	AAGTCAGATT	GATTTTTTTT	24050
AATTTGTCCA	ATTTTGTGTT	CTCAAAAACA	TAATTATAAT	CATTTATTAG	24100
AACTAGAAAT	TCTTCAGTTT	AACAACAGAA	ATAGTTATTC	ATTATGAAAA	24150
GCGAATCTGG	AGGCCCTTCAT	TGTGGTGCCA	ATCTAACCAT	TAAATTGTGA	24200
CGTTTTTCTT	TTAGGAAGCT	CTGTAGATGT	GCTATACACT	TTTGCAAACT	24250
GCTCAGGACT	GGACTTGATC	TTTGGCCTAA	ATGCGTTATT	AAGAACAGCA	24300
GATTTGCACT	GGAACAGTTC	TAATGCTCAG	TTGCTCCTGG	ACTACTGCTC	24350
TTCCAAGGGG	TATAACATTT	CTTGGGAAT	AGGCAATGGT	GAGTACCCCA	24400
GGGAACAATT	CATTAATAAG	GAGATTCCCC	ACTAGCATTA	TTTCTTTTCT	24450
TTTCTTTTTC	TTTTCTTTTT	TTTTTTTTTT	GAGACAGAGT	CTCGCACTGC	24500
TGCCAGGCT	GGAGTGCACT	GGCGCCACCT	CGGCTCACTT	GAAGCTCTGC	24550
CTCCCAAAC	GCCATTCTCC	TGCCTCAGCC	TCCCGAGTAG	CTGGGACTAC	24600
AGGCACCCGC	CACCCGCCCC	GGCTAATTTT	TTTTTTTTTT	TTTTTTTTTT	24650
TTTTTTTGCA	TTTTTAGTAG	AGACGGGGTT	TCACCGTGT	AGCCAGGATG	24700
GTCTTGATCT	CCTGACCTCG	TGATCTGCCC	TCCTCGGCT	CCCAAGTGC	24750
TGGGATTACA	GGCGTGAGCC	ACCAGGCCCG	GCTAGCATTA	TTTCTTATGA	24800
CACTTTTTTT	TTTTTTTTTG	GACGGAGTCT	CGCTCTGTCG	CCCAGGCTGG	24850
AGTGCACTGG	CGCCATCTCG	GCTCACTGCA	AGCTCCACCT	CCCAGGTTCA	24900

CGCCATTCTC CTGCTCAGC CTCCCGAGTA GCTGGGACTA CACGCACCCG 24950  
 CCACCACGCC CGGCTAATTT TTTTGTATTT TTAGTAGAGA CGGGGTTTCA 25000  
 CCGTGTAGC CAGGATGGTC TCTATATCCT GACCCCATGA TCTGCCCGCC 25050  
 TCGGCTCCC AAAGTGGTGG GATTACAGGC GTGAGCCACT GCGCCCGGCC 25100  
 AACACTCTTT TTATTATTAG CAAATATACT TCTGCTGGG CACATTCTTG 25150  
 CAAGTGCTCA ACAATGCAAC TTTTGAAGT GCATGTGGCA GAAACTCCTG 25200  
 CTGTATTAT TCCAGAACCT ATTATTGCTA ATCCAGTTT ATGTTACATT 25250  
 TGAAGTGAGA ACCAGTTGGA GCCAGCAACG TTCCAGCTC CAAAGTTCCC 25300  
 TTGAGATTTT CAGAATCACT TAACCTATT ATGCTTGGCA ACCTGGACTC 25350  
 AGCAAACTG GGAAGTCAGC AGTTTGT TTTTATCCCT TCCTTTCTCA 25400  
 GTTCTCAAA TGTGTAGTT AATCTCAGTA ACCCATTCG AACCTTCATT 25450  
 ACCTGCCCAA GCGGTCTAGA ACTTGCCAGT ATAGAATCCT ACGTGGGTCA 25500  
 AGCTCCTGAC TGTCTCCTTC TTCACTCTTT TTTTGCAAG AACTTGTAAA 25550  
 TTTTAACTAT AAGTATTCAT GATTGCGCAC ATTTATTCAA AACATAGAGT 25600  
 GCTTTTCCA CATATCAGCC AATGGAATA AGGATTAAAT GGGAAATGAA 25650  
 ATGTAGTAAT AGGATAAGCA CAAGTCTTCT TCCTGCTCAA ACTTTT TTTT 25700  
 TTTT TTTT CAGACAAGAT CTGCTCTGT TACCCAGGCT GGAGTGAGT 25750  
 GGCGTGTTC TAGCTCAATG TAACCTCCAA CTCCTGGGCT CATGCAATCT 25800  
 CTCACACTC AGCCCTCTGA TTAGCTAGGA CTACACTATG CCTAGCCAAT 25850  
 TTTT TTTT TGTCTGGTT GTGTGCCCCA GGCTGCTCG ATCTCCTGGC 25900  
 CTCAGTAAT CCTCTGCTC CGGCTTCTA AAGTGCTGGG ATTATAGGCA 25950  
 TGAGCCACTG TGCCCGTCT CAAACCTTT TTTCCAAAGT AAATGAAAGT 26000  
 ATTAGATATG GAATATAGTC TAGTTCCAG ATATCCATAT CCATTGGTTT 26050  
 ATTACCTCA TTATTAACTT CAAATTGTTT AATAGACCT CATATCTCAG 26100  
 TTATACAGTT AAAATTTTGT TTTTGT TTTT CTGGAGTATC TTATTATAA 26150  
 CTATGAGTTT TACTTTACTT ATTTATTTTA TTTTGTAGA CAGACGCTTG 26200  
 CTCGTCTACT CAGGCTGGAG TGCGGTGCG TGATCATGGC TCACTATGGC 26250  
 CTCGACCTTC TGGGCTCAG TGATCCTCTC CCTCAGCCTC CCAAGCTGAG 26300  
 ACTACAGGCA TGCACCACCA CATCTAGCTA ATTTT TTTT CCTCCATGG 26350  
 AACAGGCTT TACTATGTTA CCCAGAGTGG TCTCAAATC CTGGCTCAG 26400  
 GGGATCCTCC TGTCTCAGCC TACCAAAATG CTGGGATTAC AGGCATGAGC 26450  
 CATAGCGCCA GACCTGGTTT TACTTTCTT GACTTTGAAT TACAAGTTT 26500  
 TGTAATTTGG AAAATGTTT GTTCTTTTA AATACTGCTG TATGTTTGCT 26550  
 TTTAAATACA ACATTTCTCG ATATATATTT TGAGAATTGC TGTCTTCAG 26600  
 AACCTAAGCAG TTTCTTAAG AAGGCTGATA TTTTCATCAA TGGGTGCGAG 26650  
 TTAGGAGAAG ATTTTATICA ATTGCATAAA CTCTAAGAA AGTCCACCTT 26700  
 CAAAATGCA AAACCTCTATG GTCCTGATGT TGGTCAAGCT CGAAGAAAGA 26750  
 CGGCTAAGAT GCTGAAGAGG TAGGAACCTAG AGGATGCAGA ATCACTTTAC 26800  
 TTTTCTTCTT TTTCTTTTG AGACAGAGTC TCACTCTGTC AGCCAGACTG 26850  
 GAGTGCAAGT GTACAATCAT GGCTCACTGC AACTTCGACC TCCAGGCTC 26900  
 AAGCAATCCT CCCATCTCAG TCCACAAAAT AGCTGGGACT ACAGGTGCAC 26950  
 ATCACCACAC CTGGCTACTT TAAAAAAT TTTTGTAGA GATGGGTCT 27000  
 CCCTGTGTTG CCCAGGCTGG TCTCTTGAAT TCCTGTGCTC AAGCCATCCT 27050  
 TCCACCTCAG CCTCCAGAG TGCCAGGATT ACAGGCTAGA GCCACCACAC 27100  
 CCAGCCACCA CTTTCTTAA AAAAAAAG AGATTCTCTC TGGTAGACAA 27150  
 TCCTCAATAG TCCACATGTT ATTAAACAAT CTGCTGCCTG AATACATGAT 27200  
 TTACCAAAAA AAGGAAATTT TGACGGGTTT AGAATATCAA GGGATCTGAG 27250  
 GCAAATGTCA CCTATGATAA AATTGCTAT CAAAATTAGG AAGTTTGTGT 27300  
 TTACCTGATC CTAAAGCAGT AACCAAGCCA TTCTAGGGA ATAAACTCT 27350  
 CATGCGTATA TTGTGCATAT ATATGTATTA TATGACTGAG TGATAATAA 27400  
 ATTTT TTTT TAGCTTCTG AAGGCTGGTG GAGAAGTGAT TGATTCAAGT 27450  
 ACATGGCATC AGTAAGTATG TCTCTATT TTAATACTAG GAAAGTAAGG 27500  
 CTAGCTTAT TTATTACCTA GTATTCAAAA AGTTAGTTCA TTTAACTGCC 27550  
 AATTGACTGC AGTTCAAATA AGAAACAAAT AGTGTCTCAA GTAGCACTGT 27600  
 ACTCCAATT TAATATTAAT AAAAAAAT TTAAGTTATT TTAATAATG 27650  
 TAGTGGTTTC TATAAGATC ACTTTATACA GAAGAACAGT GCCAATTAAC 27700  
 CCATGGAACA TATAAGTAGT TAAAACCAAT TGCTTGCCAA AGAACCAGTA 27750  
 ACCCAGGAGT ACATGCTCCT GCCACTGTGT TTTTCAAGA CAGAGTAACT 27800  
 GATTTCTAGT TACTTGCTA GAATGGACTC CTCCTCATAA CTCCTTTCCA 27850  
 TCTTGGTCTT TCCTAGTAG AACTTCTACC TTTT TTTT AACAGGTGAG 27900  
 TGGGAGAGGT AAGAAGGAGA ATAAGGTGAG CAATTAACCT AAAAGCAGAA 27950  
 AGTAAAAATT GTTATTTT TTCTGAATAT TTTCTGTGTA ATTTAGCTAC 28000  
 TATTTGAATG GACGCACTGC TACCAGGGAA GATTTCTAA ACCTGATGT 28050  
 ATTGGACATT TTTATTTT CATGTGCAAAA AGTTTTCAG GTAATAGTCT 28100  
 TTTTAACTT TTTAATGTAA AACCAAGATC CTATTTTAT AGTCTAGCTA 28150  
 GTTCTAAATT CTATAGGTAT GTATATTAC ATGTTTTCT AATTTTAGAG 28200  
 AACAGCACT ATGACTTATC CACTGTAGT TTTCCCTTA GCATTGGGTC 28250  
 TTACCCATG TACGTGATTA GAAATTGAA ATATTTCCAA TAGCCTTTAG 28300  
 TAGAATTAAC TCACATAGAT GATAAGAATG GGTGGTTCA CTTCATGTT 28350  
 CTCCACAGC CTACTATTTC AATAAAGAA AGTTTCCAA GACCTAAATG 28400  
 ACTATGAACA TATTTATAA CTATATAGGA GGGGTGGGTC TAGGAATACA 28450  
 AAGTTTGAAT TGCTGTTAAT CTCAACACC ACAGTTGAAA CCACAGGTCA 28500  
 GCTTTT TTTT AATTACCATG GATACTTTTC TGTCTATAG GTGGTTGAGA 28550  
 GCACCAGGCC TGGCAAGAG GTCTGGTTAG GAGAAACAG CTCTGCATAT 28600  
 GGAGGCGGAG CGCCTTGCT ATCCGACACC TTTGCAGCTG GCTTTATGTG 28650

AGTGAAGCAG CGCTGGCCTT AGGGGTGAGA GTGCAGCTCT TCTCCATCCT 28700  
 TCTATTCTGC TGAATAGACT CCCAGCCAA AAAGCAGATC AAAGACCGTT 28750  
 TCAGTGGCTG AGCCCCAAA TTCTATGCCAG ATTTTGCAAG AAAATGATT 28800  
 ACTAAAGCTT GAGGGACATC TTTAACAAAT GTTCCAAAT AATCACTATA 28850  
 AGGATGAATT GTTTCAGAAA TTTTGGCCTT TAATTATGGC CCATAAATAT 28900  
 GTCAGTAGT CTTTACTCTA AAGAAGTACA CTGTAAAAGA ATGCATATAG 28950  
 CCGGATATGG TAGTTCCTG TAATCCCAAT ACTTTGGGAG GCCAAGGTGG 29000  
 GAGGATTGGT TGAGCCGAGG AGTTTGAGGC TGCAGTGAGT TATGATGGTG 29050  
 CCCTGCACT CTAGACTGGG CAACAGAGTG AGACTGTCTT TTTTTCCTCC 29100  
 CTCTGTCACT CAGACTGGAG GGCAGTGGCA CGATCTCACC TCACTGCAAC 29150  
 CTCTGCCTCC CGGATTGAAG CGATTCTCT GCCTCAGCGT CCTGAGTAGC 29200  
 TGGGACTACA GGAGTATCAC CGCACTGGGC TAATTTTGT ATTTTATAGTA 29250  
 GAGACGGGGT TTTGACATGT TGCCGAGCT GGTCTGAAAC CCATGAGCTC 29300  
 AAGTGATCTG CCTACCTCAG CCTTCCAAA TGCTGGGATT ACGGACATGA 29350  
 GCTACCAAGC CGGCCCACAC CCTGTCTCTT AAAAAAATA AAAATGCAAG 29400  
 TTAGAGCATA TTACAGCTTT GTCTCTCAGG AGGATACTTA GTGTATGTAG 29450  
 CTATAATTCA TAGATTCCCA AGAAGTTTAG AGCCTAAAGT ATGAGGTCCC 29500  
 ACCAGAGGGG CTATCATTAA ATTTAAAGAT TTGTTAAATC ATCTCATTGT 29550  
 CCACACCCAC AAACCTGATT GCTTTAAAT ACTGGTTAG TTACATTAG 29600  
 TAACTCTATT AGTGTCTTTA ATCTATACCT CTATATCCTC ACATTGAGAT 29650  
 TTTTCTCTT TCTCTTCCA TCTTCTCTT TTTTCTCTC ATCCTCATTC 29700  
 TTATAAGCCT AGAATACATC ACAAATCCTT TATGCCCATG GAAGCAAGAG 29750  
 GAATAAGAA TGGAGATGTT TGTTTTGCCA TTAACATAAG ATCTGGGGTG 29800  
 TCGGGGAGAA GGGGGATAGA GAAGGAGAAG TGGGAAGAGG TGTCCTAAT 29850  
 AGCTTAGGTG CAATCTGCT TATTTTACAT TTTACCCCGC CTGACTGCCA 29900  
 CTTTCTCTT AGCCTCACA CATGTGTTGT GCAGGGACCT CATAGGACCA 29950  
 GGAATTGTCT ATAGAGGTGG GAATTTGTCT CACCCTGAAA GGGATACCTC 30000  
 TAGCATGGTA ATAGTCTTCT AGGATTGTGT ATCATATGGA AAGATGTAAA 30050  
 GGGAGGGATT CTGCTGCTGC TGCTGCTGCT GCATGCAGTT GCCATTCTAT 30100  
 TTAATGACT TATTTATAAT TGATGACACT TTTCTGGCTT CTGTTAATT 30150  
 CCTCCCTCAA AGATCAATAA ACCAGAACCA GGCATGGTGG CATGCACTTG 30200  
 TGGTCTGTA ACCACCCAC AGGTTCACT TGCTGCTGT CTAGATAGAG 30250  
 CCAATTATCA AGACAGGGGA ATTGCAAAGG AGAAAGAGTA ATTTATGCAG 30300  
 AGCCAGCTGT GCAGGAGACC AGAGTTTAT TATTACTCAA ATCAGTCTCC 30350  
 CCGAACATTC GAGGATCAGA GCTTTAAGG ATAATTGGC CGGTAGGGGC 30400  
 TTAGGAAGTG GAGAGTCTG GTTGGTCAGG TTGGAGATGG AATCACAGGG 30450  
 AGTGAAGTG AGGTTTCTT GCTGTCTTCT GTTCTGGAT GGGATGGCAG 30500  
 AACTGGTTGG GCCAGATTAC CGGTCTGGGT GGTCTCAAAT GATCCACCCA 30550  
 GTTCAGGGTC TGCAAGATAT CTCAAGCACT GATCTTAGGT TTTACAACAG 30600  
 TGATGTTATC CCAGGAACA ATTTGGGGAG GTTCAGACTC TTGGAGCCAG 30650  
 AGGCTGCATT ATCCCTAAAC CGTAATCTCT AATGTTGTAG CTAATTTGTT 30700  
 AGTCTGCAA AGGTAGACT GTCCCAAGGC AAGAAGGGGG TCTTTTCAGA 30750  
 AAAGGGCTAT TATCATTTT GTTTCAGAGT CAAACCATGA ACTGAATTC 30800  
 TTCCCAAAGT TAGTTCAGCC TACACCCAGG AATGAAGAAG GACAGCTTAA 30850  
 AGGTTAGAAG CAAGATGGAG TCAATGAGGT CTGATCTCTT TCACTGTCTAT 30900  
 AATTTCTCA GTTATAATT TTGCAAGGC GGTTCAGTC CCAGCTACTT 30950  
 GGGAGGCTGA GACAGGAGGA TTAATGGAGC CCAGGAGTTT GAGGTTGCAG 31000  
 AGAGCTATGA TCACGCCACT GCACTCCAGC CTGGGTGACA GAGTGAGACC 31050  
 CTGTCTCTAA ATAAATAAAT AAGTAAATAA ATAAATACAT AAATAAAATC 31100  
 AAGATGGTGT GCAATTAGAA TTGAGCGATT TTGTTTCCAA ACCTCAAGAA 31150  
 AGCTTGGTCT TGCTCTGTC CAGGTGGCTG GATAAATTGG GCCTGTGAGC 31200  
 CCGAATGGGA ATAGAAGTGG TGATGAGGCA AGTATTCTTT GGAGCAGGAA 31250  
 ACTACCATTT AGTGGATGAA AACTTCGATC CTTTACCTGT AAGTGACCAT 31300  
 TATTTCTCTA ATTCTAGTGG AGTAGATTAA AGTCAACTCA GGACCTCTGG 31350  
 TGTAAACCTC CTATGAACAG TCAGTCTCTT CAGTAACTAG CCAATCATG 31400  
 AGATGATGAA TTAGAAGGAG CCTTAGATAG CATCCAATCT AACATTTT 31450  
 TGTGTGTTG AAGAGAAGAA ATCAAGAGCT AGGAATAACT TTTTAAAGGT 31500  
 AAGCCATTGG CAGTATAGTG TGGATTTTGT TTAAGAGGGG ATAATTGAA 31550  
 ATTTTATGAC TCATTATACA AGACAAAATA AGTTGGATT TCAAATGTTT 31600  
 TACAAAGTAA ATCAAAGTTA TAATTGCCTA CAGTACGCAA AGCTTCAAAA 31650  
 CATTTTTTAT GTTATGAAAT TGTAATTTAT TTAACCTTAA AATGAGCCAG 31700  
 TACCATGTGT TTGCTTAAAA ATCTCATGCT AAGAATTTAC TATGTTGTTA 31750  
 ATAATCTTCA AGATATTTAT GAATAAAGTC TTATTTCTAA TCCTTCTCC 31800  
 AACTGTATCT GGTGCTAAAT CAGGAAATGT TTCTTCCCAA AAAGCCTCGT 31850  
 GGAAGATCTG TATGTCTAAA TATATGTCAG GGATAATACA GATGTAGCCC 31900  
 TGCGAAGCAT GACCTTGATT TTTATAGTCT AAAATGTAT TTAGAGATAT 31950  
 CTATTTCTA AGAATAATTC CTAAGAAGAT TATTTGAATG TTAGAGGAAA 32000  
 GCTAAGAAAT TTTGCAAGA GCGTACGTGA AAATATAAGC TAGGCTTTG 32050  
 TGGTTTGTGG ATAGACTTCC CAACAAAAT GCTTTTATC TATAGTGATC 32100  
 CAAGCTTGTG GAACATATTA GTCATCTTT TTTAGAAAAA TCTTAGAAAA 32150  
 GTGATCTTGC AAAAATGGAA TTTATCTTTC CCAAGTATA TTCTGTCTATG 32200  
 TATAGAGTTA AACTAAGCAT AGTAATTTCA CCAGACAAAC ATTCAAAATC 32250  
 TACTCTGAC CTTTTATCT CATCCAAAT TTCCAGGGC CCAGACATAA 32300  
 ACCTTGCCT TACGAATCT TGTATATGC ACTAAATATG CTCTCTCTC 32350  
 AAGTTCTCA GTCAGTAGA AAAATGTGCA AGAGTAAATG GTACCTTCT 32400

CACTGTGAGA	TCCAAGAGAA	TTAGACTTAA	ACTCACTCTA	CATGTCTGTG	32450
ACTTTATTTT	ATTTCGATGA	CAGTCCTGTG	AGGTGGCAAG	GCAGGTATCT	32500
TGGATCCATT	TTTTAGATAA	GGAAGTTCAA	ATTGAGAAGA	GGTTCGATGA	32550
TTTACAGGAA	GCCATACTGT	AGTCCTATGT	TACTCTTAAA	AATCCCATTG	32600
AAATCCTGCT	TCTGAGGCCT	GCATACTTTC	TACCCCTACCA	GTCATTGACC	32650
CATGCTTATG	TCTCCCTTGA	AAACATTGAT	TCCACTCTTG	TCTCCAGTGA	32700
AAAAGTGGAA	TTTAAGCAGA	GAAACAAAAG	CCATTGTGCT	TGTTAAGTCT	32750
ACTTCCCTC	TACTTTCAG	AAGGAAAGTT	GGGGTATGTG	TGAATGGTGT	32800
ATTTATTTAT	TTATTTATTA	TTTTAAAAAT	TGATACAAGG	TCTTACTGTA	32850
TTGTGCAGGC	TGCTCTCAAA	CTCCTGGGCT	CAAGTGATCA	TCCCACCTCA	32900
GCCTCCCGAT	GTGGGATTA	CAGCATGAAC	CATTGTGCC	ACCACCGATC	32950
CGCAGTTTTT	TAAGAAAAAC	TTTTACTATA	GAAAATTTTA	ATCATATACA	33000
AAATACAGAG	GAAAGTATAT	GAACCCACTT	TAGGAGACTA	GAATATGCCA	33050
CCCCAAAATA	TGCCACTTTG	GCATAAGGAT	TATTTGAGC	TAAAGGCAAC	33100
TGGGAAGAAA	CACATAGAAG	AAAAGTTCTC	TGTCCTTCTC	CATTTCCTCA	33150
AAAGCAGGAC	ATGAATCTTA	AAAGTCCCC	TCCCTCCCTT	TCTACCAGGA	33200
AAAACAAGAG	TTAATCACTG	AAGATAACTT	CAGACCCCTT	TCAGTGTAGA	33250
GATGGCACTA	GAGGAATCTA	TATTACATAC	TCATTTATTT	TCCTTCCCAC	33300
AACCTGCCAC	CCCAGAGACT	AAAAATCCTT	TTCCTTTGTC	ATGTCTCTTG	33350
TCCAAAAATT	TGCTCTATAA	GCTGGAGTTC	TAAGCCACCT	CTTTGAGAAT	33400
TACTTGTTC	CTGGTATTTT	CTGTTAACAT	ACATGTATTA	ATATACATGT	33450
TAACAAGCTT	CTGTTTGTGT	TTCTCCTGTT	TTCTGTCTTG	TTACAGAGGT	33500
CCATCCCAAC	TAAGAACTAA	AGAGTAGGAG	GAAAAATATA	TTTCTCTCTG	33550
CATACTTTGA	TCTTGTTTAA	TCCGTAAACC	TTCCCACTTT	TCACCTCCTA	33600
CCTATTAGAT	TACTTTGAAG	CAAAATTCAG	ATATATTACT	TTATCTATAA	33650
ATATTTCACT	ATGTGCTAGG	TGTGGTGGCT	CACACCTGTA	ATCCCAACAC	33700
TTTGGGAAGC	TGAGGCAGGA	GGATCACTTG	AGCCCAAGGAG	TTCAAGACCA	33750
GCTACGGCAA	CAAAAAATCA	AAAACCTTAT	TGGGCATGGT	GGCAGATGCC	33800
TGTGGTCCCA	GCTACATGAG	AGGCTGAGGC	AGGAGGATCG	CTTAGCCCCA	33850
GGAGGTTGAG	GCTGCAGTAA	GCTGCATTCA	CACCACTGCA	CTCCAGCCTG	33900
GGTGACAGAG	TAAGACCATG	TCTCAAAAAA	ATACATATTT	TAGTATGTAT	33950
CCTTTTGTGA	AAAAACAAAT	ACTTTTATCA	TACTTTAAAT	AATAACAAATA	34000
ATTCCTTAGT	ATCACCAAAT	ATTTTGTGAG	TGTCTCACAT	TTTCCTTATT	34050
GTCTAAAAATA	TTGTTGATAG	TTATTCAAAT	CAGAATCCAA	ACAAGGTCCA	34100
TATATTACAT	TTGGTTGACA	AGTCTCTTAA	GTTGTTCAT	CTTTAAGTTC	34150
TTCTCCCTCT	TCTTTTCACT	CTTGTAAATTT	ATTAATGTGA	AAAAACAGGT	34200
AATTTGTGTT	ATAGTATTTT	CTACATTATA	GAGTTTGCTA	CATTTATTCC	34250
CTATGATATC	ATTTAGCATG	TTCTCTGTGC	CCCTGTGTTT	CCTGTAAACT	34300
GGTAGTTATA	CCTAGAAGCT	TGAGTTTATT	CAGGTTTFTA	ATTGTATTTT	34350
TTTTGCAAGA	ATTTCTTTATT	ATCTGCTTCT	GGAAGCACAG	AATGTCTGGT	34400
TGTGTCTGGT	TTTGATCTTG	ACAGCTACTG	ATGACCATTG	CCTAATCCAT	34450
TACTTTATTT	GGGTGGGGGG	AATAAGGTTT	TAAAAATAAT	TTTTTTTAAA	34500
GATTTTTTTA	ACTGTTATTT	TGAGACAGTG	TCTCATTTTG	TTTCCCAGGC	34550
TGGAGTGCAG	TGGCACAAATC	ACGGCTCACT	GCAGCCTTGA	CCTCTGGGGA	34600
TCAGGTGATC	TTCTCACCTC	AGCCTCCTGG	GTACCTGGAA	CTACAGGTGC	34650
ACACCACCAC	ACCTGGCTAA	TTTTTTGTAT	TTTGTGTACA	GAAGGGGTTT	34700
CATCATGTTT	CCCAGACTGG	TCTTGAATC	CTGGGTTCAA	GTGATCTACC	34750
CACCTCAGCT	TCCCAAAATC	CTGGGATTAC	ACTTTGGCCA	CCGTGCTGGG	34800
CCTAAATGAA	ATTATTGTCT	TCTAAACAGA	CAGAAAGTTT	ACTTTAAAAA	34850
TTTGTCTTTG	TGTGTACATG	TGTTTGTGTA	TGTGTGTGTG	TCTAAAAGTT	34900
TGGCTTTGAG	CTTTGCTTTG	AATTCTTGGA	TGAACAATAA	CCAAGAATAC	34950
TTAACTCTG	ATCATCTCTG	ACAGATATCC	CCTACAGGCT	ATGGCCTTTT	35000
GAATTGTGTC	CTCCAGTGAT	AAAAAGCAGC	AAGCACGATA	CTGCTCTCAG	35050
ATTTCATGGT	GTACATGTGT	AGGTGAAAAA	AAAAAAAAG	ATGAATCCTA	35100
TTTAAATGCC	CCCAGGATAA	CAGTGATACT	CTTTGTAGGA	TAACTATTGT	35150
CTTGCCACTG	GTTTCATTAA	ATAAGGACAT	AAGTAAAGAT	CTATTTTGTG	35200
CTCTTCTCC	CCAACCACCA	CAACTAGGAT	TATTGGCTAT	CTCTTCTGTT	35250
CAAGAAATGT	GTGGGCACCA	AGGTGTTAAT	GGCAAGCGTG	CAAGGTTCAA	35300
AGAGAAGGAA	GCTTCGAGTA	TACCTTCATT	GCACAAACAC	TGACAAGTAA	35350
GTATGAAACA	CACCTTTTAC	CAATCATCAA	GTTTGTAGTG	GTAAGCCTGT	35400
AACCTTACTC	AAACCCCTG	TTGCATGTGT	CTATACATTG	CATAAGTATA	35450
GGCAGTTGCA	ATTTAGTAAA	GTTTATATACA	ACGATTTTAT	TTTATTTTAT	35500
TTTGAAGA	AAAAATGCTAC	TTTTGTTGTT	GTTGTTTTTT	GAGACGGGGC	35550
CTGCTCGTC	ACCCAGGCTG	GAGTGCAGTG	GTGCAATCTC	AGCTCACTGC	35600
AACCTCGCC	TCCCGGTTT	AAGTGATTCT	TGAAGAGGAG	AACAATAATA	35650
ACAACAATAT	TATTTTCAAA	AGTTGTGACC	GCAGTTTCTG	GAGTTGAGAA	35700
GACATCGAGA	TTTTGTGAGC	CTCATACTCT	TGCTTTAGGT	AGCAAAAAAT	35750
GTTCTTAAAT	CTCAGGAATA	TTCTCTAGAT	AGGTTTCAAT	CTATCATTTT	35800
TGATAAGATG	ATGCTGAAAT	ACTAATCTTA	GCCAAAAAAG	ACCAGCTACC	35850
ATTTCCGATT	GTTGGGGACT	GGGAACCTCT	GATAGTGAGG	ACCCAGTAG	35900
GAAGTAGCGA	GGGGAATGGT	TTGAATGGAT	AAATTCATAA	AAAAATGTCAG	35950
TAGATTAAAT	TTTCTTATAC	ATTTCACTCT	TTTATAAAGG	CTAGGAAAAA	36000
CCCTGTGTTT	TATGGTTTAT	AATTTGAATT	CACATGAACC	CACAAAAATT	36050
GCCTTTTACC	TTCTTATGTC	TGAAAAATGGA	TAGTCTGGCT	GGCCTCTTAA	36100
CAACCAGGCT	GGCAGAGCTG	TGAGGATCTC	AGTGTGCTCT	AGCCACAGCA	36150



TTGGTAGCAT	GAACGGCAAC	ATTTTAAATT	GTGTTTTC	AATAGGAGCA	36200
CACTAGCCGT	CTAAAACGAT	CATAAAGAA	GGATACTAAG	AGGGCCCACT	36250
GTCTATTATG	ATCCTAATAC	TTAGGATGCA	TTATGGATTG	TCATTATGGA	36300
TACTAATACT	TAGGATCACA	TTTGTAATTG	AGTTTTTAAT	TGCTTAAATT	36350
AGATACATAT	TTCTATTAA	TTAACCTCTT	TGCTTTTAGT	CCAAGGTATA	36400
AAGAAGGAGA	TTTAACTCTG	TATGCCATAA	ACCTCCATAA	TGTCACCAAG	36450
TACTTGGCGT	TACCCTATCC	TTTTTCTAAC	AAGCAAGTGG	ATAAATACCT	36500
TCTAAGACCT	TTGGGACCTC	ATGGATTACT	TTCCAAGTAA	GTAATTTTCC	36550
TTGTTCATTC	CAAACTTTCA	ATAAATTAT	TGGTGTAT	CAGAATAGAG	36600
AGTTTGGACA	GGGAGCAAAA	GACAAAGTCA	ACTATATCAA	GTTCTAATAA	36650
TTCTTAATAT	TCAGGAAATT	TATGTATGAA	TACTTACTAA	TATGAGTATA	36700
ACTCATCTTA	AGAGTCTAAA	GCAAAAGGAT	GTGAACACAA	ACTAGCAGTT	36750
ATCTTAGAGA	ATAAGTTTGC	ATTTCAAAAT	AACCTTGACAT	ATCAAGATCC	36800
ACTCAACGCA	TTTAAATTAT	TTACTCTAAA	AAGACATAAT	TCTTGGTAAC	36850
ACATTCACTA	AAGCAAAATA	TACCTTTATA	TAATTGCTAT	CAAAGGTATG	36900
TGGGTGGTAT	TAAAAATATCA	TACCATGTGA	GATCAGTGTG	ATTCCTTTAC	36950
AGCATTAATT	TTTATTGGTT	AGAGTAAGAA	AAAGAATAGC	TAGAGTATAT	37000
TTCTTAAGTA	GATTCTCATA	CACCTTGGTT	TCAAAAACCA	ATTATTGACT	37050
ACATCTTATA	AAAGCCTGTA	TTCAATGGAG	TGCCAAAAAA	TGACTATGAG	37100
TCTTAAAGAG	TTAGGCATAT	AAATATTTTA	AGGTTTCTGT	TCAATGTATG	37150
TTGGAAGGAG	TTCTTTTCTC	ATGACTATTC	TCATATTGGA	GCATAAAAG	37200
AGTTTACAGG	CTTGGCGCAG	TGGCTCATGC	CTGTAATCCC	AATACTTTGG	37250
GAACTGGAAG	CAGGCAGATC	ACTTCAGCCC	AGGAGTTTGA	GACCAGCCTG	37300
GGCAATATGG	CAAAACTCTC	TCTACAAAAT	ATACCAAAAT	TAGCCAGGCG	37350
TGGTGGTGCA	TGCCCTGTAGT	CCCAGCTACT	TGGGAAGCTG	AGGTGGGAGG	37400
ATTGCTTGAG	CCCAGGGGGG	TCATGGCTGC	AGTGAGCTGT	GATGGTGCCT	37450
CTGTCAACCA	GCCTGGGTGA	CAGAGTGAGA	CCCTGTCTCA	AAAAAATAAA	37500
TAAATAAAAA	TAAAGAGTTT	ACAAAATCTT	CACCATCTCC	TCCCATCTTT	37550
GCAAAATGCCA	CATAAGTGAT	GTGTTCCAGG	ACTATTAGCC	TCCGAACCTG	37600
AGGCAGTACA	GTAAGCACGC	TTTCTCCAAA	GTCTGTCTCC	CCACAGACAA	37650
ACATTATTTA	CACCTGGGTAC	TGCTCTTTTA	TTTTTTCCCC	TCTATGCTTT	37700
ATTTTACTAT	AACCTAATATC	ATATAACATG	TAATAGGAAA	AAGGCAGGGT	37750
CGGGGAGAG	ATCCAGAAAT	CTTCCCAAGA	GCCTTTCCAA	CATAGCCTCT	37800
GTAGACATTT	TTCTTTCTTT	CTTTTCTTTT	TTTTTTTTTT	TTCTGAGACA	37850
GAGTCTCACT	CTGTGTGTTA	GGCTAGAGTG	CAGTGGCGTG	ATCTAGGCTC	37900
ACTGCAACCT	CCGCTCTCTG	GGTTCAGGCA	ATTCTCCAC	CTCAGCCTCC	37950
CTAGTAGCTG	GGATTAGAGG	CATGCATCAC	CACGCTGGCC	TAATTTTGTG	38000
ATTTTATAGTA	GAGATGAGGT	TTCAACATGT	GGGCCAGGCT	GGTCTTGAAC	38050
TCCTGACCTC	AAGTGATCCA	CCTGCCCTAG	CCTCCCAAG	TGCTAGGATT	38100
ACACGAGTGA	GCCACCGTGC	CCTGCCCTTA	TTACATTCTG	ATCACACATT	38150
TCATGTTTTA	TAATTGGAAA	ACTGGTGAAA	TTATAGACAA	TGTTTTGTTC	38200
CCCTAAATTC	TCCTTGATGA	GTATATATTA	CTTACACTCT	TCGTCTTTTA	38250
AAATTTTGCA	AAATAGTATC	CTAGATAAGT	TTATGAGTGC	ACAGTCTGTA	38300
CGCTTACTCA	TATTAATGAC	CTCGGAGAGT	TAAACAACAG	TCACCTTTAA	38350
AAATTTATAT	TATCATTTAT	ATTATTTTGT	AGGCGGGGGT	CTCATTCTGT	38400
CTCCAGGCTC	GGAGAGTAGT	GGTGCGGTCA	CAGCTCACTG	CAGCCACGCG	38450
TACCTGGGCT	CAAGTGATCC	TTCTCTCTCA	GCCTTCTGAG	TAGCTGAGAC	38500
CACAGGCTTA	TGCTACCACA	CCTGGCTAAT	TTTTTAACTT	TTTGTAGAGA	38550
CGATGTCTCA	TTATGTGTGC	CAGGCTGGTC	TCAAACTCCT	AAGCTCAAGT	38600
GATCTTCTCT	AGCTCTCCAA	AGTGCTGGGA	TTACAGGCAT	GAAAAACTGC	38650
ACCCAGCCCT	AAAAATTATT	AGGGTCTCTG	ATAGTAAGAC	TTTAATAAAT	38700
ATTTAAATGA	ACATCTGGTT	TTTTTAAAAA	AAAAATAGAG	ACAAGGTCTC	38750
ACTATATTGC	CCAAGCTGGT	CTCGAACTCC	TGGACTCACG	CAATCCTGCT	38800
GCCTTAGCCG	CCCAAAGTGC	TGGGATTACA	GGCATGACCC	ACCTCATCTG	38850
GGCTGAGTGA	ACATATTTT	AACATAAAGG	CCGTATTTTA	TATTTATCTC	38900
ATACATTTTG	CCCAGCATCC	CCATTTCGCG	CGAATCTGTT	GCTTGCTAAT	38950
TCCTTCCAGC	TTCATTTTCAT	CTGAAATTTG	ACAAACATCT	TCTATTTCTT	39000
TGTCGTCTAG	TTATTGACTT	CAGAATATAA	AATAAAACAC	TATACCCAAA	39050
TTAAACCCCA	CCCTCATTTG	CCAGCCTGAT	GTGAAAAATA	TCAGCATACA	39100
TTAAGCTTAC	CCTTGATATA	TGTGTAGCAT	CTTTTAGATA	AATATACAGC	39150
TGATTAAGCA	ATATAGCCTG	ATGGTATAAT	ATCTTGCCCA	TGTACCTCAT	39200
CTTATCTCCA	GCAGGATTAA	TTACAGTGTA	TCAGATTTAC	CTTTAAACTT	39250
TGTAGCAAAA	TATCCTCTCC	AAAAGCATAT	CTAAAACCTT	TGTGTGTACT	39300
CTTGCAAGTT	TCTTAATTTT	ATGCAGAAC	GGCTCTTACC	ACTGTTAGCT	39350
GGAGATATTT	TCAAGACCTA	TTTTTGTTTG	TGGTTTCTGT	ATGATGGTCA	39400
TGGCATTTCC	CCCTTCACTC	CATCTAAAAA	TGAGGTGAT	ACAGGCTTTT	39450
AAACAAACCC	AACCTCATATA	GACTGAGTAC	AACGTCAATG	CAGGCATGCT	39500
AACCTCTGCT	ACAATCATGG	GCGTGTCTAT	GATATGTCTT	AAGTTACAGA	39550
ACACAGGGCT	GAGCGTCTCA	TTAGGTCAAA	ATGTAAACCA	GTTTTTCTGC	39600
TCACGTATGC	TTAATGAGGA	CAGGCTGTGA	GAGATTTCTT	TAAGGAAAAAC	39650
AAATATATAA	TAATGCTACA	TGGAAAAATA	TCTAACATTA	GAGAATTAAG	39700
TAAATAAACT	AATATACTCA	CACCATGGAA	TCTTGTGCAG	ACATTAATAA	39750
TATGTAGTGG	ATGGATGTTT	AATGGTGTGA	GAAAAAGTTA	GGATGTGCTG	39800
GGGTGGGGGG	AAGAAATCAAG	TTTTAAGAAA	ATACAGTATA	CCCATACTTA	39850
AGTAAAAAAA	AAAAAAAGG	TATGTACAGT	CATGTGTTGC	TTAATGATGG	39900

GGATACATTC	CGAGAAATGT	GTGATAGGT	GATTTTCATCC	TTGTGTGAAC	39950
ATCATAGAGT	GAACCTTACAC	AAACCTAGAT	GGTCTAGCCT	ACTATGTATC	40000
TAGGCTATAT	GACTAGCCTG	TTGCTCCTAG	GCTACAAACC	TGTAAAGCAT	40050
GTTACTGTAG	CGAATATACA	AATACTTAAC	ACAATGGCAA	GCTATCATTG	40100
TGTTAAGTAG	TTGTGTATCT	AAACATATCT	AAAACATAGA	AAACTAATGT	40150
GTGTGTCTAC	AATGTTACAA	TGACTATGAC	ATTGCTAGGC	AATAGGAATT	40200
ATAATTTTAT	CCTTTTATGG	AACCACACTT	ATATATGCGG	TCCATGGTGG	40250
ACCAAAACAT	CCTTATGTGG	CATATGACTG	TATACATGTA	CACAAAAAAT	40300
AGATGAAAGA	ATGAATATAC	ATCAAAATAT	TTAAAATGGT	TATAATGACT	40350
TAGGTACTTT	TTATTTATCT	TAGTAATAAT	AATGATGATA	GATAATACTT	40400
TTATAGTGGT	TACTATATAA	AAGACACTGT	TATAAGTGTT	CTACATACTT	40450
TACATGTATT	ACCTAAATGA	TATAAATATA	ACTCTGACAG	TAACTAATCT	40500
TATACGTTCT	CTTTTCTTTT	TTTTTTTTTT	CTTTTTTTAG	ACAGAATCTT	40550
GCTCTACCAG	GCTGGAGTGC	AGGGTGCAAT	CTCGGCTCAC	TGCAACCTCC	40600
GCCTCCCAGG	TTCAAACGAT	TCTCATGTCT	CAGCCTCCTG	AGTAGCTGGG	40650
ACTACAGGCA	CACACCACCA	TGCCCGGCTA	ATTTTTGTAT	TTTTGGGTAG	40700
AGATGGAGTT	TTGCCATGTT	GGCCAGGCTG	ATCTTGAAC	CCTGGCCTCA	40750
AGTGATCTGC	CTGCCTCAGC	CTCCCAAAGT	GCTGGGATTA	CAGGTGTGAA	40800
CCACTGTGCT	CGGCCTAATC	TTACAAGTTT	TCAATATTTA	AAGAGTGCTA	40850
ACTTTGTGGA	CAATATAAAA	CATATTTGAG	AAAAAGAGAT	ATAAGCATCT	40900
TATTTAGAAT	TATGAAAAAT	TCAATAGACC	TACAGCCGAC	TAAAGCTTTT	40950
CTTCATAAGC	TCTTGCCTAT	ATTGATTGCG	TCTGTGAAT	ATGCATTAAT	41000
TTGATTTAAA	TAATAAGTAT	GTATAAGAAA	TAACACTTTT	CCTTAATTTT	41050
TAAGAACGTT	CAACAGTTT	TAATTTGAAT	TCCAATAGTG	AAATACATAG	41100
AAAAATATAA	ATTTTCTGTA	GTTTAGCCAA	ATTGTTTTTG	TTTACCACCA	41150
GCATTCTACC	AAAATTTCTT	AATAACAGTA	AGAAAATGAA	TGCATACCTC	41200
CTGCAGGGAG	AGGGGAGTTA	GGCAGTTTAT	GGCATAAGT	ACAAGTGAGA	41250
AAITTCATTG	GCTACCAAT	ACGCTAAAT	CATAAAAACT	GCATTCAATT	41300
CTATATATCT	ATTTTCTTTA	CATAAAAAAG	GTTTCAATTA	TTGGCCATTA	41350
AATAAAATAG	CCACCAATCC	AGAAGTTGTG	TCATGTTTAT	CCTTTTTATA	41400
CCACCATCAT	ATTGCCTATT	ATATAGATTG	TGTGTGTTCC	ATTTTCTGTA	41450
ATGGGCCAGA	CAGTAAGTAT	TTCTGGCTTT	GGAGTCCATA	TGGTCTCTAT	41500
CATAACTACT	CATCTCTGCC	ATTGTAGCTT	AAAGATTATC	TAGGTCAAAT	41550
GCCTAAGTGA	TATAGTGTG	AAATACAAGT	TATATAATAT	AGGCTGCCAC	41600
AAAAAAAAT	TTATTTGGTC	TAAAAAAGAT	TTCATGACTT	TTGTAGCAGC	41650
ATGGGTGGGG	CATGCACCAC	TTGGTTAACT	CGGTGTATCT	TTCTCCTTTG	41700
CAGATCTGTC	CAACTCAATG	GTCTAACTCT	AAAGATGGTG	GATGATCAAA	41750
CCTTGCACC	TTAATGGAA	AAACCTCTCC	GGCCAGGAAG	TTCACTGGGC	41800
TTGCCAGCTT	TCTCATATAG	TTTTTTTGTG	ATAAGAAATG	CCAAAGTTGC	41850
TGCTTGCAAT	TGAAAAATAA	ATATACTAGT	CCTGACACTG	AATTTTTCAA	41900
GTATATAAG	AGTAAAGCAA	CTCAAGTTAT	AGGAAAGGAA	GCAGATACCT	41950
TGCAAAAGCAA	CTAGTGGGTG	CTTGAGAGAC	ACTGGGACAC	TGTCAGTGCT	42000
AGATTTAGCA	CAGTATTTTG	ATCTCGCTAG	GTAGAACACT	GCTAATAATA	42050
ATAGCTAATA	ATACCTTGTT	CCAAATAC	CTTAGCATTT	TGCATGTTTT	42100
ACTTTTATCT	AAAGTTTTGT	TTTGTTTTAT	TATTTATTTA	TTTATTTATT	42150
TTGAGACAGA	ATCTCTCTCT	GTCAACCCAGG	CTGGAGTGCC	ATGGTGCGAT	42200
CTTGGCTCAC	TGCAACTTTA	AGCAATTCTC	CTGCCTCAGC	TTCTGTAGTA	42250
GCTGGGATTA	TAGGCGTGTG	CCACCACGCC	CAGCTACTTT	CTATATTTTT	42300
TGTAGAGATG	GAGTTTCGCC	ATATTGGCCA	AGCTGGTCTC	GAACTCCTGT	42350
CCTCGAACTC	CTTGCCTCAA	GTGATCCACC	CGCCTCAGCC	TCTCAAAGTG	42400
CTGGGATTAC	AGGTGTGAGC	CACCACACCC	AGCAGTGT	TATTTTGTAG	42450
ACAGGGTATC	ATTCGTGTTG	CCAGGCTTGA	GTGCAGTGGT	GCAATCATAG	42500
ATCACTGCAG	CCTTTTAACT	CCTGGGCTCA	AGTCATCCTC	CTGCTTAGCC	42550
TCCCAAGTAG	CTAGGACCAC	AGACACATGC	CATCACACTT	GGCTATTTTT	42600
AAAAAATTTT	TTGTAGAGAT	GGGGTCTCGC	TATGTTACCC	AAACTGGTCC	42650
TGAATCCTG	GACTCAATTG	ATCCTCCAC	CTTGGCCTTC	CAGGTGCTGG	42700
GATTTCTTTG	GGAGTACAGC	ATGGTACAGC	AGGAGATCAT	TTGATGTTAC	42750
CTCTGTGCAG	TGTTGTCTAGT	CAGCGAAAGA	CTATAATACC	TGTGGGGACA	42800
GCGATTAGCC	ACCACAACCA	GTCTTTATTT	AAAGTTATTA	AAAATGGCTG	42850
GCGCAGTGG	CTCACACTG	TAATCCTAGC	ACTTTGGGAG	GCCGAGGCAG	42900
ATGGATCACC	TGACGTGAGG	AATTTGAGAC	CAGCCTGGCC	AACATGGTGA	42950
AACCCCATCT	CTACTAAAAA	ATACAAAAAT	TAGCTGGGTG	TGGTCTGTGA	43000
GTCCAGCTA	CTTGGGAGGC	TGGGGCAGGA	GAATTAAGTG	AACCCAGGAG	43050
GCAGAGGTTG	CAGTGAGCCG	AGATTGTGCC	ACTGCACTCC	AGCCTGGGTG	43100
ACAGAGAGAG	ATTCCATCTC	AAAAAAACAA	GTTATTTAAA	ATGTATATGA	43150
ATGCTCCTAA	TATGGTCAGG	AAGCAAGGAA	GCGAAGGATA	TATTATGAGT	43200
TTTAAAGAGG	TGCTTAGCTG	TATATTTATC	TTTCAAAATG	TATTAGAAGA	43250
TTTTAGAATT	CTTTCCTTCA	TGTGCCATCT	CTACAGGCAC	CCATCAGAAA	43300
AAGCATACTG	CCGTTACCGT	GAAACTGGTT	GTAAAAGAGA	AACTATCTAT	43350
TTGCACCTTA	AAAGACAGCT	AGATTTTGCT	GATTTTCTTC	TTTCGGTTTT	43400
CTTTGTGACG	AATAATATGT	GAGAGGACAG	ATTGTTAGAT	ATGATAGTAT	43450
AAAAAATGGT	TAATGACAAAT	TCAGAGGCGA	GGAGATTCTG	TAAACTTAAA	43500
ATTACTATAA	ATGAAATTGA	TTTGTCAAGA	GGATAAATTT	TAGAAAACAC	43550
CCAATACCTT	ATAACTGTCT	GTTAATGCTT	GCTTTTCTC	TACCTTTCTT	43600
CCTTGTTTCA	GTTGGGAAGC	TTTTGGCTGC	AAGTAACAGA	AACTCCTAAT	43650

30

TCAAAATGGCT TAAGCAATAA GGAAATGTAT ATTCCACAT AACTAGACGT 43700  
 TCAAAACAGGC CAGGCTCCAG CACTTCAGTA CGTCACCAGG GATCTGGGTT 43750  
 CTTCCAGCT CTCTGCTCTG CCATCTTTAG CGCTGGCTTC ATTCTCAGAC 43800  
 TCTGGTAGCA TGATGGCTGT AGCTGTTTCA TGGGCCCTT CAAACCTCAT 43850  
 AGCAACCAGA GGAAGAAAAT GAGCCATTTT TTGAGTCTCC TTCATAGACT 43900  
 TGAATAACTC TTTTTCAGAG CTTCTCACAG CAAACCTCTC CTCATGTCTC 43950  
 CTCATGTCTT ATTGTTTACA AATGGGTAAT GTGGCCATTT CACCAGTCAC 44000  
 TGCCAAACAC AACGAGGTTT CTATAATTGT CTCTGAGTAA CCCTTTGGAA 44050  
 TGGAGAGGGT GTTGGTCAGT CTACAACTG AACACTGCAG TTCTGCGCTT 44100  
 TTTACCAGTG AAAAAATGTA ATTATTTTCC CCTCTTAAGG ATTAATATTC 44150  
 TTCAAATGTA TGCCTGTTAT GGATATAGTA TCTTTAAAT TTTTATTTT 44200  
 AATAGCTTTA GGGGTACACA CTTTTGCTT ACAGGGGTGA ATTGTGTAGT 44250  
 GGTGAAGACT CGGCTTTTAA TGTACTGTG ACCTGAGTGA TGTACATTGT 44300  
 ACCCAATAGG TAATTTTCA TCCATTACCC TCCTTCCGCC CTCCTCCCTT 44350  
 CTGAGTCTCC AACATCCCTT ATACCACTGT GTATGTTCTT GTGTACCTAC 44400  
 AGCTAAGCTT CCACITATAA GTGAGAACAT GCAGTATTGT GTTTTCCATT 44450  
 CTTGAGTTAC TTCCCTTAGG ATAACAGCCC CCAGTTCGT CCAAGTTGCT 44500  
 GCAAAATACA TTATCTTCTT TTATGGCTGA GTAATAGTCC ATGGTACATA 44550  
 TATACCAAGT TTTCTTTATC CACTTATCAG TTGATGGACA CTTAGGTTAA 44600  
 TTCCATTCAA TTTTATTCAA TTTAAGTATA TTTGTAAGGA GCTAAAGCTG 44650  
 AAAATTAAAT TTTAGATCTT TCAATACTCT TAAATTTTAT ATGTAAGTGG 44700  
 TTTTATATTT TTACATTTG AAATAAAGTA ATTTTATAA CCTTGATATT 44750  
 GTATGACTAT TCTTTTAGTA ATGTAAAGCC TACAGACTCC TACATTTGGA 44800  
 ACCACTAGTG TGTGTTTCA CCCCTTGTTA TACTATCAGG ATCCTCGA 44848

## (2) INFORMATION FOR SEQ ID NO:43:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 2396  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:43

TTCTAGTTG CTTTATGCCA ATGTCGGATC AGGTTTTTCA AGCGACAAAG 50  
 AGATACTGAG ATCTGGGCA GAGGACATCC TAGCTCGGTC AGATTTGGGC 100  
 AGGCTCAAGT GACCAAGTGC TTAAGGCAGA AGGGAGTCGG GGTAGGGTCT 150  
 GGCTGAACCC TCAACCGGGG CTTTAACTC AGGGTCTAGT CCTGGCGCCA 200  
 AATGATGAG ACCTAGAAAA GGTGACAGAG TCGCAGGAC ACCAGGAAGC 250  
 TGGTCCACC CCTGCGGGC TCCCGGGCGC TCCCTCCCA GGCTCCGAG 300  
 GATCTTGGAT TCTGGCCACC TCCGACCCCT TTGGATGGGT GTGGATGATT 350  
 TCAAAAGTGG ACGTGACCGC GCGGAGGGG AAAGCCAGCA CGGAAATGAA 400  
 AGAGAGCGAG GAGGGAGGG GCGGAGGGG AGGGCGCTAG GGAGGGACTC 450  
 CCGGAGGGG TGGGAGGGAT GGAGCGCTGT GGGAGGGTAC TGAGTCTTGG 500  
 CGCCAGAGGC GAAGCAGGAC CGGTTGCAGG GGGCTTGAGC CAGCGCGCCG 550  
 GCTGCCCGAC CTCTCCCGGC AGCGGGCGGT CCAGCCAGGT GGGATGCTGA 600  
 GGCTGCTGCT GCTGTGGCTC TGGGGGCCGC TCGGTGCCCT GGCCAGGGC 650  
 GCCCCGCGG GGACCGCGC GACCGACGAC GTGGTAGACT TGGAGTTTTA 700  
 CACCAAGCGG CGCTCCGAA GCGTGAGTCC CTCGTTCTG TCCATCACCA 750  
 TCGACGCCAG CTGGCCACC GACCCGCGCT TCCTCACCTT CTGGGCTCT 800  
 CCAAGGCTCC GTGCTCTGGC TAGAGGCTTA TCTCTGCAT ACTTGAGATT 850  
 TGGCGGCACA AAGACTGACT TCCTTATTTT TGATCCGGAC AAGGAACCGA 900  
 CTTCCGAAGA AAGAAGTTAC TGGAAATCTC AAGTCAACCA TGATATTTGC 950  
 AGGTCTGAGC CGGTCTCTGC TGCCTGTTG AGGAACTCC AGGTGGAATG 1000  
 GCCCTTCCAG GAGCTGTTGC TGCTCCGAGA GCAGTACCAA AAGGAGTTCA 1050  
 AGAACAGCAC CTAATCAAGA AGCTCAGTGG ACATGCTCTA CAGTTTGGCC 1100  
 AAGTGCTCGG GGTAGACCT GATCTTTGGT CTAATGCGT TACTACGAAC 1150  
 CCCAGACTTA CGGTGGAACA GcTCCAACGC CCAGCTTCTC CTTGACTACT 1200  
 GCTCTTCCA GGGTTATAAC ATcTCTGGG AACTGGGCAA TGAGCCCAAC 1250  
 AGTTTCTGGA AGAAAGCTCA CATTCTCATC GATGGGTTGC AGTTAGGAGA 1300  
 AGACTTTGTG GAGTTGCATA AACTTcTACA AAGGTCAGCT TTCCAAAATG 1350  
 CAAAACCTTA TGGTCTGAC ATCGGTCAGC CTCGAGGGAA GACAGTTAAA 1400  
 CTGCTGAGGA GTTCTCTGAA GGCTGGCGGA GAAGTGATCG ACTCTCTTAC 1450  
 ATGGCATCAC TATTACTTGA ATGGACGCAT CGCTACCAA GAAGATTTTC 1500

TGAGCTCTGA TGCCTGGAC ACTTTTATTC TCTCTGTGCA AAAAATTCTG 1550  
 AAGGTCACCTA AAGAGATCAC ACCTGGCAAG AAGGTCTGGT TGGGAGAGAC 1600  
 GAGCTCAGCT TACGGTGGCG GTGCACCCTT GCTGTCCAAC ACCTTTGCAG 1650  
 CTGGCTTTAT GTGGCTGGAT AAATTGGGCC TGTCAGCCCA GATGGGCATA 1700  
 GAAGTCGTGA TGAGGCAGGT GTTCTTCGGA GCAGGCAACT ACCACTTAGT 1750  
 GGATGAAAAC TTTGAGCCTT TACCTGATTA CTGGCTCTCT CTCTGTTCa 1800  
 AGAAACTGGT AGGTCCCAGG GTGTTACTGT CAAGAGTGAA AGGCCAGAC 1850  
 AGGAGCAAAC TCCGAGTGTA TCTCCACTGC ACTAAGTCT ATCACCCACG 1900  
 ATATCAGGAA GGAGATCTAA CTCTGTATGT CCTGAACCTC CATAATGTCA 1950  
 CCAAGCACTT GAAGGTACCG CCTCCGTTGT TCAGGAAACC AGTGATACG 2000  
 TACCTTCTGA AGCCTTCGGG GCCGGATGGA TTACTTTCCA AATCTGTCCA 2050  
 ACTGAACGGT CAAATTCTGA AGATGGTGA TGAGCAGACC CTGCCAGCTT 2100  
 TGACAGAAAA ACCTCTCCCC GCAGGAAGTG CACTAAGCCT GCCTGCCTTT 2150  
 TCCTATGGTT TTTTGTGCAT AAGAAATGCC AAAATCGCTG CTTGTATATG 2200  
 AAAATAAAAG GCATACGGTA CCCCTGAGAC AAAAGCCGAG GGGGGTGTTA 2250  
 TTCATAAAAC AAAACCCTAG TTAGGAGGC CACCTCCTTG CCGAGTTCCA 2300  
 GAGCTTCGGG AGGGTGGGT ACACCTCAGT ATTACATTCA GTGTGGTGT 2350  
 CTCTCTAAGA AGAATACTGC AGGTGGTGAC AGTTAATAGC ACTGTG 2396

## (2) INFORMATION FOR SEQ ID NO:44:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 535  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:44

Met Leu Arg Leu Leu Leu Trp Leu Trp Gly Pro Leu Gly Ala  
 5 10 15  
 Leu Ala Gln Gly Ala Pro Ala Gly Thr Ala Pro Thr Asp Asp Val  
 20 25 30  
 Val Asp Leu Glu Phe Tyr Thr Lys Arg Pro Leu Arg Ser Val Ser  
 35 40 45  
 Pro Ser Phe Leu Ser Ile Thr Ile Asp Ala Ser Leu Ala Thr Asp  
 50 55 60  
 Pro Arg Phe Leu Thr Phe Leu Gly Ser Pro Arg Leu Arg Ala Leu  
 65 70 75  
 Ala Arg Gly Leu Ser Pro Ala Tyr Leu Arg Phe Gly Gly Thr Lys  
 80 85 90  
 Thr Asp Phe Leu Ile Phe Asp Pro Asp Lys Glu Pro Thr Ser Glu  
 95 100 105  
 Glu Arg Ser Tyr Trp Lys Ser Gln Val Asn His Asp Ile Cys Arg  
 110 115 120  
 Ser Glu Pro Val Ser Ala Ala Val Leu Arg Lys Leu Gln Val Glu  
 125 130 135  
 Trp Pro Phe Gln Glu Leu Leu Leu Leu Arg Glu Gln Tyr Gln Lys  
 140 145 150  
 Glu Phe Lys Asn Ser Thr Tyr Ser Arg Ser Ser Val Asp Met Leu  
 155 160 165  
 Tyr Ser Phe Ala Lys Cys Ser Gly Leu Asp Leu Ile Phe Gly Leu  
 170 175 180  
 Asn Ala Leu Leu Arg Thr Pro Asp Leu Arg Trp Asn Ser Ser Asn  
 185 190 195  
 Ala Gln Leu Leu Asp Tyr Cys Ser Ser Lys Gly Tyr Asn Ile  
 200 205 210  
 Ser Trp Glu Leu Gly Asn Glu Pro Asn Ser Phe Trp Lys Lys Ala  
 215 220 225  
 His Ile Leu Ile Asp Gly Leu Gln Leu Gly Glu Asp Phe Val Glu

32

230	235	240
Leu His Lys Leu Leu Gln Arg Ser Ala Phe Gln Asn Ala Lys Leu		
245	250	255
Tyr Gly Pro Asp Ile Gly Gln Pro Arg Gly Lys Thr Val Lys Leu		
260	265	270
Leu Arg Ser Phe Leu Lys Ala Gly Gly Glu Val Ile Asp Ser Leu		
275	280	285
Thr Trp His His Tyr Tyr Leu Asn Gly Arg Ile Ala Thr Lys Glu		
290	295	300
Asp Phe Leu Ser Ser Asp Ala Leu Asp Thr Phe Ile Leu Ser Val		
305	310	315
Gln Lys Ile Leu Lys Val Thr Lys Glu Ile Thr Pro Gly Lys Lys		
320	325	330
Val Trp Leu Gly Glu Thr Ser Ser Ala Tyr Gly Gly Gly Ala Pro		
335	340	345
Leu Leu Ser Asn Thr Phe Ala Ala Gly Phe Met Trp Leu Asp Lys		
350	355	360
Leu Gly Leu Ser Ala Gln Met Gly Ile Glu Val Val Met Arg Gln		
365	370	375
Val Phe Phe Gly Ala Gly Asn Tyr His Leu Val Asp Glu Asn Phe		
380	385	390
Glu Pro Leu Pro Asp Tyr Trp Leu Ser Leu Leu Phe Lys Lys Leu		
395	400	405
Val Gly Pro Arg Val Leu Leu Ser Arg Val Lys Gly Pro Asp Arg		
410	415	420
Ser Lys Leu Arg Val Tyr Leu His Cys Thr Asn Val Tyr His Pro		
425	430	435
Arg Tyr Gln Glu Gly Asp Leu Thr Leu Tyr Val Leu Asn Leu His		
440	445	450
Asn Val Thr Lys His Leu Lys Val Pro Pro Pro Leu Phe Arg Lys		
455	460	465
Pro Val Asp Thr Tyr Leu Leu Lys Pro Ser Gly Pro Asp Gly Leu		
470	475	480
Leu Ser Lys Ser Val Gln Leu Asn Gly Gln Ile Leu Lys Met Val		
485	490	495
Asp Glu Gln Thr Leu Pro Ala Leu Thr Glu Lys Pro Leu Pro Ala		
500	505	510
Gly Ser Ala Leu Ser Leu Pro Ala Phe Ser Tyr Gly Phe Phe Val		
515	520	525
Ile Arg Asn Ala Lys Ile Ala Ala Cys Ile		
530	535	

## (2) INFORMATION FOR SEQ ID NO:45:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2396  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:45

	TT TCT AGT	8
TGC TTT TAG CCA ATG TCG GAT CAG GTT TTT CAA GCG ACA AAG AGA		53
TAC TGA GAT CCT GGG CAG AGG ACA TCC TAG CTC GGT CAG ATT TGG		98
GCA GGC TCA AGT GAC CAG TGT CTT AAG GCA GAA GGG AGT CGG GGT		143
AGG GTC TGG CTG AAC CCT CAA CCG GGG CTT TTA ACT CAG GGT CTA		188
GTC CTG GCG CCA AAT GGA TGG GAC CTA GAA AAG GTG ACA GAG TGC		233
GCA GGA CAC CAG GAA GCT GGT CCC ACC CCT GCG CGG CTC CCG GGC		278

33

GCT CCC TCC CCA GGC CTC CGA GGA TCT TGG ATT CTG GCC ACC TCC	323
GCA CCC TTT GGA TGG GTG TGG ATG ATT TCA AAA GTG GAC GTG ACC	368
GCG GCG GAG GGG AAA GCC AGC ACG GAA ATG AAA GAG AGC GAG GAG	413
GGG AGG GCG GGG AGG GGA GGG CGC TAG GGA GGG ACT CCC GGG AGG	458
GGT GGG AGG GAT GGA GCG CTG TGG GAG GGT ACT GAG TCC TGG CGC	503
CAG AGG CGA AGC AGG ACC GGT TGC AGG GGG CTT GAG CCA GCG CGC	548
CGG CTG CCC CAG CTC TCC CGG CAG CGG GCG GTC CAG CCA GGT GGG	593
ATG CTG AGG CTG CTG CTG CTG TGG CTC TGG GGG CCG CTC GGT GCC	638
Met Leu Arg Leu Leu Leu Leu Trp Leu Trp Gly Pro Leu Gly Ala	
5 10 15	
CTG GCC CAG GGC GCC CCC GCG GGG ACC GCG CCG ACC GAC GAC GTG	683
Leu Ala Gln Gly Ala Pro Ala Gly Thr Ala Pro Thr Asp Asp Val	
20 25 30	
GTA GAC TTG GAG TTT TAC ACC AAG CGG CCG CTC CGA AGC GTG AGT	728
Val Asp Leu Glu Phe Tyr Thr Lys Arg Pro Leu Arg Ser Val Ser	
35 40 45	
CCC TCG TTC CTG TCC ATC ACC ATC GAC GCC AGC CTG GCC ACC GAC	773
Pro Ser Phe Leu Ser Ile Thr Ile Asp Ala Ser Leu Ala Thr Asp	
50 55 60	
CCG CGC TTC CTC ACC TTC CTG GGC TCT CCA AGG CTC CGT GCT CTG	818
Pro Arg Phe Leu Thr Phe Leu Gly Ser Pro Arg Leu Arg Ala Leu	
65 70 75	
GCT AGA GGC TTA TCT CCT GCA TAC TTG AGA TTT GGC GGC ACA AAG	863
Ala Arg Gly Leu Ser Pro Ala Tyr Leu Arg Phe Gly Gly Thr Lys	
80 85 90	
ACT GAC TTC CTT ATT TTT GAT CCG GAC AAG GAA CCG ACT TCC GAA	908
Thr Asp Phe Leu Ile Phe Asp Pro Asp Lys Glu Pro Thr Ser Glu	
95 100 105	
GAA AGA AGT TAC TGG AAA TCT CAA GTC AAC CAT GAT ATT TGC AGG	953
Glu Arg Ser Tyr Trp Lys Ser Gln Val Asn His Asp Ile Cys Arg	
110 115 120	
TCT GAG CCG GTC TCT GCT GCG GTG TTG AGG AAA CTC CAG GTG GAA	998
Ser Glu Pro Val Ser Ala Ala Val Leu Arg Lys Leu Gln Val Glu	
125 130 135	
TGG CCC TTC CAG GAG CTG TTG CTG CTC CGA GAG CAG TAC CAA AAG	1043
Trp Pro Phe Gln Glu Leu Leu Leu Leu Arg Glu Gln Tyr Gln Lys	
140 145 150	
GAG TTC AAG AAC AGC ACC TAC TCA AGA AGC TCA GTG GAC ATG CTC	1088
Glu Phe Lys Asn Ser Thr Tyr Ser Arg Ser Ser Val Asp Met Leu	
155 160 165	
TAC AGT TTT GCC AAG TGC TCG GGG TTA GAC CTG ATC TTT GGT CTA	1133
Tyr Ser Phe Ala Lys Cys Ser Gly Leu Asp Leu Ile Phe Gly Leu	
170 175 180	
AAT GCG TTA CTA CGA ACC CCA GAC TTA CGG TGG AAC AGC TCC AAC	1178
Asn Ala Leu Leu Arg Thr Pro Asp Leu Arg Trp Asn Ser Ser Asn	

185	190	195	
GCC CAG CTT CTC CTT GAC TAC TGC TCT TCC AAG GGT TAT AAC ATc			1223
Ala Gln Leu Leu Leu Asp Tyr Cys Ser Ser Lys Gly Tyr Asn Ile			
200	205	210	
TCC TGG GAA CTG GGC AAT GAG CCC AAC AGT Ttc TGG AAG AAA GCT			1268
Ser Trp Glu Leu Gly Asn Glu Pro Asn Ser Phe Trp Lys Lys Ala			
215	220	225	
CAC ATT CTC ATC GAT GGG TTG CAG TTA GGA GAA GAC TTT GTG GAG			1313
His Ile Leu Ile Asp Gly Leu Gln Leu Gly Glu Asp Phe Val Glu			
230	235	240	
TTG CAT AAA CTT cTA CAA AGG TCA GCT TTC CAA AAT GCA AAA CTC			1358
Leu His Lys Leu Leu Gln Arg Ser Ala Phe Gln Asn Ala Lys Leu			
245	250	255	
TAT GGT CCT GAC ATC GGT CAG CCT CGA GGG AAG ACA GTT AAA CTG			1403
Tyr Gly Pro Asp Ile Gly Gln Pro Arg Gly Lys Thr Val Lys Leu			
260	265	270	
CTG AGG AGT TTC CTG AAG GCT GGC GGA GAA GTG ATC GAC TCT CTT			1448
Leu Arg Ser Phe Leu Lys Ala Gly Gly Glu Val Ile Asp Ser Leu			
275	280	285	
ACA TGG CAT CAC TAT TAC TTG AAT GGA CGC ATC GCT ACC AAA GAA			1493
Thr Trp His His Tyr Tyr Leu Asn Gly Arg Ile Ala Thr Lys Glu			
290	295	300	
GAT TTT CTG AGC TCT GAT GCG CTG GAC ACT TTT ATT CTC TCT GTG			1538
Asp Phe Leu Ser Ser Asp Ala Leu Asp Thr Phe Ile Leu Ser Val			
305	310	315	
CAA AAA ATT CTG AAG GTC ACT AAA GAG ATC ACA CCT GGC AAG AAG			1583
Gln Lys Ile Leu Lys Val Thr Lys Glu Ile Thr Pro Gly Lys Lys			
320	325	330	
GTC TGG TTG GGA GAG ACG AGC TCA GCT TAC GGT GGC GGT GCA CCC			1628
Val Trp Leu Gly Glu Thr Ser Ser Ala Tyr Gly Gly Gly Ala Pro			
335	340	345	
TTG CTG TCC AAC ACC TTT GCA GCT GGC TTT ATG TGG CTG GAT AAA			1673
Leu Leu Ser Asn Thr Phe Ala Ala Gly Phe Met Trp Leu Asp Lys			
350	355	360	
TTG GGC CTG TCA GCC CAG ATG GGC ATA GAA GTC GTG ATG AGG CAG			1718
Leu Gly Leu Ser Ala Gln Met Gly Ile Glu Val Val Met Arg Gln			
365	370	375	
GTG TTC TTC GGA GCA GGC AAC TAC CAC TTA GTG GAT GAA AAC TTT			1763
Val Phe Phe Gly Ala Gly Asn Tyr His Leu Val Asp Glu Asn Phe			
380	385	390	
GAG CCT TTA CCT GAT TAC TGG CTC TCT CTT CTG TTC AAG AAA CTG			1808
Glu Pro Leu Pro Asp Tyr Trp Leu Ser Leu Leu Phe Lys Lys Leu			
395	400	405	

GTA GGT CCC AGG GTG TTA CTG TCA AGA GTG AAA GGC CCA GAC AGG	1853
Val Gly Pro Arg Val Leu Leu Ser Arg Val Lys Gly Pro Asp Arg	
410 415 420	
AGC AAA CTC CGA GTG TAT CTC CAC TGC ACT AAC GTC TAT CAC CCA	1898
Ser Lys Leu Arg Val Tyr Leu His Cys Thr Asn Val Tyr His Pro	
425 430 435	
CGA TAT CAG GAA GGA GAT CTA ACT CTG TAT GTC CTG AAC CTC CAT	1943
Arg Tyr Gln Glu Gly Asp Leu Thr Leu Tyr Val Leu Asn Leu His	
440 445 450	
AAT GTC ACC AAG CAC TTG AAG GTA CCG CCT CCG TTG TTC AGG AAA	1988
Asn Val Thr Lys His Leu Lys Val Pro Pro Pro Leu Phe Arg Lys	
455 460 465	
CCA GTG GAT ACG TAC CTT CTG AAG CCT TCG GGG CCG GAT GGA TTA	2033
Pro Val Asp Thr Tyr Leu Leu Lys Pro Ser Gly Pro Asp Gly Leu	
470 475 480	
CTT TCC AAA TCT GTC CAA CTG AAC GGT CAA ATT CTG AAG ATG GTG	2078
Leu Ser Lys Ser Val Gln Leu Asn Gly Gln Ile Leu Lys Met Val	
485 490 495	
GAT GAG CAG ACC CTG CCA GCT TTG ACA GAA AAA CCT CTC CCC GCA	2123
Asp Glu Gln Thr Leu Pro Ala Leu Thr Glu Lys Pro Leu Pro Ala	
500 505 510	
GGA AGT GCA CTA AGC CTG CCT GCC TTT TCC TAT GGT TTT TTT GTC	2168
Gly Ser Ala Leu Ser Leu Pro Ala Phe Ser Tyr Gly Phe Phe Val	
515 520 525	
ATA AGA AAT GCC AAA ATC GCT GCT TGT ATA TGA AAA TAA AAG GCA	2213
Ile Arg Asn Ala Lys Ile Ala Ala Cys Ile	
530 535	
TAC GGT ACC CCT GAG ACA AAA GCC GAG GGG GGT GTT ATT CAT AAA	2258
ACA AAA CCC TAG TTT AGG AGG CCA CCT CCT TGC CGA GTT CCA GAG	2303
CTT CGG GAG GGT GGG GTA CAC TTC AGT ATT ACA TTC AGT GTG GTG	2348
TTC TCT CTA AGA AGA ATA CTG CAG GTG GTG ACA GTT AAT AGC ACT	2393
GTG	2396

## (2) INFORMATION FOR SEQ ID NO:46:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 385  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:46

CGGCCGCTGC TGCTGCTGTG GCTCTGGGGG CGGCTCCGTG CCCTGACCCA	50
AGGCACTCCG GCGGGGACCG CGCCGACCAA AGACGTGGTG GACTTGGAGT	100
TTTACACCAA GAGGCTATT CAAAGCGTGA GTCCCTCGTT CCTGTCCATC	150
ACCATGACG CCACTCTGGC CACCGACCCT CGGTTCCCTCA CCTTCCTGAG	200
CTCTCCACGG CTTCGAGCCC TGTCTAGAGG CTTATCTCCT GCGTACTTGA	250
GATTTGGCGG CACCAAGACT GACTTCCTTA TTTTGTATCC CAACAACGAA	300
CCCACCTCTG AAGAAAGAAG TTA CTGGCAA TCTCAAGACA ACAATGATAT	350



36

TTGCGGGTCT GACCGGGTCT CCGCTGACGT GTTGA

385

## (2) INFORMATION FOR SEQ ID NO:47:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 541  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: double  
(D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:47

AAATCAGGAC	ATATCCTTCA	CTTATTTGCC	TCTTGGTCAT	ATTGGAGGCA	50
TTTGTATTCA	TTTTTAATAA	CCCTCAAAAT	AGTGCATGCA	AAGTGCTAAG	100
CGTCATTTGC	CACATGGTGC	CATTAACTGT	CACCACCTGC	AGTGGTCTAC	150
TTAGAGAACA	CCGCACTGGA	TGTTAACT	GAAGCGCGTG	CCCCGCCCTC	200
CCGAGGCTCT	GGATCCAGCG	TTGAAGCTTG	CCCCGCCCTC	CCGAGGCTCT	250
GGATCCAGCA	CTGGAGCATG	CCCCGCCCTC	CCGAGGCTCT	GGAGCTTGCT	300
AAGGAGTCCG	CTCCCTACCG	CTGGGGTTTT	GCTTTATTCT	TATGAATGAC	350
ACCCCTGACC	GCITTCGTCT	CAGGGGTACT	GTAAATGCCTT	TTATTTTCAT	400
ATACAAGCTG	CGATTTTGGC	ATTTCTTATG	ACAAAAAACC	CATAGGAAAA	450
GGCGGGCAGC	CTTAGTGAGC	TTCTGCGGG	GAGAGGTTTT	TCTGTTAGAG	500
CTGGCANGGT	CTGCTCATCG	ACCATCTTCA	GGCCTCGTGC	C	541

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US00/03542

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(7) : C12N 15/56, 15/63, 1/21, 9/24, 15/11

US CL : 536/23.1, 23.2; 435/200, 325, 252.3, 320.1; 424/94.61

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 536/23.1, 23.2; 435/200, 325, 252.3, 320.1; 424/94.61

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EAST, MEDLINE, BIOSIS, CAPLUS, SCISEARCH, EMBASE, JAPIO, PATOWEP, PATOSWO search terms: heparanase, gene or sequence

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X - Y	US 5,362,641 A (FUKS et al.) 08 November 1994, see entire document.	21-25 ----- 1-20, 26-28
X, P	US 5,968,822 A (PECKER et al.) 19 October 1999, see entire document.	1-28
X - Y	WO 95/04158 A1 (THE UPJOHN COMPANY) 09 February 1995, see entire document.	21-25 ----- 1-24, 26-28
X, P	WO 99/11798 A1 (INSIGHT STRATEGY & MARKETING LTD.) 11 March 1999, see entire document.	1-28

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* "A"	Special categories of cited documents: document defining the general state of the art which is not considered to be of particular relevance	*T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*B"	earlier document published on or after the international filing date	*X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*O"	document referring to an oral disclosure, use, exhibition or other means	*A document member of the same patent family
*P"	document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

12 JUNE 2000

Date of mailing of the international search report

24 JUL 2000

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

RICHARD HUTSON

Telephone No. (703) 308-0196

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US00/03542

## Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
  
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
  
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/03542

### BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claim(s) 1-7, 19, 20 and 28, drawn to a nucleic acid encoding a polypeptide having heparanase activity.

Group II, claim(s) 8-18, drawn to antisense oligonucleotides of a polynucleotide which encodes a polypeptide having heparanase activity.

Group III, claim(s) 21-25, drawn to polypeptide having heparanase activity.

Group IV, claim(s) 26, drawn to a method of identifying a chromosome region harboring a heparanase gene.

Group V, claim(s) 27, drawn to a method of eliciting anti-heparanase antibodies in vivo.

The species listed above do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, the species lack the same or corresponding special technical features for the following reasons: The listed inventions share a technical relationship of a polypeptide having heparanase activity, but this does not constitute a special technical feature because Fuks et al. (Fuks et al. US Patent No: 5,362,641) teach a polypeptide having heparanase activity.